

THE ROLE OF ADAPTIVE STRESS RESPONSES IN HIV  
REPLICATION AND MACROPHAGE-MEDIATED NEUROTOXICITY

Stephanie A. Cross

A DISSERTATION

In

Neuroscience

Presented to the Faculties of the University of Pennsylvania

In

Partial Fulfillment of the Requirements for the

Degree of Doctor of Philosophy

2011

Supervisor of Dissertation

---

Dennis L. Kolson, M.D., Ph.D., Professor, Neurology

Graduate Group Chairperson

---

Rita Balice-Gordon, Ph.D., Professor of Neuroscience

Dissertation Committee

Peter B. Crino, M.D., Ph.D., Associate Professor of Neurology  
Virginia M. Lee, M.D., M.B.A., John H. Ware 3rd Professor in Alzheimer's Research  
Kelly L. Jordan-Sciutto, Ph.D., Associate Professor of Pathology  
Glenn Rall, Ph.D., Associate Professor

## **Dedication**

For my mom, Charlie.

## **Acknowledgments**

I have so many people that I need to thank and acknowledge. Graduate school is a challenging and difficult process and I never would have made it through, or had so many good times along the way, if it weren't for the help, support and friendship that so many people have shown me. To each and every person who has touched my life over the past 5 years, I sincerely thank you for the lessons you have taught me. I have learned so very much about life, but maybe most of all, have realized that I have so much more to learn.

First and foremost, I need thank Dennis for his mentorship, support, encouragement and enthusiasm. I never would have finished this PhD if he didn't decide to take a chance on me, and I will forever be thankful that I landed in his lab. I am so incredibly grateful for having the opportunity to learn from him and I never believed that I would find such a great mentor. I am constantly amazed and inspired by his enthusiasm for medicine and science – the MD/PhD training is a long one, and thanks to Dennis, I now recognize that holding on to the joy of scientific discovery and desire to improve medicine is necessary for anyone to finish the program and become a physician-scientist. I also hope to one day achieve a similar level of calmness, perspective and professionalism that Dennis exemplifies so readily.

I also need to acknowledge each and every person who has helped me with my project, the day-to-day problems and small (and large) issues that comprise a typical graduate student's training. To Pat and Lorraine, thank you for being so awesome. You keep the lab running smoothly, have been invaluable in supporting my science and helped me navigate some tough times. I couldn't have asked for better colleagues. Thanks to everyone in the Crino Lab, Jordan-Sciutto Lab, Soldan Lab and Kolson Lab – you helped shape my graduate training in so many ways, thanks for everything.

To all of my friends – thank you for your support and friendship. It's been a crazy few years for me, but I have so many fond memories and great times to look back on. To the MD/PhDs, matriculating class of 2004, I honestly believe that we are the best class ever and I never would have wanted a different group to go through this process with. I need to thank Tony Chi, Abby Olsen, Ana Cristancho, Ksenia Orlova, Renuka Nayak, Erica Dwyer, Marta Rowh, Alex Chavez and Amy Tsou especially. You are a formidable group of people and I know you will all do well with whatever path you chose to take in life. Natalia, MV and Sam – I consider you some of the best friends I have made at Penn and I am thankful that I have had the opportunity to get to know you. You guys have made lab so much fun, and I can never describe how much that has meant to me over the years. And to Chelain, Tedi, Alaena, Jen and Andrew – you guys are keepers for life. Thanks for sticking with me, I know I've had more ups and downs than anyone would have ever predicted. Your constant friendship and support has meant so much to me over the years, I really don't know what I have done without you.

I never would have ever made it to graduate school if not for the help and support of my family. My mom will never get to see this thesis or my graduation, but I know that I never would have been able to do this without her 28 years of support. I miss her immensely, but the lessons she taught me have served me well and I hope that they continue to do so. I have never known a stronger, more intelligent woman and I can only hope to live up to the potential of which she believed me capable. To my Dad and my sister – you two are now my rock and I love you both. Thank you for your support, sometimes it felt like the only thing that was holding me together.

I would like to thank the MD/PhD program, especially Skip Brass, and my thesis committee members. Your support and encouragement made all the difference and without it I never would have brushed myself off and kept on trying. To each and every one of you, thank you.



Finally, I need to acknowledge Maggie Krall. Maggie is undeniably the heart and soul of the MD/PhD program here at Penn and has been supportive from an academic/coordinating perspective for so many others in addition to myself. However, Maggie is also one of the most amazing people that I have ever met in my life. I don't have the words to describe how truly wonderful she is or what her support has meant to me. She inspires me to be a better person, has spent many hours helping me talk through issues that felt too dreadful and immense to handle alone, and helped keep me sane during a time in my life when everything, including myself, was in total disarray. She has seen me at my worst, and I could never have achieved my best without her help. I can never thank Maggie enough for her support, ever.

## ABSTRACT

### THE ROLE OF ADAPTIVE STRESS RESPONSES IN HIV REPLICATION AND MACROPHAGE-MEDIATED NEUROTOXICITY

Stephanie A. Cross

Dennis L. Kolson, M.D., Ph.D.

Despite antiretroviral therapy (ART), HIV infection promotes cognitive dysfunction and neurodegeneration through persistent inflammation and neurotoxin release from infected and/or activated macrophages. Inflammation and immune activation within both the central nervous system (CNS) and periphery correlate with disease progression and morbidity in ART-treated individuals. Accordingly, drugs targeting these pathological processes are needed for effective, adjunctive therapy. Using our *in vitro* model of HIV-mediated neurotoxicity, in which HIV infected monocyte-derived macrophages (HIV/MDM) release excitatory neurotoxins, we demonstrate that HIV infection dysregulates adaptive stress responses, including the antioxidant response and the unfolded protein response (UPR). HIV infected macrophages have dramatic reductions in heme oxygenase-1 (HO-1) levels. Activation of the antioxidant response attenuates HIV replication and restoration of HO-1 expression, specifically, reduces neurotoxin release from HIV/MDM, even with robust HIV replication. We propose that dysregulation of the antioxidant response during HIV infection drives macrophage-mediated neurotoxicity and that pharmacological inducers of the antioxidant response could serve as adjunctive neuroprotectants and HIV disease modifiers in ART-treated individuals. Additionally, we found that HIV infection activates the UPR in macrophages, increasing phosphorylated eIF2 $\alpha$  in our *in vitro* system and macrophagic BiP in HAND frontal cortex. Pharmacological induction of the UPR, which attenuates viral replication, enhances macrophage-mediated neurotoxicity. Therefore, processes that induce the UPR in macrophages may enhance neurotoxin production and contribute to pathological processes

underlying HAND. Understanding how HIV infection affects adaptive stress responses and neurotoxin production pathways in the macrophage will improve our ability to develop effective adjunctive therapies for the neurological consequences of HIV infection.

## **Table of Contents**

List of Tables .....	xi
List of Figures .....	xii
List of Abbreviations .....	xiv
 <b>CHAPTER 1: INTRODUCTION</b> .....	 1
Figure 1-1 .....	7
 <b>CHAPTER 2: NEUROAIDS AS AN INFLAMMATORY DISORDER</b> .....	 9
Abstract .....	10
Introduction .....	11
2.1 - Pathological Characteristics of NeuroAIDS .....	11
Table 2-1 .....	17
2.2 – Biology of HIV Infection and Invasion of the Brain .....	18
Figure 2-1 .....	20
2.3 – Mechanisms of HIV-Induced Neurodegeneration: Roles for Chemokines, Chemokine Receptors, and Inflammation .....	22
2.4 – Inflammatory Markers in NeuroAIDS and Other Neuroinflammatory Diseases .....	26
2.5 – Therapeutic Considerations .....	32
Table 2-2 .....	35
 <b>CHAPTER 3: MATERIALS AND METHODS</b> .....	 38
Table 3-1 .....	46
Table 3-2 .....	47

## **CHAPTER 4: THE ANTIOXIDANT RESPONSE ATTENUATES HIV REPLICATION AND MACROPHAGE-MEDIATED NEUROTOXICITY ..... 48**

Abstract	49
Introduction	50
Results	53
Figure 4-1	54
Figure 4-2	55
Table 4-1	56
Figure 4-3	57
Figure 4-4	59
Figure 4-5	61
Figure 4-6	64
Figure 4-7	66
Figure 4-8	68
Figure 4-9	70
Discussion	71
Work In Progress	76
Figure 4-10	77
Figure 4-11	79
Figure 4-12	81
Figure 4-13	83

## **CHAPTER 5: THE UNFOLDED PROTEIN RESPONSE ATTENUATES HIV REPLICATION AND ENHANCES MACROPHAGE-MEDIATED NEUROTOXICITY ..... 84**

Abstract	85
Introduction	87
Figure 5-1	89

Results .....	98
Figure 5-2 .....	101
Figure 5-3 .....	103
Figure 5-4 .....	104
Figure 5-5 .....	107
Figure 5-6 .....	109
Figure 5-7 .....	112
Figure 5-8 .....	113
Figure 5-9 .....	115
Discussion .....	116
 <b>CHAPTER 6: DISCUSSION</b> .....	 122
Summary of Work .....	123
Implications .....	126
 <b>BIBLIOGRAPHY</b> .....	 130

### **List of Tables**

<b>Table 2.1.</b>	Changes in the neuropathology of HAND in the era of HAART
<b>Table 2.2.</b>	Therapeutics under consideration for HAND
<b>Table 3.1.</b>	Summary of case data for autopsy tissue obtained from the National Neuro-AIDS Tissue Consortium brain bank for BiP and HLA-DR staining.
<b>Table 3.2.</b>	Summary of case data for autopsy tissue obtained from the National Neuro-AIDS Tissue Consortium brain bank for phosphorylated eIF2 $\alpha$ and HAM56 staining.
<b>Table 4.1.</b>	Reduction in RT activity, relative to vehicle control, with increasing concentrations of DMF and MMF among individual donors.

## **List of Figures**

- Figure 1.1** The signaling pathways comprising the antioxidant response and the Unfolded Protein Response (UPR).
- Figure 2.1** Role for inflammation via cytokines and chemokines in HIV neuroinvasion and neurodegeneration.
- Figure 4.1** Dimethyl fumarate and monomethyl fumarate attenuate HIV replication in human MDM.
- Figure 4.2** Dimethyl fumarate attenuates 89.6 HIV replication in human MDM.
- Figure 4.3** Dimethyl fumarate and efavirenz have additive effects on attenuation of HIV replication and neurotoxin production in human MDM.
- Figure 4.4** DMF and MMF reduce HIV/MDM mediated neurotoxicity.
- Figure 4.5** DMF inhibits NF- $\kappa$ B nuclear translocation, DNA binding and TNF $\alpha$  production in human MDM.
- Figure 4.6** DMF restores the imbalance in the antioxidant response caused by HIV infection.
- Figure 4.7** HO-1 does not mediate the attenuation of HIV replication or NF- $\kappa$ B signaling induced by DMF.
- Figure 4.8** HO-1 induction reduces neurotoxin production in HIV/MDM without affecting HIV replication.
- Figure 4.9** DMF and MMF reduce CCL2 induced chemotaxis in human monocytes.
- Figure 4.10** tBHQ, an inducer of the antioxidant response, attenuates HIV replication and macrophage-mediated neurotoxicity.
- Figure 4.11** Hemin, an inducer of HO-1 and the antioxidant response, attenuates macrophage-mediated neurotoxicity without altering HIV replication.
- Figure 4.12** DMF decreases the cell surface expression of CCR5 and CXCR4 on human monocytes.
- Figure 4.13** DMF, but not efavirenz, added to an established HIV infection decreases macrophage-mediated neurotoxicity without altering HIV replication.
- Figure 5.1** The signaling pathways comprising the Unfolded Protein Response (UPR).
- Figure 5.2** HIV infection induces phosphorylation of eIF2 $\alpha$  in human MDM.
- Figure 5.3** HIV infection increases levels of macrophage-specific BiP in the human frontal cortex of HAND cases.



- Figure 5.4** HIV infection does not alter levels of macrophage specific phosphorylated-eIF2 $\alpha$  in the human frontal cortex of HAND cases.
- Figure 5.5** Thapsigargin attenuates HIV replication and enhances neurotoxin production in human MDM.
- Figure 5.6** Pharmacological induction of phosphorylated eIF2 $\alpha$  and disruption of Ca<sup>2+</sup> stores in the endoplasmic reticulum enhances neurotoxin release in human MDM.
- Figure 5.7** DMF and MMF activate the antioxidant response and decrease levels of phosphorylated eIF2 $\alpha$  in HIV-infected human macrophages.
- Figure 5.8** MMF activates the antioxidant response and decreases levels of Grp94 and BiP in uninfected human macrophages.
- Figure 5.9** Levels of phosphorylated eIF2 $\alpha$  do not correlate with HIV replication or neurotoxin production in HIV-infected human macrophages.

### List of Abbreviations

<b>ARE</b>	antioxidant response element
<b>ART</b>	antiretroviral therapy
<b>BBB</b>	blood-brain barrier
<b>CCL2</b>	chemokine (C-C motif) ligand 2
<b>CLT</b>	clotrimazole
<b>CoPP</b>	cobalt (III) protoporphyrin IX chloride
<b>DMF</b>	dimethyl fumarate
<b>DPI</b>	day post infection
<b>eIF</b>	eukaryotic initiation factor
<b>EFZ</b>	efavirenz
<b>ER</b>	endoplasmic reticulum
<b>GPX1</b>	glutathione peroxidase 1
<b>HAND</b>	HIV-associated neurocognitive disorders
<b>HO-1</b>	Heme oxygenase-1
<b>HPI</b>	hour post infection
<b>IL</b>	interleukin
<b>IFN</b>	interferon
<b>MAP2</b>	microtubule-associated protein 2
<b>MDM</b>	monocyte-derived macrophages
<b>MMF</b>	monomethyl fumarate
<b>NF-<math>\kappa</math>B</b>	nuclear factor kappa-light-chain-enhancer of activated B cells
<b>NQO1</b>	NAD(P)H quinone oxidoreductase 1
<b>Nrf2</b>	nuclear factor erythroid 2-related factor 2
<b>PBMC</b>	peripheral blood mononuclear cell
<b>ROS</b>	reactive oxygen species
<b>RT</b>	reverse transcriptase
<b>SnMP</b>	tin (IV) mesoporphyrin IX dichloride
<b>tBHQ</b>	<i>tert</i> -butylhydroquinone
<b>TNF<math>\alpha</math></b>	tumor necrosis factor- $\alpha$
<b>UPR</b>	unfolded protein response

## **CHAPTER 1**

### **INTRODUCTION**

At the end of 2009, an estimated 30.8 million adults and 2.5 million children were living with HIV globally (WHO, 2010). Current antiretroviral therapy (ART) regimens for HIV infection have greatly improved virological control and clinical outcomes in infected individuals. However, despite the efficacy of ART, the prevalence of HIV-associated neurocognitive disorders (HAND) has persisted, with prevalence and associated morbidity estimated at up to 50% (McArthur et al., 2003; Robertson et al., 2007; Sacktor et al., 2002). While ART will remain the mainstay of HIV therapy, there is a need for adjunctive neuroprotective therapies that address the pathological processes persisting in ART-treated individuals.

HIV-1 infection of the central nervous system (CNS) can result in cognitive, motor, and behavioral abnormalities, collectively known as HIV-associated neurocognitive disorders (HAND) (Antinori et al., 2007; McArthur et al., 2010). Pathological processes in both the brain and periphery, involving multiple cell types, contribute to the neurological complications of AIDS. Early in the course of infection, HIV traffics into the brain via infected monocytes and lymphocytes (Dunfee et al., 2006). Despite ART, HIV persists in the CNS in parenchymal microglia and perivascular macrophage reservoirs (Ho et al., 1985; Koenig et al., 1986; Petito et al., 1986). And because HIV cannot infect neurons, HIV-induced neuronal damage is mediated indirectly by the release of neurotoxins from infected and/or activated macrophages, microglia and astrocytes. *In vitro* and *in vivo* studies have identified viral proteins, pro-inflammatory cytokines, interferons, excitatory amino acids, phospholipids, and reactive oxygen species as neurotoxic factors produced by macrophages/microglia following HIV infection (Boven et al., 1999; Brenneman et al., 1988; Brew et al., 1995; Gelbard et al., 1993; Gelbard et al., 1994; Gendelman et al., 1998; Genis et al., 1992; Jiang et al., 2001; Maragos et al., 2003; Scorziello et al., 1998; Song et al., 2003; Wesselingh et al., 1993). Furthermore, the severity of pre-mortem HAND correlates with increased numbers of microglia and macrophages in the CNS, supporting the hypothesis that these cell types are principal mediators of HIV-induced neurological impairment (Anthony et al., 2005; Glass et al., 1995; Petito et al., 1986).

The persistence of HAND in individuals effectively controlled for systemic viral replication is incompletely explained, although recent evidence suggests that prolonged inflammation in both the CNS and periphery may be responsible (Ancuta et al., 2008; Brenchley et al., 2006b; Eden et al., 2007). Multiple pro-inflammatory cytokines, including interleukin (IL)-1 $\beta$ , tumor necrosis factor-alpha (TNF- $\alpha$ ), and IL-6 are elevated in the CNS and/or CSF of patients with HAND (Achim et al., 1993; Foli et al., 1997; Oster et al., 1987; Perrella et al., 1992). The pro-inflammatory environment within the CNS is a result of cytokine release from immune activated astrocytes (Kramer-Hammerle et al., 2005; Sabri et al., 2003) and monocytes/macrophages stimulated by direct viral infection, shed viral proteins or proinflammatory mediators (Rappaport et al., 1999; Sundar et al., 1991). Inflammation can alter the permeability of the blood-brain barrier, enhance entry of infected monocytes into the CNS, and drive peripheral processes that contribute to the neurological complications of HIV. Accordingly, drugs targeting inflammatory-mediated processes in the CNS and systemic compartments are needed for effective, adjunctive therapy in HAND.

Numerous physiological and pathological stimuli can initiate and/or propagate the inflammatory response. Of the many relevant pathways, this work focuses on the interactions between oxidative stress, endoplasmic reticulum (ER) stress and inflammation. The ER, as a protein folding compartment and dynamic calcium store, is primed to sense cellular stress and initiate adaptive stress responses, including the unfolded protein response (UPR). Intracellular calcium signals and free radicals, such as reactive oxygen species (ROS) and nitric oxide (NO), act as messengers in coordinating several of the adaptive stress responses. Calcium release from the ER, during states of stress, can induce the generation and release of ROS from the mitochondria. This in turn activates the antioxidant response, inflammatory pathways and with sustained activation, cellular apoptosis. Furthermore, ROS can feedback onto the ER and target calcium channels and resident-ER chaperones to exacerbate ER calcium release and ER stress. As a consequence, misfolded proteins accumulate in the ER, induce the UPR, and activate signaling cascades to promote the inflammatory response, antioxidant response, apoptosis and other

cellular adaptive stress responses (Ron and Walter, 2007; Zhang and Kaufman, 2008). The effect of HIV infection on the antioxidant response, the UPR and the interactions between these adaptive stress responses and inflammation has not yet been explored.

The antioxidant response maintains redox balance and counteracts oxidative damage by inducing the transcriptional upregulation of proteins that are involved in detoxification of reactive oxygen species (ROS). These genes have a common promoter element, the antioxidant response element (ARE), and are regulated by nuclear factor erythroid 2-related factor 2 (Nrf2) (Figure 1.1). Following exposure to ROS or electrophiles, Nrf2 translocates to the nucleus to drive expression of numerous genes, including heme oxygenase-1 (HO-1), NAD(P)H quinone oxidoreductase 1 (NQO1), glutathione peroxidase 1 (GPX1), and genes responsible for the synthesis of glutathione, the principal antioxidant produced by the cell. Recent studies have demonstrated that viral infection can affect the cellular oxidation state and induce the activation of the antioxidant response (Chen et al., 2011; Schaedler et al., 2010). HIV infected patients have evidence of increased ROS production and depressed levels of glutathione (Dworkin et al., 1986), suggesting that the antioxidant response is altered following HIV infection.

We hypothesized that HIV infection of macrophages would result in suppression of the antioxidant response and that induction of the antioxidant response would attenuate HIV replication. Furthermore, restoration of the cellular redox state in HIV-infected macrophages could dampen inflammatory processes if they are driven by virus-induced generation of excessive ROS. The studies presented in this body of work begin to characterize the consequences of HIV infection on the antioxidant response and the link between the dysregulation of the antioxidant response and macrophage-mediated neurotoxicity. Most importantly, we demonstrate that HIV infection dramatically reduces levels of HO-1 in macrophages and that restoration of HO-1 decreases macrophage mediated neurotoxicity. We also show that induction of the antioxidant response attenuates NF- $\kappa$ B and TNF $\alpha$  signaling, a major mediator of inflammation and immune

activation. Our results provide evidence that therapeutics that ameliorate the dysregulation of the antioxidant response following HIV-infection should be considered as adjunctives to ART for the treatment and prevention of HAND.


We also examined the role of HIV-infection of macrophages on activation of the UPR and the interaction between activation of the UPR and the antioxidant response. Virus replication and assembly can strain the capacity of the endoplasmic reticulum (ER) and result in the accumulation of misfolded proteins and induction of the UPR. The UPR is designed to eliminate misfolded proteins and promote cellular recovery by attenuating translation and upregulating the expression of chaperones, degradation factors, and regulators of metabolic and redox states. Activation of the signaling pathways comprising the UPR can also induce the transcription of inflammatory genes via activation of NF- $\kappa$ B pathways, among others.

The UPR is a quality control mechanism that can be activated during physiological stress (e.g., glucose deprivation, oxidative stress) and viral infection (Figure 1.1). The UPR attempts to eliminate excessive misfolded proteins in the ER through two mechanisms; 1) attenuation of protein translation, in order to reduce the flux of proteins entering into the ER and 2) induction of chaperone proteins and degradation factors to refold and/or eliminate misfolded proteins. Three proteins have been identified as sensors of ER stress: protein kinase RNA-like ER kinase (PERK), activating transcription factor 6 (ATF6), and inositol-requiring enzyme 1 (IRE1). Under normal conditions, the ER chaperone immunoglobulin heavy-chain-binding protein (BiP) is bound to the sensor domain of PERK, ATF6 and IRE1 in the ER lumen. When misfolded proteins accumulate in the ER, BiP is sequestered away from these sensors to bind to misfolded proteins, resulting in activation of PERK, ATF6 and IRE1 (Bertolotti et al., 2000). Activation of PERK initiates a signaling cascade that results in the inhibition of new protein translation, mediated by phosphorylated eIF2 $\alpha$ , and the induction of genes important for cellular recovery from ER stress, as mediated by ATF4 transcriptional activity. ATF6 activates genes carrying ER stress response

elements (ERSEs) in their transcriptional promoters (Mori, 2000). ERSEs are found in genes encoding for cellular chaperones including BiP, protein-disulfide isomerase (PDI) and calreticulin (Harding et al., 2003). Phosphorylation of IRE1 results in the alternative splicing of X box binding protein 1 (XBP-1) mRNA, which encodes an active transcription factor capable of inducing genes regulated by the ERSE (Lee et al., 2003). In addition, sXBP-1 can also activate the transcription of genes containing UPR elements (UPRE), thereby enhancing the capacity of ER-stressed cells to degrade irrevocably misfolded proteins via ER-associated protein degradation (ERAD) (Hosokawa et al., 2001; Yoshida et al., 2003). Chronic activation of the UPR, which occurs when the cell cannot manage or recover from ER stress, results in the activation of signaling pathways that will commit the cell to apoptosis (Ferri and Kroemer, 2001; Oyadomari et al., 2002a).

In some cases, viral infection induces ER stress and activates components of the three branches of the UPR. Enveloped viruses utilize the ER as the primary site of envelope glycoprotein biogenesis and several viruses undergo genomic replication and particle assembly in the ER compartment. Increasing the burden on the ER, especially at times of high viral production, could increase the accumulation of misfolded proteins in the ER lumen and activate the UPR. However, while activation of the UPR is essential for host cell survival during viral infection, some of the consequences of UPR activation, such as inhibition of protein translation and enhanced ERAD, would be detrimental to viral replication. Consequently, many viruses that induce the UPR have evolved mechanisms to regulate pathways of the UPR in order to promote efficient viral replication. While UPR activation has been reported following cellular infection by several different viruses, the effect of HIV infection on activation of the UPR and the role of the UPR in modulating HIV replication has not yet been investigated.





**Figure 1.1 The signaling pathways comprising the antioxidant response and the Unfolded Protein Response (UPR).** The antioxidant response maintains the redox state of the cell and counteracts oxidative damage through induction of proteins that are involved in detoxification of reactive oxygen species (ROS). The transcription of these genes is regulated by Nrf2-dependent transcription of the antioxidant response element (ARE). During states of low oxidative stress, Nrf2 is kept transcriptionally inactive by Keap1, which sequesters Nrf2 in the cytoplasmic compartment. Following exposure to ROS or electrophiles, Keap1 is degraded by the proteasome and Nrf2 translocates to the nucleus to drive expression of HO-1, NQO1, and glutathione peroxidase 1 (GPX1), among others. Accumulation of misfolded protein in the ER causes ER stress and results in the activation of the UPR, an adaptive stress pathway. PERK, IRE1 and ATF6 have sensor domains that monitor levels of misfolded protein in the ER lumen. Following activation, each of these pathways initiates a signaling cascade that results in the transcriptional upregulation of genes that will help the cell reduce, manage or recover from ER stress. Phosphorylation of eIF2α by activated PERK results in the attenuation of all cap-dependent protein translation, in order to reduce the incoming burden of new proteins into the ER. Increased levels of phosphorylated eIF2α also result in the translational upregulation of ATF4, a transcription factor that increases amino acid response element (AARE)-regulated genes in order to modulate cellular metabolism and redox state during cellular recovery from ER stress. ATF4 also increases levels of GADD34, which restores protein translation by mediating eIF2α dephosphorylation as part of a negative feedback loop. In addition to PERK, three other kinases (PKR, GCN2 and HRI) can phosphorylate eIF2α and initiate the UPR, although instead of misfolded protein levels, they respond to double stranded RNA, amino acid limitation and heme levels, respectively. ATF6, upon activation, translocates to the Golgi where it is cleaved into an active form, which also functions as a transcriptional regulator of ER stress-element (ERSE)-regulated genes. ERSEs regulate genes encoding for a variety of cellular chaperones, which promote the proper folding of misfolded proteins in the ER, including BiP and GRP94. Activation of IRE1 results in its auto-phosphorylation and the subsequent activation, by a splicing event, of XBP1 mRNA. Spliced XBP1 (sXBP1) encodes a transcription factor capable of upregulating genes with ERSEs or UPR elements (UPREs). Transcriptional upregulation of UPRE-regulated genes results in increased levels of EDEM, which enhances the degradation of misfolded proteins in the ER by ER-associated degradation (ERAD). If the UPR is unable to manage the stress to the ER, the cell will undergo CHOP-mediated apoptosis.

We hypothesized that HIV infection would induce the UPR during times of high viral replication, when viral protein translation, glycoprotein biogenesis and viral particle assembly would be highest. We found that HIV infection increases levels of phosphorylated eIF2 $\alpha$  in our *in vitro* model system and that macrophagic BiP was increased in the macrophages of HAND frontal cortex. Interestingly, pharmacological induction of the UPR, which attenuates viral replication, is associated with increased macrophage-mediated neurotoxicity. This finding has important implications for the development of adjunctive therapies for HAND. Therapeutics or processes that induce the UPR in macrophages, regardless of the effect on HIV replication, could enhance neurotoxin production and contribute to the pathological processes underlying HAND.

HIV infection results in the alteration of several adaptive stress pathways in the macrophage. Suppression of the antioxidant response and induction of the UPR both associate with high levels of viral replication and neurotoxin production. And while pharmacological induction of either stress pathway attenuates HIV replication, only induction of the antioxidant response results in decreased macrophage-mediated neurotoxicity. Therefore, activation of the UPR by inflammation, oxidative stress or ART is predicted to enhance neurotoxin production in macrophages, regardless of the level of HIV replication. Understanding how HIV infection affects adaptive stress responses and neurotoxin production pathways in the macrophage will continue to improve our ability to develop effective adjunctive therapies for HAND.

## CHAPTER 2

### NEUROAIDS AS AN INFLAMMATORY DISORDER<sup>1</sup>

---

<sup>1</sup> This work was originally published in *The Neurology of AIDS, 3<sup>rd</sup> Edition*. Cook DR\*, Cross SA\*,

## **Abstract**

The pathological basis for the neurological complications of AIDS involves complex interactions between multiple cell types within both the brain and the periphery. This review focuses on the role for inflammation in the development of HIV-associated neurocognitive disorders (HAND). Specifically, evidence for inflammation in HAND pathogenesis in the pre-antiretroviral therapy (ART) and post-ART eras is discussed. The biology of HIV infection and subsequent CNS invasion is emphasized, with particular focus on chemokines and chemokine receptors. Chronic neuroinflammation has also been implicated in other infectious and non-infectious CNS disorders, of which multiple sclerosis (MS), Alzheimer's disease (AD), Parkinson's disease (PD) and HTLV-1-associated myelopathy/tropical spastic paraparesis (HAM/TSP) are reviewed. The status of current adjunctive therapies for HAND and their relationship to such neuroinflammatory disorders is also included. Understanding common pathways of neuroinflammation will lead to the identification of biomarkers for classification, diagnosis, and clinical prognosis, as well as to the development of novel treatment modalities for HAND and other neuroinflammatory disorders of the CNS.

## **Introduction**

Advances in antiretroviral therapies for HIV infection have improved virological control and greatly increased the life expectancy of infected individuals. However, despite the availability of these potent antiviral drugs, the prevalence of HIV-associated neurocognitive disorders (HAND) has persisted and even increased (McArthur et al., 2003; Robertson et al., 2007; Sacktor et al., 2002), emphasizing the need for effective adjunctive neuroprotective therapies. Inflammation in the brain and in the periphery is a driving force in the neuropathogenesis of HAND, MS, HAM/TSP, and probably also AD and PD. As these conditions share common features in their pathogenic processes, identifying common mechanisms and developing targeted treatment modalities will not only benefit HAND patients, but likely also be effective in other neuroinflammatory disorders.

## **2.1 - Pathological Characteristics of NeuroAIDS**

### *Clinical Characteristics of HIV-Associated Neurocognitive Disorders (HAND)*

According to recent estimates, over 33 million people are currently living with HIV-1 worldwide (WHO, 2009). HIV-1 infection has devastating consequences for the immune system, resulting in immunodeficiency marked by profound CD4<sup>+</sup> T-cell depletion. In addition, neurologic disorders involving the CNS and the peripheral nervous system (PNS) affect between 40–70% of HIV-positive individuals at some point during the course of infection (McArthur et al., 2005). Although opportunistic infections of the CNS and PNS associated with HIV-induced immunodeficiency have become far less common because of the availability of antiretroviral therapy (ART) (Habata et al., 1999; Mamidi et al., 2002; Roulet, 1999), the prevalence of primary HIV-induced neurological disorders has increased (Antinori et al., 2007). Conditions directly induced by HIV-1 include peripheral neuropathies, vacuolar myelopathies, and HIV-associated neurocognitive disorders (HAND) (Antinori et al., 2007; Childs et al., 1999; McArthur et al., 2005; McArthur et al., 2003).

HAND are comprised of three conditions of increasingly severe cognitive impairment and interference with activities of daily living: (1) HIV-associated asymptomatic neurocognitive impairment (ANI), (2) HIV-associated mild neurocognitive disorder (MND), and (3) HIV-associated dementia (HAD) (Antinori et al., 2007; Grant, 2008). The diagnosis of ANI, the least severe of the HAND conditions, describes individuals who have mild cognitive impairment, revealed by formal neuropsychological testing, that does not affect day-to-day functioning and does not meet criteria for delirium or dementia. In the more severe MND, increased cognitive impairment interferes with daily functioning as determined by self-reporting or by observation of others. Again, individuals with MND do not meet the criteria for delirium or dementia. HAD, the most severe form of HAND, is typically diagnosed during end-stage HIV infection, primarily in patients with low CD4+ T-cell counts. In HAD, cognitive impairment is associated with marked interference with day-to-day functioning and HAD is considered a significant independent risk factor for death due to AIDS (Liner et al., 2008). Since the inception of ART, the incidence of HAD, defined as the percentage of new HAD cases diagnosed in a given year, has decreased (McArthur et al., 2004; Nath et al., 2008; Sacktor et al., 2002). However, the prevalence of HAD, defined as the overall number of HAD cases, is rising owing to the increased life span of HIV patients (McArthur et al., 2003; Robertson et al., 2007; Sacktor et al., 2002). There is also evidence of ongoing neurodegeneration that may manifest as ANI or MND in patients without evidence of active HIV disease, contributing to the continued prevalence of HAND in the ART era. Moreover, the onset of HAND during ART-controlled clinical latency underscores the need for adjunctive neuroprotective therapies, because no current therapies would be expected to improve neurologic outcomes in the absence of viral replication (Antinori et al., 2007; Brew, 2004; Sacktor et al., 2002).

#### *Pathology of HAND: Evidence for Neuroinflammation*

HIV-1 enters the brain early in the course of infection via infected macrophages and lymphocytes (Ho et al., 1985; Koenig et al., 1986; Petito et al., 1986). Subsequently, the virus persists in the

CNS primarily in perivascular macrophages and microglia and increased numbers of macrophages and microglia have been found to correlate with the severity of HAND (Glass et al., 1995). In general, intrathecal replication of HIV-1 is controlled by CD8+ T cells (McCrossan et al., 2006; Sadagopal et al., 2008), and cerebral spinal fluid (CSF) viral load has been found to correlate with both viral load in the brain and the degree of cognitive dysfunction in HAND (Brew et al., 1997; Ellis et al., 2000; Ellis et al., 1997; Ellis et al., 2002; McArthur et al., 1997). Importantly, in addition to the initial neuroinvasion and infection of macrophages and microglia, it is believed that HIV infection and immune activation in the periphery and ongoing neuroinvasion of activated monocytes play a role in the development of HAND (Banks et al., 2006; Gartner, 2000).

The pathological hallmarks of HIV infection in the brain in the pre-ART era, collectively termed HIV encephalitis (HIVE), include monocyte infiltration and accumulation of perivascular monocyte-derived macrophages (MDM), formation of microglial nodules and multinucleated giant cells (syncytia) due to HIV-driven fusion of MDM/microglia, widespread reactive microgliosis and astrogliosis, and myelin pallor indicative of oligodendrocyte damage (Adle-Biassette et al., 1999; Gendelman et al., 1994; Lawrence and Major, 2002; Masliah et al., 1997; Petito et al., 1986; Wiley and Achim, 1994). Since the initiation of widespread ART (1996/7), however, at least one study suggests the presence of “burnt out” HIVE in some individuals dying with AIDS (see below) (Gray et al., 2003). MDM and microglia are the primary CD4+ cells in the CNS and the major sources of productive HIV infection in the brain (Gonzalez-Scarano and Martin-Garcia, 2005; Kaul et al., 2001; Kolson and Gonzalez-Scarano, 2000; McArthur et al., 2003) and clinical disease severity correlates more strongly with the amount of monocyte infiltration and MDM/microglia activation than with the quantity of infected cells or viral load (Adle-Biassette et al., 1999; Glass et al., 1995), suggesting that MDM/microglia play a predominant role in the neuroinflammation and neurotoxicity seen in HAND. Immune activation of MDM/microglia is demonstrated by expression of CD14 (lipopolysaccharide receptor), CD16, CD68, and MHC class II in vivo (Anderson et al.,

2002; Bell, 2004; Fischer-Smith et al., 2004; Swindells et al., 1999). Furthermore, CSF markers of immune activation and inflammation are commonly detected in individuals with HAND. These markers include CCL2 (monocyte chemoattractant protein-1, MCP-1) (Chang et al., 2004; Conant et al., 1998),  $\beta$ 2 microglobulin (Brew et al., 1992; Brew et al., 1996; Enting et al., 2000; McArthur et al., 1992), quinolinic acid (Achim et al., 1993; Brew et al., 1995; Heyes et al., 1991; Heyes et al., 2001), arachidonic acid metabolites (Genis et al., 1992; Griffin et al., 1994), oxidative stress markers (Haughey et al., 2004; Schifitto et al., 2009a), and platelet activating factor (PAF) (Gelbard et al., 1994).

Although most studies demonstrate that neurons are not infected by HIV, neuronal loss is common in HAND and post-mortem studies of HAND patients have revealed morphological changes in neurons including loss of synaptic density, dendritic simplification, and vacuolization (Masliah et al., 1997; Petito et al., 1986). This neuronal damage induced by HIV infection affects multiple regions of the brain and several neuronal subtypes. HIV antigen is commonly detected in the thalamus, basal ganglia, and central white matter and neuronal damage and loss has been reported in the frontal cortex, cerebellum, putamen, and substantia nigra, although the distribution of HIV antigen might be altered in individuals receiving ART (Everall et al., 1991; Ketzler et al., 1990).

#### *Changes in Neuropathology of HAND in the Era of ART*

The most effective current therapy for HAND is treatment of the underlying HIV infection with ART. Neuropsychological performance is improved in AIDS patients treated with ART and the CNS penetration of antiretroviral drugs directly correlates with decreased CSF viral loads and improved neurocognitive performance (Letendre et al., 2008). However, some recent studies suggest that ART drugs might demonstrate CNS neurotoxicity in treated patients, with associated poorer neurocognitive performance (Marra et al., 2009; Schweinsburg et al., 2005). Unfortunately, although ART can improve cognition, it does not fully eradicate impairments. In addition, patients



who have had ART exposure before developing HAND do not appear to respond as well to a change in their ART regimen, indicating the presence of drug-resistant viruses in the CNS or the emergence of a burn-out phase of the disease (Brew et al., 2007).

The era of ART has changed the neuropathology of HIV-1 infection (Table 2.1) (Anthony et al., 2005; Boisse et al., 2008; Brew, 2004). Before the introduction of ART, neuroinflammation was frequently observed in HIV-infected patients and usually increased throughout the progression of disease from the asymptomatic stage to AIDS and HAD. Inflammation is less severe during ART, but it appears to be persistent within the macrophage/microglial populations (Gray et al., 2003). Furthermore, ART seems to have limited the severity of pathological changes characteristic of HIVE. As described by Gray and colleagues (Gray et al., 2003), the persistent pathological findings in ART-experienced individuals include neuronal loss with apoptosis, astrocytosis, myelin pallor, and at least a few activated microglia and perivascular macrophages. Distinctly absent are multinucleated giant cells and microglial nodules.

Although for the most part ART has limited the persistent infiltration of HIV-infected lymphocytes into the CNS (Anthony and Bell, 2008), it should be noted that an exception to this occurs during neurologic immune reconstitution inflammatory syndrome (neuroIRIS). NeuroIRIS is a relatively rare (less than 1% of those who initiate ART) consequence of the introduction of ART in highly immunosuppressed patients and is marked by a severe deterioration in neurologic status that is characterized by massive lymphocytosis, extensive demyelination, and white matter damage (Anthony et al., 2005; Boisse et al., 2008). In neuroIRIS there is a paradoxical clinical deterioration in spite of improved CD4+ T-cell counts and decreased viral loads (McCombe et al., 2009). Despite the overall effectiveness of ART in limiting the infiltration of infected cells into the CNS, neuroinflammation still persists. However, the primary sites of neuroinflammation are different (Table 2.1); a strong involvement of the basal ganglia was observed pre-ART, whereas post-ART specimens display prominent signs of inflammation in the hippocampus and adjacent

parts of the entorhinal and temporal cortex (Anthony et al., 2005; Ho et al., 1985). Notably, these autopsy studies have demonstrated microglial activation in brains of individuals treated with ART comparable to those of patients with fully developed, pre-ART AIDS, although HIVE (when defined as the presence of HIV-infected multinucleated giant cells) is much less common now. Overall, these studies confirm the notion that neuroinflammation continues to be associated with HIV CNS infection in ART-experienced individuals, albeit without HIVE.

**Table 2.1 - Changes in the neuropathology of HAND in the era of HAART**

	<b>Pre HAART</b>	<b>Post HAART</b>
CSF viral load	viral load correlates with cognitive dysfunction (Brew et al., 1997; Ellis et al., 2000; Ellis et al., 1997; Ellis et al., 2002; McArthur et al., 1997)	CSF viral load is decreased with treatment (Letendre et al., 2008)
cognitive dysfunction	severe in HAD and improved when patients begin HAART (Letendre et al., 2008)	variable, from mild to severe (Antinori et al., 2007; McArthur et al., 2004)
predictive biomarkers of dementia	CSF viral load, decreased CD4 <sup>+</sup> T-cells, increased levels of CCL2 (Bandaru et al., 2007; von Giesen et al., 2005)	increased levels of CCL2, $\beta$ 2 microglobulin, quinolinic acid, arachidonic acid metabolites, oxidative stress markers, platelet activating factor (Bandaru et al., 2007)
major pathological findings	multinucleated giant cells, microglial nodules, neuronal loss, astrocytosis, myelin pallor, activated microglia and perivascular macrophages	neuronal loss, astrocytosis, myelin pallor, activated microglia and perivascular macrophages.
multinucleated giant cells and microglial nodules	present	absent
neuroinflammation	severe and progressive	less severe, but chronic
site of neuroinflammation	basal ganglia (Ho et al., 1985)	hippocampus and entorhinal and temporal cortex (Anthony et al., 2005; Ho et al., 1985)
microglial activation	present	present
CNS complications from Tx	N/A	NeuroIRIS (rare <1%) (Boisse et al., 2008)

## **2.2 - Biology of HIV Infection and Invasion of the Brain**

As mentioned above, HIV-1 traffics into the brain early in the course of infection via infected monocytes and lymphocytes (Dunfee et al., 2006). Despite retroviral therapy, HIV-1 persists in the CNS throughout the duration of infection in parenchymal microglia and perivascular macrophages (Ho et al., 1985; Koenig et al., 1986; Petito et al., 1986). Because increased numbers of microglia and macrophages correlate with the severity of pre-mortem HAND, these cell types probably mediate neurological impairment (Anthony et al., 2005; Glass et al., 1995; Petito et al., 1986). Multiple pro-inflammatory cytokines, including interleukin (IL)-1 $\beta$ , tumor necrosis factor-alpha (TNF- $\alpha$ ), and IL-6 are elevated in the CNS and/or CSF of patients with HAD (Achim et al., 1993; Folini et al., 1997; Oster et al., 1987; Perrella et al., 1992). The pro-inflammatory environment within the CNS is a result of cytokine release from monocytes/macrophages stimulated by either direct viral infection or by shed viral proteins (Rappaport et al., 1999; Sundar et al., 1991) and astrocyte activation without productive infection but with immune activation (Kramer-Hammerle et al., 2005; Sabri et al., 2003). Inflammatory mediators modulate the permeability of the blood-brain barrier, the entry of infected monocytes into the CNS, and peripheral processes that contribute to the neurological complications of HIV. Thus, understanding the mechanisms responsible for HIV entry into the CNS and the modulation of replication within the monocyte/macrophage reservoir are important for developing targeted therapeutics for HAND.


### *Integrity of the Blood-Brain Barrier (BBB): Role for Neuroinflammation*

The blood-brain barrier (BBB) separates the CNS from the periphery and modulates the traffic of low-molecular-weight nutrients, peptides, proteins, and cells in and out of the brain. The integrity and traffic across the BBB can be impacted by many factors, including HIV-dependent cytotoxicity towards cellular BBB components, chemotactic gradients, and the regulation of adhesion molecules and tight junction proteins (Figure 2.1). Progressive HIV infection and immune compromise result in the breakdown of the BBB (Dallasta et al., 1999; Kanmogne et al., 2002;

Persidsky et al., 2000), which permits the entry of free virus, lymphocytes, and infected and/or activated monocytes into the CNS.

The BBB is composed of brain microvascular endothelial cells (BMECs) that are connected by intercellular junctions to form a semipermeable monolayer. HIV infection increases the permeability of the BBB by compromising the integrity of the tight junctions. Brain regions with HIVE demonstrate an accumulation of activated/infected perivascular macrophages and a decrease in the tight junction membrane proteins zonula occludens (ZO-1) and occludin (Dallasta et al., 1999; Persidsky et al., 2006). TNF- $\alpha$  production by activated/infected macrophages can directly increase the permeability of the BBB to free virus (Fiala et al., 1997). However, these profound structural changes of the BBB are late events associated with encephalitis and HIV neuroinvasion is an early and continuing process.

HIV enters the CNS compartment through infected monocytes that cross the BBB via trans-endothelial migration, and this process is particularly enhanced by monocyte immune activation (Figure 2.1). Infected and immune-activated monocytes induce adhesion molecules on BMECs, thereby increasing transmigration during HIV infection. In HIVE brain tissue, viral load and pro-inflammatory cytokines correlate with levels of the adhesion molecules E-selectin and VCAM-1 (Hurwitz et al., 1994; Nottet et al., 1996; Persidsky et al., 1997; Sasseville et al., 1994). Exposure to TNF- $\alpha$  stimulates astrocytes to produce ICAM-1, VCAM-1, IG9, and E-selectin, all of which promote monocyte attachment and transmigration (Hurwitz et al., 1994). Following HIV infection, inflammatory cytokines promote the expression of adhesion molecules on BMECs and thereby promote the transmigration of activated/infected monocytes into the brain. Whether compromise to the BBB is as common now in the ART era as in the pre-ART (HIVE) era remains to be determined.



**Figure 2.1 Role for inflammation via cytokines and chemokines in HIV neuroinvasion and neurodegeneration.** HIV-infected monocytes cross the BBB, differentiate into macrophages, and productively infect microglia and other macrophages. Infected and/or activated macrophages/microglia and immune activated astrocytes release pro-inflammatory cytokines which promote monocyte neuroinvasion by increasing expression of adhesion molecules required for monocyte attachment and increasing BBB permeability. This pro-inflammatory environment is enhanced by LPS- and cytokine-induced systemic immune activation, which promotes monocyte recruitment and further BBB permeability. Together, the resulting neuroinflammation can directly and indirectly cause neuronal loss. Infected and/or activated macrophages/microglia and astrocytes also release chemokines, which can serve as both neurotoxic pro-inflammatory factors and neuroprotectants against HIV-induced neurotoxicity. Cytokines and chemokines can undergo reciprocal modulation by additional neurotoxic factors released from infected and/or activated macrophages/microglia, including excitatory amino acids (EAAs), phospholipids and reactive oxygen species (ROS).

### *Inflammation in the Periphery*

Pathology in the periphery may also be involved in the development of HAND (Figure 2.1) (Gartner, 2000). Neurological complications following HIV infection have been associated with an expanded population of circulating blood monocytes expressing markers of activation (Pulliam et al., 1997). These activated CD14<sup>+</sup>/CD16<sup>+</sup> monocytes produce TNF- $\alpha$  and IL-1 $\beta$ , which contribute to inflammation in the periphery and further activation of other immune cells (Thieblemont et al., 1995). CD16<sup>+</sup> monocytes are particularly susceptible to HIV infection, are capable of tissue invasion, and they compose the majority of the accumulated perivascular macrophages in patients with HIVE (Ellery et al., 2007; Jaworowski et al., 2007; Shiramizu et al., 2005). Peripheral activation of monocytes leads to an enhancement of infected and invasive monocytes that can be recruited into the CNS by chemokines (chemotactic cytokines), as discussed in detail below.

In the gastrointestinal tract, HIV infection also causes microbial translocation, in which HIV-driven depletion of gut-associated lymphoid tissue (GALT) results in leakage of bacteria into the bloodstream and subsequently increases systemic bacterial lipopolysaccharide (LPS) (Brenchley et al., 2006b; Douek, 2007). LPS increases monocyte transmigration into the CNS by contributing to the systemic immune activation of chronic HIV infection. In addition to priming peripheral monocytes for neuroinvasion, bacterial LPS can also compromise the integrity of the BBB. In vitro studies demonstrate that LPS-stimulated macrophages create gaps between endothelial cells of an artificial BBB, resulting in enhanced monocyte transmigration (Persidsky et al., 1997; Wang et al., 2008; Zhou et al., 2006). Notably, studies in SIV-infected rhesus macaques and HIV-infected humans correlate higher levels of plasma LPS, LPS-binding protein, and soluble CD14 with increased HAND severity (Ancuta et al., 2008) and more rapid progression towards AIDS (Brenchley et al., 2006b). Thus, the neurological complications of HIV are not exclusively mediated by pathological processes within the CNS, and components of systemic immune activation also play a significant role in the development of HAND.

In summary, inflammatory mediators contribute to HIV infection of the brain by affecting components of both the CNS and the peripheral compartments. Pro-inflammatory cytokines activate a neuroinvasive subset of monocytes in the periphery, compromise the integrity of the BBB, and promote the transmigration of infected monocytes into the CNS. In addition to mediating HIV neuroinvasion, inflammatory mediators also promote the accumulation of macrophages in the brain. These cells serve as a primary reservoir for HIV in the CNS and contribute to the pathogenesis of HAND via the release of neurotoxic and inflammatory cellular products, which can activate noninfected cells (macrophages/microglia, astrocytes). Understanding the role of inflammation in mediating HIV infection of the CNS could provide new therapeutic targets for HAND and other neuroinflammatory diseases.

### **2.3 - Mechanisms of HIV-Induced Neurodegeneration: Roles for Chemokines, Chemokine Receptors, and Inflammation**

Monocyte infiltration and macrophage/microglia activation are thought to initiate HAND pathogenesis through both systemic and CNS inflammatory signaling (Kaul et al., 2001; Kraft-Terry et al., 2009; Yadav and Collman, 2009). Following systemic inflammation and monocyte infiltration, activated and/or infected macrophages/microglia within the CNS can release a variety of neurotoxic factors, including viral proteins (gp120, Tat), pro-inflammatory cytokines (TNF- $\alpha$ , IL-1 $\beta$ , IL-6), interferons (IFN- $\alpha$ , IFN- $\beta$ , IFN- $\gamma$ ), excitatory amino acids (glutamate, quinolinic acid), phospholipids (platelet activating factor, arachidonic acid), and reactive oxygen species (Boven et al., 1999; Brenneman et al., 1988; Brew et al., 1995; Gelbard et al., 1993; Gelbard et al., 1994; Gendelman et al., 1998; Genis et al., 1992; Jiang et al., 2001; Maragos et al., 2003; Scorziello et al., 1998; Song et al., 2003; Wesselingh et al., 1993). Many of these factors can undergo reciprocal modulation by chemokines. Indeed, chemokines and chemokine receptors expressed within the CNS have central roles in HIV neuropathogenesis, from the function of chemokine receptors in mediating infection of the macrophage/microglia reservoir, to the seemingly dichotomous roles of chemokines as neurotoxic pro-inflammatory factors and neuroprotectants



against HIV-induced neurotoxicity (Collman and Yi, 1999; Doms, 2000; Gonzalez-Scarano and Martin-Garcia, 2005; Martin-Garcia et al., 2002). Here, we focus on the  $\alpha$ -,  $\beta$ -, and  $\delta$ -chemokine subfamilies in relation to HAND pathogenesis.

### *$\alpha$ -Chemokines*

The  $\alpha$ -chemokines, which bind CXCR chemokine receptors, affect HAND neuropathogenesis by enhancing neuroinvasion, promoting astrocyte activation, and directly acting as neurotoxic or neuroprotective factors (Figure 2.1). Increased expression of several  $\alpha$ -chemokines has been documented in the CSF and brain tissue of HAD patients, including CXCL10 ( $\gamma$ -interferon-inducible protein 10, IP-10) and CXCL12 (stromal cell-derived factor-1, SDF-1) (Cinque et al., 2005; Kolb et al., 1999; Rostasy et al., 2003; Zhang et al., 1998). In vitro studies demonstrate that CXCL10 is expressed in microglia and astrocytes, and that gp120, Tat, TNF- $\alpha$  and IFN- $\gamma$  can act independently or synergistically to increase CXCL10 release (Asensio et al., 2001; D'Aversa et al., 2004; Dhillon et al., 2008a; Kutsch et al., 2000; Williams et al., 2009a; Williams et al., 2009b). In turn, soluble CXCL10 can directly induce apoptosis in neurons (Sui et al., 2004; Sui et al., 2006a; van Marle et al., 2004) and increase neuroinflammation through leukocyte recruitment into the CNS (Asensio et al., 2001; Dhillon et al., 2008a; Kolb et al., 1999).

Like CXCL10, CXCL12 is a potent chemoattractant that can increase recruitment, adhesion, and transendothelial migration of monocytes into the CNS (Malik et al., 2008; Peled et al., 1999; Peng et al., 2006; Rostasy et al., 2003; Wu et al., 2000). CXCL12 is expressed in astrocytes, microglia, and neurons, and exposure to LPS or IL-1 $\beta$  from HIV-infected and/or activated macrophages can increase CXCL12 release (Bajetto et al., 1999; Ohtani et al., 1998; Peng et al., 2006).

Conversely, CXCL12 can trigger astrocytic release of TNF- $\alpha$  and glutamate, causing neuronal damage and apoptosis (Bezzi et al., 2001). Like CXCL10, CXCL12 can also act as a direct neurotoxin. It can undergo proteolytic cleavage by matrix metallic proteinase-2, which changes its co-receptor specificity from CXCR4 to CXCR3, the receptor for CXCL10, and enhances its

neurotoxicity (Vergote et al., 2006; Zhang et al., 2003). The neurotoxic properties of native, non-cleaved CXCL12 are controversial as CXCL12 exposure produces either neuroprotective or neurodegenerative responses depending on the experimental conditions (Hesselgesser et al., 1998; Kaul and Lipton, 1999; Khan et al., 2004; Lazarini et al., 2000; Zheng et al., 1999a; Zheng et al., 1999b).

### *$\beta$ -Chemokines*

Like  $\alpha$ -chemokines,  $\beta$ -chemokines, which bind CCR receptors, can mediate neurotoxic and neuroprotective effects against HIV-induced neurotoxicity (Figure 2.1). Several ligands in the  $\beta$ -chemokine subfamily are expressed at increased levels during HIV infection, including CCL2, CCL3 (macrophage inflammatory protein-1 $\alpha$ , MIP-1 $\alpha$ ), CCL4 (MIP-1 $\beta$ ), and CCL5 (regulated on activation, normal T cell expressed and secreted (RANTES; (Kelder et al., 1998)). Interestingly, CCL2 CSF levels increase in SIV-infected macaques and HIV-infected individuals prior to neurocognitive impairments and correlate with severity of dementia, suggesting that CCL2 can be a predictive marker for clinical HAND (Cinque et al., 1998; Dhillon et al., 2008b; Kelder et al., 1998; Ragin et al., 2006; Seignin et al., 2004; Seignin et al., 2007; Zink et al., 2001; Zink et al., 1998). In vitro studies demonstrate that CCL2 is expressed and released from endothelial cells, macrophages, microglia, and astrocytes during HIV infection and in response to gp120, Tat, TNF- $\alpha$ , IL-1 $\beta$ , IFN- $\beta$ , and IFN- $\gamma$  (Choe et al., 2001; Conant et al., 1998; D'Aversa et al., 2004; Gu et al., 1997; Guillemin et al., 2003; Lehmann et al., 2006; McManus et al., 2000; Mengozzi et al., 1999). HIV infection of macrophages increases their expression of the CCL2 receptor, CCR2, and concomitantly increases CCL2-mediated recruitment and transmigration of HIV-infected monocytes into the CNS (Eugenin et al., 2006; Park et al., 2001). Thus, inflammation following HIV infection establishes a cycle of monocyte/macrophage activation and neuroinvasion mediated by CCL2. However, CCL2 also provides neuroprotection against Tat-induced neurotoxicity (Eugenin et al., 2003; Yao et al., 2009), suggesting that increased expression of CCL2 during HIV infection may play a destructive and/or protective role. Additionally, the potential

CNS destructive role for CCL2 in HIV infection thus appears to be driven through its enhancement of monocyte transendothelial migration and not direct effects on neurons.

While elevated CCL2 CSF levels are associated with an increased risk of HAND, the association of CCL3, CCL4, and CCL5 with HAND is unclear (Letendre et al., 1999). CCL3, CCL4, and CCL5 can all serve as ligands for the CCR5 receptor and can suppress CCR5-mediated HIV infection (Cocchi et al., 1995). Studies also demonstrate that CCL3, CCL4, and CCL5 are expressed and released from microglia and astrocytes during HIV infection and in response to Tat (Cota et al., 2000; D'Aversa et al., 2004; El-Hage et al., 2005; Si et al., 2002), and can ameliorate excitatory amino acid- and gp120-induced neurotoxicity in vitro (Bruno et al., 2000; D'Aversa et al., 2004; Kaul and Lipton, 1999; Meucci et al., 1998; Meucci et al., 2000). The neuroprotective effects of CCL5 could be mediated by CCL2 induction (Eugenin et al., 2003). Together, these studies suggest that  $\beta$ -chemokines play important roles in modulating neuroinflammation and neurotoxicity during HIV infection and that they could express either protective or destructive effects within the CNS inflammatory microenvironment.

#### *$\delta$ -Chemokines*

CX3CR1 and its unique ligand, CX3CL1 (fractalkine), are the only known receptor-ligand pair in the  $\delta$ -chemokine subfamily. Furthermore, CX3CL1 is the only chemokine expressed in higher amounts in the CNS than in peripheral tissues, suggesting its critical role in modulation of HAND pathogenesis (Figure 2.1) (Bajetto et al., 2001; Cotter et al., 2002; Re and Przedborski, 2006). It is expressed as a membrane-anchored form on the cell surface or as a soluble form, which can be proteolytically released from the cell. CX3CL1 expression is elevated in serum, CSF, and brain tissue of HAD patients (Erichsen et al., 2003; Pereira et al., 2001; Sporer et al., 2003) and in vitro studies suggest that exposure to purified HIV virus, gp120, TNF- $\alpha$ , IL-1 $\beta$ , IFN- $\gamma$ , or glutamate increases expression of soluble and membrane-bound forms of CX3CL1 in neurons and astrocytes (Chapman et al., 2000a; Erichsen et al., 2003; Maciejewski-Lenoir et al., 1999; Pereira

et al., 2001; Sporer et al., 2003). Both forms of CX3CL1 can serve as potent CNS leukocyte chemoattractants and can mediate monocyte adhesion and transendothelial migration across the BBB (Ancuta et al., 2003; Chapman et al., 2000b; Harrison et al., 1998; Imai et al., 1997; Tong et al., 2000). In addition to recruiting activated peripheral blood monocytes, fractalkine can trigger the production of IL-6 and CCL2 by CD16<sup>+</sup> monocytes (Ancuta et al., 2003). Elevated CCL2 levels can lead to further recruitment of monocytes to the brain while IL-6 can activate and promote the differentiation of monocytes into macrophages, which could promote neurodegeneration. CX3CL1 differentially modulates other neuroinflammatory factors as well, increasing secretion of TNF- $\alpha$  and IL-8, from monocytes and macrophages (Ancuta et al., 2003; Ancuta et al., 2006; Cotter et al., 2002), while inhibiting TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 release from LPS-activated microglia and attenuating cytokine-mediated neuronal loss in vivo and in vitro (Cardona et al., 2006; Mizuno et al., 2003).

Like  $\beta$ -chemokines, CX3CL1 can also promote neuronal survival, as it can protect against gp120- and Tat-induced neurotoxicity in vitro (Deiva et al., 2004; Limatola et al., 2005; Meucci et al., 1998; Meucci et al., 2000). While it is unclear what factors maintain the in vivo balance between these neurotoxic and neuroprotective functions of CX3CL1 and other chemokines, the dysregulation of chemokine signaling likely significantly contributes to HAND pathogenesis. These processes are central to neurodegenerative diseases such as HIV infection and multiple sclerosis, and have implications for other neuroinflammatory diseases (Kaul and Lipton, 2006; Li and Ransohoff, 2008; Ransohoff, 1999; Ransohoff and Zamvil, 2007; Savarin-Vuillat and Ransohoff, 2007).

## **2.4 - Inflammatory Markers in NeuroAIDS and Other Neuroinflammatory Diseases**

Chronic neuroinflammation and elevated levels of pro-inflammatory cytokines and chemokines are associated with several neurodegenerative disorders of the CNS including HAND, MS, AD, PD, and HAM/TSP (Block and Hong, 2005; Grant et al., 2002; Mrak and Griffin, 2005; Sawada et

al., 2006). In the CNS, macrophages/microglia are the principal mediators of inflammation and, when activated, secrete pro-inflammatory cytokines (including TNF- $\alpha$ , IL-1 $\beta$ , and IL-6), chemokines (including CCL2 and CCL3), and adhesion molecules (ICAM-1, VCAM-1) that promote inflammation. Regardless of the factor initiating microglia activation, a chronic inflammatory response in the brain can contribute to the death of vulnerable neuronal populations and understanding common mechanisms in these neuroinflammatory disorders could identify common targets for broadly protective drugs. Additionally, more reliable biomarkers for neuroinflammatory diseases are needed in order to make early and accurate diagnoses, monitor the course of disease progression, and predict a patient's response to therapy.

#### *Inflammation in Neurodegenerative Diseases Associated with Aging*

HAND shares clinical features with normal aging including the deterioration of cognitive abilities and working memory (Alirezai et al., 2008; Kaul, 2009). Of interest, diffusion tensor imaging studies, where diffusion is an indicator of neuroinflammation, have shown altered water diffusion within specific brain regions in HIV-infected patients in comparison to normally aging control patients (Chang et al., 2008). These data suggest that even a well-controlled HIV infection may accelerate aging and promote neurodegeneration. In addition, there are several pathological features shared between HAND and neurodegenerative diseases associated with aging, including AD and PD (Brew et al., 2009; Chang et al., 2008; Esiri et al., 1998; Khanlou et al., 2009). These include neuroinflammation, oxidative stress, and cellular degradation pathways (Brew et al., 2009; Lovell and Markesbery, 2007; Nath et al., 2008).

The neuropathology of HAND is clearly distinguishable from AD at autopsy, although there are some shared features. HAND demonstrates less atrophy and fewer neurofibrillary tangles in plaques, which are the hallmarks of AD. Amyloid beta (A $\beta$ ) deposition, which precedes symptoms in AD, has also been described in the brains of HIV-infected individuals, although its potential role in HAND symptoms is controversial (Brew et al., 2009; Esiri et al., 1998). A $\beta$  deposition has pro-

inflammatory effects that accelerate neurodegeneration in vitro (Craft et al., 2006). In AD brains, microglia localize to amyloid plaques (McGeer et al., 1987) and upregulate human leukocyte antigen-DR (HLA-DR) and complement in addition to several pro-inflammatory cytokines that are also implicated in HAND pathogenesis (IL-1 $\beta$ , IL-6, and TNF $\alpha$ ) (Eikelenboom et al., 2002; McGeer and McGeer, 2010; McGeer et al., 1987; Sheng et al., 1998). Notably, a transgenic mouse model for a familial AD mutation of amyloid precursor protein recapitulates this proinflammatory effect of A $\beta$  deposition and astrocytes and microglia expressing IL-1 $\beta$ , IL-6, TGF- $\beta$ , and TNF- $\alpha$  surround plaques (Qiao et al., 2001). In addition to common expression of cytokines that are believed to be involved in the pathogenesis of AD and HAND, LPS concentrations are elevated in both disorders (Ancuta et al., 2008; Herber et al., 2006).

As with AD, HAND also shares some similar neuropathological features with PD. The presence of  $\alpha$ -synuclein-positive inclusions in the cell bodies (Lewy bodies) and processes (Lewy neuritis) in the substantia nigra is the neuropathological hallmark of PD. Lewy bodies have also been observed in autopsied brains of HIV-infected individuals (Brew et al., 2009; Esiri et al., 1998; Kaul, 2009; Khanlou et al., 2009) and dopamine deficiency is found in both disorders. However, whether neuroinflammation in PD is a consequence or a cause of the selective loss of dopaminergic neurons in the substantia nigra is unknown. Evidence for neuroinflammation in PD includes increased CSF levels of proinflammatory cytokines, including TNF- $\alpha$ , IL-1 $\beta$ , IL-6, TGF- $\beta$ , and IFN- $\gamma$  (Hunot et al., 1999; Vawter et al., 1996). Moreover, single nucleotide polymorphisms associated with increased production of cytokines and chemokines are overrepresented in PD cohorts and may confer increased susceptibility to PD (Hakansson et al., 2005a, b; Kruger et al., 2000).

Although the role for neuroinflammation in AD and PD is not completely understood, inflammatory mediators may be used as both biomarkers and targets for drug development for these disorders. For example, platelet inflammatory biomarkers, like cyclooxygenase 2 (COX-2) and

phospholipase A2 could be exploited as peripheral inflammatory biomarkers (Casoli et al., 2010). COX-2 and its homolog COX-1 are pro-inflammatory proteins that are used by activated microglia to synthesize a variety of inflammatory mediators and are elevated in AD brains. Both COX-1 and COX-2 are targets of non-steroidal anti-inflammatory drugs (NSAIDs), and there is a link between chronic use of NSAIDs and a reduced risk for AD and PD (Breitner et al., 1995; Chen et al., 2003). Moreover, biomarkers identified in AD, PD, and other diseases related to aging may also be relevant in HAND. In a recent study,  $\beta$ -Amyloid(1–42)(A $\beta$ 42) measurements in the CSF of HAND patients were found to be similar to those found in patients with AD and significantly decreased compared to those in normal controls and HIV-infected individuals with normal cognitive function (Clifford et al., 2009). However, HAND patients had normal or slightly depressed levels of CSF tau and tau phosphorylated at threonine 181 (p-Tau181), which distinguished them from patients with AD (Clifford et al., 2009). The detection of pro-inflammatory proteins in the periphery may, in the future, be exploited as useful biomarkers in a wide spectrum of neurodegenerative disorders.

#### *Other Chronic Progressive, Inflammatory Disorders of the CNS with Known or Suspected Viral Etiology*

Neuroinflammation is a major component of other disorders of the CNS with known or suspected viral etiology including HAM/TSP and MS. The human retrovirus HTLV-1 (human T-lymphotropic virus type I) causes HAM/TSP in a small percentage (<5%) of infected individuals. The incubation period between infection with HTLV-I and the development of HAM/TSP is typically long (20–30 years) and the clinical hallmark of this chronic progressive neurologic disorder is a gradual onset of lower extremity weakness (McFarlin and Blattner, 1991; Osame et al., 1990). HTLV-I is predominantly CD4+T-cell-tropic. However, CD8+ T cells, astrocytes, monocytes/macrophages, and microglia may also become infected and serve as viral reservoirs (Hoffman et al., 1992; Nagai et al., 2001a; Nagai et al., 2001b; Watabe et al., 1989). As in HAND, monocytes/macrophages are believed to contribute to the pathogenesis of HAM/TSP. However,

in contrast to the pathogenesis of HAND/HIV-1 infection of the CNS, much of the neurodegeneration resulting from HTLV-I infection is caused directly by HTLV-I Tax-specific cytotoxic T cells (CTLs) restricted to immunodominant epitopes of HTLV-I gene products (predominantly Tax) (Elovaara et al., 1993; Jacobson et al., 1990b). These HTLV-I-specific CTLs are readily detected in the peripheral blood lymphocytes (PBLs), CSF, and active inflammatory lesions of HAM/TSP patients (Greden et al., 1998; Levin et al., 1997). HTLV-I Tax-specific CTLs secrete many proinflammatory cytokines and chemokines including IFN- $\gamma$ , TNF- $\alpha$ , CCL3, CCL4, and IL-16 (Biddison et al., 1997). In addition, HTLV-I Tax trans-activates many host genes and HTLV-I-infected astrocytes secrete high levels of IL-1 $\alpha$ , IL-6, TNF- $\alpha$ , and matrix metalloproteinases (MMPs) (Szymocha et al., 2000a; Szymocha et al., 2000b). Increased levels of HTLV-I tax mRNA, an increased frequency of HTLV-I Tax-specific CD8<sup>+</sup> T cells, and high proviral loads correlate with disease severity in HAM/TSP and provide good biomarkers of disease progression (Yamano et al., 2002). Other candidate HAM/TSP biomarkers, CD244 (a signaling lymphocyte activation molecule [SLAM] family receptor) and SLAM-associated protein (SAP), were found to be significantly higher in HAM/TSP compared to asymptomatic carriers and uninfected individuals and both may be used as biomarkers for neurodegeneration in HTLV-I-infected individuals (Enose-Akahata et al., 2009).

Another prototypic neuroinflammatory disease with some neuropathological similarities to HAND is MS. MS is the most common inflammatory disease of the CNS, with a prevalence that ranges between 2 and 150 per 100,000 (Rosati, 2001). The etiology of MS is unknown. In part, this is attributable to the variability of this disease, suggesting that many factors may be involved in the development of MS (reviewed in (Soldan et al., 2008)). However, it is generally believed that genetic, immunological, and environmental factors contribute to MS pathogenesis. Infectious agents have been implicated in the pathogenesis of MS for over 100 years but no single causative agent has been identified. Again, unlike HAND, MS is chiefly a T-cell-mediated neuroinflammatory disorder. However, in both MS and HAND an inflammatory cascade



contributes to disease pathogenesis. Although the neuroinflammatory nature of MS has been confirmed at all stages, inflammation tends to decrease after decades of disease, as irreversible neuronal degeneration accumulates. Such neuroinflammation is detectable through brain magnetic resonance imaging (MRI) as the presence of gadolinium-enhancing lesions, which represent areas of inflammation associated with leakiness of capillary endothelial cells and infiltration of serum components into the brain parenchyma. Analysis of CSF often reveals markers of neuroinflammation, as does examination of autopsied brain specimens (see below).

The overexpression of several proinflammatory cytokines, including TNF- $\alpha$ , IFN- $\gamma$ , IL-12, IL-6 and CXCL10 have been demonstrated in MS brain specimens (Bartosik-Psujek and Stelmasiak, 2005; Drulovic et al., 1998; Drulovic et al., 1997; Frohman et al., 2006; Miljkovic et al., 2002; Ubogu et al., 2006). Importantly, an increase in TNF- $\alpha$  expression in peripheral blood mononuclear cells has been found to precede MS relapses and inflammatory activity (Rieckmann et al., 1995). In contrast, anti-inflammatory cytokines (IL-4, IL-10, and TGF- $\beta$ ) and other chemokines (CCL2 and CCL5) are downregulated during MS disease exacerbations (Mahad et al., 2002a, b; Malmestrom et al., 2006; Rieckmann et al., 1995). The altered expression of these cytokines promotes disease pathogenesis by upregulating MHC and adhesion molecule expression on endothelial and glial cells, activating macrophages, recruiting TH-1 cells, and/or by directly damaging oligodendrocytes and myelin sheaths. In addition, soluble adhesion molecules such as ICAM-1 and E-selectin are elevated in the sera of MS patients while soluble vascular cell adhesion molecules VCAM-1 and E-selectin are increased in the CSF of MS patients, thereby promoting the trafficking of activated T cells into the CNS (Dore-Duffy et al., 1995).

Another long-standing diagnostic biomarker for MS is the presence of oligoclonal bands, which represent intrathecally expressed immunoglobulin. While the presence of oligoclonal bands clearly suggests neuroinflammation in the CSF, the MS-relevant reactive antigens have not yet been identified. CSF oligoclonal bands are also present in HAM/TSP and in these individuals they have

been found to be chiefly directed against HTLV-I (Jacobson et al., 1990a). More reliable biomarkers that can be assayed in the peripheral blood and CSF of MS patients are still being sought due to the high cost of frequent MRI and its poor reliability to detect neuronal degeneration, axonal loss, and to some extent, spinal cord lesions.

There are many molecules with the potential for more specific clinical diagnostic and prognostic application in MS including auto-antibodies, virus antibodies, transcription factors, molecules in the nitric oxide pathway, neuronal breakdown products, apolipoprotein E (a marker for cognitive dysfunction in AD), molecules in the amyloid precursor protein pathway, cytokines, and chemokines among others (reviewed in (Harris and Sadiq, 2009)). Several biomarkers of MS inflammatory disease activity are also of interest in HAND, including IL-6 and osteopontin, which are both upregulated in HAND and in MS lesions (Burdo et al., 2008; Cannella and Raine, 1995; Frei et al., 1991) and CCL2, which is downregulated during MS exacerbations and increased in HAND (Kelder et al., 1998). Further assessment of these biomarkers and the use of proteomics and microarray technologies to discover new and specific biomarkers for neuroinflammatory diseases will ultimately improve diagnosis and treatment of these disabling neuroinflammatory disorders.

## **2.5 - Therapeutic Considerations**

Given the complexity of the pathogenesis of HIV-associated neurodegeneration, multiple cell types and cellular processes are under investigation as therapeutic targets. Ongoing clinical trials are examining drugs that target overall neuronal survival in addition to reducing neuroinflammation. It is likely that a combination of classes of drugs will be necessary to ameliorate the neurocognitive decline in HIV patients receiving ART.

ART is increasingly effective in reducing morbidities and mortality in HIV-1-infected patients.

While ART has significantly decreased the incidence of HAD, presumably by lowering systemic

viral loads, current efforts are focused on improving the penetration of ART past the BBB. Improved penetration or retention of bioavailable ART drugs in the CNS may slow HAND progression by decreasing CNS viral replication and reducing concomitant release of neurotoxins from infected and activated macrophages/microglia (Ellis et al., 2007; Letendre et al., 2008; Spitzenberger et al., 2007). However, a recent study demonstrated that ART regimens with strong CNS penetration and reduced CSF viral loads were associated with poorer neurocognitive outcomes (Marra et al., 2009). While further clinical studies are needed, this study highlights the importance of considering drug toxicity when promoting ART in the CNS. Furthermore, effective therapies for HAND will likely require combination therapy that not only targets viral load, but also addresses the indirect pathways known to contribute to HIV-associated neurodegeneration.

Clinical similarities between HIV-associated neurodegeneration and other neurodegenerative diseases have prompted investigation into currently approved neuroprotective therapies in HAND (Table 2.2). Memantine (Namenda), approved in the treatment for Alzheimer's disease, is a non-competitive NMDA receptor antagonist that also increases levels of brain-derived neurotrophic factor (BDNF) and conserves dopamine function in SIV-infected macaques (Meisner et al., 2008). Both in vitro and in vivo animal studies have shown that memantine inhibits gp120 and Tat-induced neurotoxicity (Anderson et al., 2004; Nath et al., 2000; Toggas et al., 1996). A short-term clinical trial in HAND patients demonstrated that memantine improved neuronal metabolism (as judged by magnetic resonance spectroscopy/MRS), indicative of neuroprotection, but did not cause significant neurocognitive improvement (Schifitto et al., 2007a). Nevertheless, this study suggests a potential beneficial effect of memantine, even following short-term administration, although a longer-term follow-up study in this patient cohort failed to reveal a clinically demonstrable neurological benefit (Zhao et al., 2010).

Selegiline (Deprenyl), a monoamine oxidase B (MAO-B) inhibitor used in the treatment of early-stage Parkinson's disease, has shown some promise in clinical trials for HAND. Selegiline is

proposed to act as a neuroprotectant by reducing the antioxidant burden of the cell (Magyar and Szende, 2004). Early trials with orally or transdermally administered selegiline demonstrated some improvement in psychomotor speed (Sacktor et al., 2000; The Dana Consortium, 1998). However, recent studies have shown neither a reduction in biomarkers of oxidative stress nor evidence of cognitive improvement with short-term (24 weeks) transdermal selegiline (Schifitto et al., 2009a; Schifitto et al., 2007b). While memantine and selegiline have shown some effectiveness, they are clearly not potential monotherapies for HAND. In addition to using such drugs prior to neurocognitive decline, effective neuroprotective therapies will likely have to be used in combination and over the duration of viral infection in order to have maximal clinical benefit.

**Table 2.2 Therapeutics under consideration for HAND**

<b>Generic (Brand) Name</b>	<b>Molecular Target</b>	<b>Effects/Role</b>	<b>References</b>
Memantine (Namenda)	NMDA receptor antagonist	Increases BDNF levels, conserves dopamine function, inhibits gp120 and Tat-induced neurotoxicity, improves neuronal metabolism, short-term treatment provided no neurocognitive impairment	(Anderson et al., 2004; Meisner et al., 2008; Nath et al., 2000; Schifitto et al., 2007a; Toggas et al., 1996; Zhao et al.)
Selegiline (Deprenyl)	MAO-B Inhibitor	Proposed to reduce antioxidant burden of the cell, may improve psychomotor speed, no clinical evidence of cognitive improvement over 24 week treatment	(Magyar and Szende, 2004; Sacktor et al., 2000; Schifitto et al., 2009a; Schifitto et al., 2007b; The Dana Consortium, 1998)
Sodium valproate and lithium	GSK-3 $\beta$ inhibitor	Reduces neurotoxicity, improved neuropsychological performance	(Ances et al., 2008; Dou et al., 2003; Everall et al., 2002; Letendre et al., 2006; Schifitto et al., 2006; Schifitto et al., 2009b; Tong et al., 2001)
SSRIs – citalopram, paroxetine	Serotonin transporter	May decrease HIV viral levels in CSF, improved adherence to ART	(Ances et al., 2008; Letendre et al., 2007)
Minocycline	5-lipoxygenase and others	Decreases CCL2 levels in CSF, improves encephalitis, suppresses HIV replication and inhibits secretion of TNF $\alpha$ , IFN $\gamma$ and IL-2 by lymphocytes	(Colovic and Caccia, 2003; Copeland and Brooks, 2010; Si et al., 2004; Szeto et al., 2010; Zink et al., 2005)
PMS-601	platelet-activating factor (PAF) receptor antagonist	Reduces neurotoxicity, microgliosis and TNF $\alpha$ , CCL3, CCL4 and CCL5 secretion by macrophages	(Eggert et al., 2009b; Martin et al., 2000)
Copolymer-1 (Copaxone)		Decreases microgliosis, astrogliosis, neurotoxicity and TNF $\alpha$ and IL-12 levels	(Gorantla et al., 2007; Gorantla et al., 2008)
CEP-1347		Anti-apoptotic, decreases monocyte secretion of TNF $\alpha$ , reduces microgliosis and neurotoxicity	(Bodner et al., 2002; Eggert et al., 2009a; Sui et al., 2006b)

In addition to therapies for neurodegenerative diseases, compounds in clinical use for neuropsychiatric disorders are also under investigation in HAND (Ances et al., 2008). Sodium valproate (VPA) and lithium are approved for treatment of bipolar disorder and related mood disorders, and both inhibit glycogen synthase kinase-3 $\beta$  and provide neuroprotection against HIV-induced toxicity in vitro and in mouse models of HIVE (Dou et al., 2003; Everall et al., 2002; Tong et al., 2001). Several small pilot studies have demonstrated improved neuropsychological performance in HAND patients following short-term VPA or lithium therapy (Letendre et al., 2006; Schifitto et al., 2006; Schifitto et al., 2009a). Additional studies using selective serotonin reuptake inhibitors (SSRIs), including citalopram and paroxetine, are also under consideration as adjunctive therapies for HAND (Ances et al., 2008; Letendre et al., 2007).

Other adjunctive therapies for HAND are focused on targeting inflammation cascades that contribute to neurotoxicity. Minocycline is a broad-spectrum tetracycline antimicrobial that is currently in phase I clinical trials for HAND (Copeland and Brooks, 2010). Minocycline is capable of efficiently crossing the BBB and can suppress HIV replication in microglia, macrophages, and lymphocytes (Colovic and Caccia, 2003; Si et al., 2004; Szeto et al., 2010; Zink et al., 2005). In addition, minocycline has anti-inflammatory properties and can inhibit the secretion of the inflammatory cytokines TNF- $\alpha$ , IFN- $\gamma$ , and IL-2 by lymphocytes (Szeto et al., 2010). Experimental studies using SIV-infected macaques demonstrated that minocycline decreased CSF levels of CCL2, a marker of CNS inflammation, and decreased the severity of encephalitis (Zink et al., 2005).

In addition to minocycline, several other therapies that have anti-HIV and anti-inflammatory effects are also being considered as adjunctive therapy for HAND. PMS-601, a platelet-activating factor (PAF) receptor antagonist, reduces neurotoxicity, microgliosis, and macrophage secretion of the inflammatory mediators TNF- $\alpha$ , CCL3, CCL4, and CCL5 in vitro and in a mouse model of HIVE (Eggert et al., 2009b; Martin et al., 2000). Copolymer-1 (COP-1 or Copaxone) is a clinically

approved immune modulator used in the treatment of MS. In addition to decreasing levels of TNF- $\alpha$  and IL-12, COP-1 decreases microgliosis, astrogliosis, and neurotoxicity in a mouse model of HIVE (Gorantla et al., 2007; Gorantla et al., 2008). CEP-1347 is an anti-apoptotic immune modulator that decreases monocyte secretion of TNF- $\alpha$  and macrophage secretion of chemokines, including CCL4 and CXCL10 (Eggert et al., 2009a; Sui et al., 2006b). CEP-1347 reduces microgliosis and HIV-mediated neurotoxicity in vitro and reduces signs of HIVE in a mouse model (Bodner et al., 2002; Eggert et al., 2009a; Sui et al., 2006b). As our understanding of the complex nature of HAND pathogenesis evolves, it is becoming clear that adjunctive therapies that address not only viral burden in the CSF but also the contribution of processes such as inflammation are critical for successful clinical management of HAND. In addition, the development of adjunctive therapies for HAND will contribute to and likely improve the clinical management of other neuroinflammatory disorders, including AD, PD, and MS.

## **CHAPTER 3**

### **MATERIALS AND METHODS**



### *Reagents*

Stock solutions of dimethyl fumarate and monomethyl fumarate (Sigma, St. Louis, MO), *tert*-butylhydroquinone (tBHQ; Acros Organics/Thermo Fischer Scientific, Geel, Belgium), thapsigargin (Tocris Bioscience, Ellisville, MO), Sal003 (Calbiochem/EMD Biosciences, La Jolla, CA), clotrimazole (Sigma) were prepared in DMSO and stored at -20°C until use. Tin (IV) mesoporphyrin IX dichloride (SnMP), cobalt (III) protoporphyrin IX chloride (CoPP), and hemin (Frontier Scientific, Logan, UT) were prepared in 1N NaOH and stored at -20°C until use. Stock solutions of Ara-C (Sigma), phytohemagglutinin (PHA; Sigma), TNF $\alpha$  (R&D Systems, Minneapolis, MN) and CCL2 (Peprotech, Rocky Hill, NJ) were prepared in filter-sterilized distilled water and stored at -20°C. Stock solutions of efavirenz (NIH AIDS Research and Reference Reagent Program, Germantown, MD) were prepared in DMSO and frozen at -80°C until use.

### *Isolation and culture of human monocyte-derived macrophages (MDM)*

All human studies were reviewed and approved by the Institutional Review Board at the University of Pennsylvania. Human monocytes were prepared from PBMCs of healthy donors and isolated by Ficoll density gradient centrifugation as previously described (Chen et al., 2002; O'Donnell et al., 2006). Monocytes were plated at  $1 \times 10^6$  cells per well to Cell-Bind 6-well plates (Corning, Lowell, MA) and cultured in DMEM supplemented with 10% fetal bovine serum (FBS), 10% horse serum, 1% non-essential amino acids with 50 U/mL penicillin/streptomycin at 37°C, 6% CO<sub>2</sub>. Cells were cultured for 7-8 days and visually inspected for MDM differentiation before use in HIV-infection experiments. MDM were cultured for 7-10 days before use in non-infectious experiments.

### *HIV infection of MDM*

MDM were pretreated with efavirenz, MMF, DMF or tBHQ, at the indicated concentrations, for 1 hour. All wells were normalized for the vehicles appropriate for drug treatments (DMSO and/or NaOH). Differentiated MDM were exposed to 50ng (p24 ELISA, equivalent to  $1.82 \pm 0.22$

kcpm/ $\mu$ L by reverse transcriptase (RT) activity assay) of HIV-1 Jago (R5 strain) or 89.6 (R5/X4 strain) for 24 hours. HIV-Jago is a macrophage tropic, CSF isolate from a patient with confirmed HIV-associated dementia (Chen et al., 2002). Virus stocks were prepared by the University of Pennsylvania Center for AIDS Research Virology Core. For some experiments, HIV/MDM were exposed to drug treatments, or appropriate vehicle controls, for 6 (0.5 $\mu$ M thapsigargin) or 24 hours (10 $\mu$ M clotrimazole or 50 $\mu$ M Sal003) beginning at day 6 post infection. Following drug exposure, wells were washed x1 with DMEM before cells were put into fresh, drug-free media for the remainder of the infection time course. Supernatants from HIV-infected or non-infected (Mock) MDM were collected every 2-4 days and stored at -80°C. Supernatants were monitored for HIV replication by quantifying viral RT activity, as analyzed by the amount of radiolabeled deoxythymidine incorporation.

#### *Subcellular fractionations and Western blot analysis*

For whole cell lysate collection, cells were rinsed twice with ice-cold PBS and lysed in 75mM Tris-HCl (pH 6.8), 15% glycerol, 3.75mM EDTA, 3% SDS and supplemented with Complete Protease Inhibitor Cocktail (Roche Applied Science, Indianapolis, IN) and PhosSTOP phosphatase inhibitor cocktail (Roche Applied Science).

To assess for nuclear translocation of NF- $\kappa$ B proteins, differentiated MDM were treated with DMF for 24 hours, exposed to TNF $\alpha$  (1 ng/mL) for 10 minutes and fractionated. To prepare nuclear extracts, cells were rinsed twice in ice-cold PBS and lysed on ice for 10 minutes in 10mM HEPES (pH 7.9), 10mM KCl, 10mM EDTA, 1mM DTT, 0.4% Nonidet P-40, supplemented with protease and phosphatase inhibitors. Nuclei were pelleted for 3 minutes at 16,000  $\times$  g and the supernatant (cytoplasmic fraction) was collected and stored at -20°C. The nuclear pellet was resuspended in 20mM HEPES (pH 7.9), 400mM NaCl, 1mM EDTA, 10% glycerol, 1mM DTT, protease and phosphatase inhibitors and incubated at 4°C on a rocking platform at 200rpm for 2 hours. After centrifugation at 16,000  $\times$  g for 5 minutes, supernatants (nuclear fractions) were collected and

stored at -20°C. All protein concentrations were determined by the Detergent Compatible (DC) protein assay (Bio-Rad Laboratories, Hercules, CA).

Cell lysates were subjected to SDS-PAGE as previously described (O'Donnell et al., 2006) using the following antibodies: rabbit anti-HO-1 (Stressgen/Enzo Life Sciences, Farmingdale, NY), mouse anti-NQO1 (Abcam, Cambridge, MA), mouse anti-Nrf2 (R&D Systems), rabbit anti-RelB (Cell Signaling Technologies, Danvers, MA), rabbit anti-NF-κB p65 (Cell Signaling), rabbit anti-NF-κB p50 (Cell Signaling), rabbit anti-poly (ADP-ribose) polymerase (PARP) (Cell Signaling), rat anti-GRP94 (Stressgen), mouse anti-BiP (BD Transduction Laboratories/BD Biosciences, Franklin Lakes, NJ), rabbit anti-phosphorylated eIF2α (Invitrogen, Carlsbad, CA), mouse anti-eIF2α (Cell Signaling), rabbit anti-β-tubulin (Cell Signaling), mouse anti-GAPDH (Advanced Immunochemical, Long Beach, CA), and species-specific HRP-conjugated secondary antibodies (Jackson ImmunoResearch Laboratories, West Grove, PA or Cell Signaling). For densitometry analysis, films were scanned and a fixed cursor area centered over each band was assessed for pixel density using ImageJ (NIH, Bethesda, MD).

#### *MDM-mediated neurotoxicity*

Rat cerebrocortical neuronal cultures were prepared from embryos of Sprague-Dawley rats at day 17 of gestation, as previously described (O'Donnell et al., 2006). All procedures were within the ARRIVE guidelines for animal research, and in accordance with protocols approved by the University of Pennsylvania Institutional Animal Care and Use Committee. Cells were plated in tissue culture dishes pre-coated with poly-L-lysine (Peptides International, Louisville, KY) and maintained in neurobasal media plus B27 supplement (Invitrogen) at 37°C and 5% CO<sub>2</sub>. Forty-eight hours after plating, cells were treated with 10μM Ara-C. After 7 days *in vitro* (DIV), approximately one-half volume of fresh media was added to the cells in order to counteract effects of evaporation. All cultures were used between 14 and 16 DIV.

Cell-based microtubule-associated protein 2 (MAP2) ELISAs were performed on primary rat cerebrocortical cells plated at a density of  $6 \times 10^4$  cells per well in 96-well plates. Following a 24 hour exposure to HIV/MDM supernatant, cultures were fixed and fluorescently labeled as described (Wang et al., 2007; White et al., 2011) using the following reagents: mouse anti-MAP2 (Covance, Princeton, NJ), goat anti-mouse  $\beta$ -lactamase TEM-1 conjugate (Invitrogen), and Fluorocillin Green substrate (Invitrogen). Fluorescence intensity was measured using a fluorometric plate reader with the 480/520 nm filter set. Macrophage supernatant was applied at a 1:10-1:50 dilution; the dilution that gave values within the linear range of the assay is presented.

#### *Immunofluorescence*

Primary rat cerebrocortical cells were plated at a density of  $2 \times 10^5$  cells per 35mm dish with glass coverslips. Following exposure to HIV/MDM supernatant for 24 hours, cultures were fixed and fluorescently labeled as described (O'Donnell et al., 2006) using the following reagents: mouse anti-MAP2 (Sigma) and species-specific Cy3-conjugated secondary antibody (Jackson ImmunoResearch Laboratories), and Hoescht 33342 (Invitrogen).

#### *LDH assay*

Soluble lactate dehydrogenase (LDH) in HIV/MDM culture supernatant was measured using the Cytotoxicity Detection KitPLUS (Roche Applied Science) according to manufacturer's instructions.

#### *Electrophoretic Mobility Shift Assay (EMSA)*

Following 24 hours of pretreatment with DMF, human MDM were exposed to 1 ng/mL TNF $\alpha$  for 10 minutes and nuclear protein extracts were isolated as described. 8 $\mu$ g of nuclear protein was assessed for NF- $\kappa$ B-DNA binding with an EMSA kit (Panomics, Santa Clara, CA), used according to manufacturer's directions. The labeled oligonucleotide for NF- $\kappa$ B p50 binding, 5'-AGTTGAGGGGACTTCCCAGGC-3', was used.

#### *Cytokine detection in culture supernatants*

The concentration of TNF $\alpha$  in culture supernatants was detected using an ELISA kit (Invitrogen) and used according to manufacturer's instructions. Uninfected MDM were treated with 0.067% DMSO (vehicle), DMF or tBHQ for 24 hours prior to exposure to 10  $\mu$ g/mL PHA for 6 hours. Supernatants were collected and frozen at -80°C until assayed.

#### *Chemotaxis assay*

Monocyte chemotaxis was assayed using the Chemicon QCM 96-well (5 $\mu$ m pore size) Migration kit (Millipore, Temecula, CA) according to manufacturer's directions. Freshly isolated human monocytes were plated at a density of  $2 \times 10^5$  cells/well in serum- and growth factor-free culture media to the upper chamber in the presence of DMF, MMF or vehicle (0.02% DMSO). CCL2 (300 ng/mL) was added to the lower chamber and cells were incubated at 37°C and 6% CO<sub>2</sub> for 6 hours (Chiou et al., 2011; Eugenin et al., 2006; Janic et al., 2008). Exposure of monocytes to 300 ng/mL CCL2 for 6 hours most consistently induced chemotaxis, with an average of  $35.3 \pm 20.2\%$  above baseline. All cells that had migrated through the insert, including those adhered to bottom of the membrane, were collected. For quantification, cells were lysed and labeled with CyQuant GR dye. Fluorescence was read with the 480/520 nm filter set on a fluorometric plate reader.

#### *Flow cytometry*

Human PBMCs were cultured in RPMI supplemented with 10% FBS and 50 U/mL penicillin/streptomycin at 37°C and 5% CO<sub>2</sub>. Following 6 or 36 hours of treatment with the indicated concentrations of DMF or DMSO vehicle, cells were washed with ice-cold FACS buffer (PBS, 1% BSA, 0.1% NaN<sub>3</sub>) and stained with CD11b-PE (clone ICRF44, eBioscience, San Diego, CA), CD14-PE/Cy7 (M5E2, BioLegend, San Diego, CA), CCR2-PerCP/Cy5.5 (TG5, BioLegend), CD4-FITC (OKT4, BioLegend), CD195/CCR5-Alexa Fluor 700 (HEK/1/85a, BioLegend) and CD184/CXCR4-PerCP/Cy5.5 (12G5, BioLegend) antibodies. Mouse IgG2a-PerCP/Cy5.5 (MOPC-173, BioLegend) was used as the isotype control for CCR2 staining. Antibody-stained cell

suspensions were pretreated with 4'6'-diamidino-2-phenylindole (DAPI) to identify dead cells. Flow cytometry was performed on a LSR-II (BD Biosciences). Doublets were excluded using forward side scatter-height versus forward side scatter-width and side scatter-height versus side scatter-width parameters. Data were analyzed using FlowJo (Tree Star, Ashland, OR). Monocytes were identified as CD11b<sup>+</sup>CD14<sup>+</sup> cells.

#### *Indirect fluorescence assay*

For immunofluorescent staining, paraffin-embedded tissue sections from the frontal cortices of control ( $n = 2$ ) and HIV(+) ( $n = 12$ ) human autopsy cases were obtained from the National Neuro-AIDS Tissue Consortium. Glass slides containing paraffin-embedded sections (5  $\mu$ m) were deparaffinized and rehydrated as described previously (Chalovich et al., 2005). Endogenous peroxidase activity was inactivated using 3% H<sub>2</sub>O<sub>2</sub> in methanol. Antigen unmasking was performed by target retrieval solution (Dako Corporation, Carpinteria, CA) at 95°C for 1 h. Tissue sections were then blocked with 10% normal goat serum in PBS. Mouse monoclonal antibodies to BiP (1:150, BD BioSciences) and the macrophage/microglia marker, HLA-DR (MHC class II protein, clone CR3/43, 1:1000, Sigma) were used in IFA for macrophagic BiP. Rabbit polyclonal antibodies to phosphorylated eIF2 $\alpha$  (1:1000, Invitrogen) and the macrophage/microglia marker, HAM56 (human alveolar macrophage, clone 56, 1:100, Dako Corporation) were used in IFA for macrophagic phosphorylated eIF2 $\alpha$ . The tyramide amplification system (New England Biolabs, Beverly, MA) was used according to manufacturer's directions for the detection of HLA-DR and phosphorylated eIF2 $\alpha$ . DNA was visualized by DAPI staining (5 mM, Molecular Probes, Carlsbad, CA). Slides were mounted in Vectashield (Vector Laboratories, Burlingame, CA) and analyzed by laser confocal microscopy on a Bio-Rad Radiance 2100 equipped with Argon, Green He/Ne, Red Diode and Blue Diode lasers (Bio-Rad), as described previously (Strachan et al., 2005). All images shown were captured with uniform threshold and intensity settings.

Post-acquisition analysis for immunofluorescent staining was performed using MetaMorph 6.0 image analysis software (Universal Imaging, Inc, Downingtown, PA, USA). Eight 212  $\mu\text{m}$   $\times$  212  $\mu\text{m}$  images were captured randomly from the areas of positive staining in the midfrontal cortical grey matter for each case. HLA-DR, HAM56 and phosphorylated eIF2 $\alpha$  area was determined by the measurement of the number of the pixels positive for the desired signal. Total macrophage specific BiP was determined by the measurement of integrated pixel intensity of BiP over pixels positive for HLA-DR, where integrated pixel intensity is defined as total pixel intensity per image times the area of pixels positive for the signal. Total HAM56 and phosphorylated eIF2 $\alpha$  intensity was determined by the measurement of integrated pixel intensity for HAM56 or phosphorylated eIF2 $\alpha$  per image, where the integrated pixel intensity is defined as total pixel intensity per image times the area of pixels positive for the signal. Total macrophage specific phosphorylated eIF2 $\alpha$  was determined by the measurement of integrated pixel intensity of phosphorylated eIF2 $\alpha$  over pixels positive for HAM56. Data is also presented for phosphorylated eIF2 $\alpha$  colocalization normalized to the area of pixels positive for HAM56 (macrophagic phosphorylated eIF2 $\alpha$  intensity per HAM56 area), to account for differences in HAM56 expression due to effects of disease.

### *Statistics*

All quantifications are expressed as mean  $\pm$  standard error of mean. Statistical comparisons were made by Student's *t*-test, one-way ANOVA plus Newman-Keuls *post hoc* test or *post hoc* test for linear trend, as indicated. All graphs were generated and statistical analyses were performed using GraphPad Prism software (San Diego, CA), and values of  $p < 0.05$  were considered significant.

**Table 3.1** Summary of case data for autopsy tissue obtained from the National Neuro-AIDS Tissue Consortium brain bank for BiP and HLA-DR staining.

<b>Case</b>	<b>HIV</b>	<b>Gender</b>	<b>Age (yrs)</b>	<b>Post mortem interval (hrs)</b>	<b>Neurocognitive diagnosis</b>
HIV1	+	M	44	8.5	HAND
HIV2	+	M	33	20	HAND
HIV3	+	M	51	5	HAND
HIV4	+	M	33	6	HAND
HIV5	+	F	31	9	HAND
HIV6	+	M	49	12	HAND
HIV7	+	N/A	46	2.75	Normal
HIV8	+	N/A	45	13	Normal
HIV9	+	N/A	44	16.4	Normal
HIV10	+	N/A	34	5	HAND
HIV11	+	N/A	32	14	HAND
HIV12	+	N/A	57	5.5	HAND
Control1	-	M	44	21.5	Normal
Control2	-	M	52	17.5	Normal



**Table 3.2** Summary of case data for autopsy tissue obtained from the National Neuro-AIDS Tissue Consortium brain bank for phosphorylated eIF2 $\alpha$  and HAM56 staining.

Case	HIV	Gender	Age (yrs)	Post mortem interval (hrs)	Neurocognitive diagnosis
HIV1	+	M	35	9.25	HAND
HIV2	+	F	34	5	HAND
HIV3	+	M	57	5.5	HAND
HIV4	+	M	32	14.5	HAND
HIV5	+	M	38	5.5	HAND
HIV6	+	M	32	14	HAND
HIV7	+	M	36	2.5	HAND
HIV8	+	M	37	11.5	HAND
HIV9	+	M	31	8.83	HAND
HIV10	+	M	49	67.33	HAND
HIV11	+	M	42	27.33	HAND
HIV12	+	M	50	21	HAND
HIV13	+	M	43	unknown	HAND
HIV14	+	M	50	18	Normal
HIV15	+	M	46	2.75	Normal
Control1	-	M	46	27.65	Normal
Control2	-	F	47	19.18	Normal
Control3	-	M	40	14.66	Normal
Control4	-	F	51	21.75	Normal

## CHAPTER 4

### THE ANTIOXIDANT RESPONSE ATTENUATES HIV REPLICATION AND MACROPHAGE-MEDIATED NEUROTOXICITY<sup>2</sup>

---

<sup>2</sup> This work was originally published in *The Journal of Immunology*. Cross SA, Cook DR, Chi AWS, Vance PJ, Kolson LL, Wong BJ, Jordan-Sciutto KL, Kolson DL. 2011. Dimethyl fumarate, an immune modulator and inducer of the antioxidant response, suppresses HIV replication and macrophage-mediated neurotoxicity; a novel candidate for HIV-neuroprotection. *J. Immunol.* Vol. 187 (10): pp-pp. Copyright © 2011 The American Association of Immunologists, Inc.

## **Abstract**

Despite antiretroviral therapy (ART), HIV infection promotes cognitive dysfunction and neurodegeneration through persistent inflammation and neurotoxin release from infected and/or activated macrophages/microglia. Furthermore, inflammation and immune activation within both the central nervous system (CNS) and periphery correlate with disease progression and morbidity in ART-treated individuals. Accordingly, drugs targeting these pathological processes in the CNS and systemic compartments are needed for effective, adjunctive therapy. Using our *in vitro* model of HIV-mediated neurotoxicity, in which HIV infected monocyte-derived macrophages (MDM) release excitatory neurotoxins, we show that HIV infection dysregulates the macrophage antioxidant response and reduces levels of heme oxygenase-1 (HO-1). Furthermore, restoration of HO-1 expression in HIV-infected MDM reduces neurotoxin release without altering HIV replication. Given these novel observations, we have identified dimethyl fumarate (DMF), used to treat psoriasis and showing promising results in clinical trials for multiple sclerosis, as a potential neuroprotectant and HIV disease-modifying agent. DMF, an immune modulator and inducer of the antioxidant response, suppresses HIV replication and neurotoxin release. Two distinct mechanisms are proposed; inhibition of NF- $\kappa$ B nuclear translocation and signaling, which could contribute to the suppression of HIV replication, and induction of HO-1, which is associated with decreased neurotoxin release. Finally, we found that DMF attenuates CCL2-induced monocyte chemotaxis, suggesting that DMF could decrease recruitment of activated monocytes to the CNS in response to inflammatory mediators. We propose that dysregulation of the antioxidant response during HIV infection drives macrophage-mediated neurotoxicity and that DMF could serve as an adjunctive neuroprotectant and HIV disease modifier in ART-treated individuals.

## Introduction

HIV-1 infection of the central nervous system (CNS) can result in cognitive, motor, and behavioral abnormalities, collectively known as HIV-associated neurocognitive disorders (HAND) (Antinori et al., 2007; McArthur et al., 2010). Early in the course of infection, HIV traffics into the brain via infected monocytes and lymphocytes (Dunfee et al., 2006) and despite antiretroviral therapy (ART) persists in parenchymal microglia and perivascular macrophages (Ho et al., 1985; Koenig et al., 1986; Petito et al., 1986). HIV infection of the CNS results in the immune activation of resident glia, and because HIV cannot infect neurons, neuronal damage is mediated by neurotoxins released by these infected and/or activated macrophages, microglia and astrocytes. Although the severity of HAND has been significantly reduced through the widespread use of ART, the prevalence and associated morbidity remain high (~50%) (Robertson et al., 2007; Sacktor et al., 2002). The persistence of HAND in individuals effectively controlled for systemic viral load is incompletely explained, although recent evidence suggests that prolonged inflammation in both the CNS and periphery may be responsible (Ancuta et al., 2008; Brenchley et al., 2006b; Eden et al., 2007).

Chronic systemic inflammation is tightly linked to morbidity and mortality in ART-treated patients, which suggests that adjunctive anti-inflammatory drugs or immune modulators may improve clinical outcomes. Despite undetectable plasma viral loads, measures of systemic inflammation correlate to cerebral spinal fluid (CSF) immune activation, CNS inflammation and HAND (Ancuta et al., 2008; Brenchley et al., 2006b; Eden et al., 2007). It has been proposed that elevated peripheral inflammation mediates neurocognitive decline by increasing the transendothelial migration of infected and/or activated monocytes into the brain (Ancuta et al., 2008; Persidsky et al., 1997). An increased number of microglia and macrophages in the CNS correlates with the severity of pre-mortem HAND, demonstrating the importance of these cell types in mediating neurological impairment (Anthony et al., 2005; Glass et al., 1995; Petito et al., 1986). Some of the most striking evidence linking peripheral inflammation to HAND derives from the strong

association between early and persistent damage caused to gut-associated lymphoid tissue (GALT) by HIV infection (or SIV infection in macaques), increased microbial translocation, systemic immune/monocyte activation and HAND progression (Ancuta et al., 2008; Brenchley and Douek, 2008; Brenchley et al., 2006a; Brenchley et al., 2006b). Therefore, reducing inflammation in the periphery as well as within the CNS is expected to improve neurocognitive impairment in HIV-infected patients.

Fumaric acid esters (FAEs), including dimethyl fumarate (DMF) and its primary *in vivo* metabolite monomethyl fumarate (MMF), are a class of compounds that have anti-inflammatory and immune modulating effects *in vitro* and *in vivo*. Fumaderm, a formulation of DMF and other FAEs, has been used in Europe since 1995 as an effective treatment for psoriasis; its mechanism of action is attributed to modulation of T cell activation and infiltration into plaques (Hoxtermann et al., 1998). DMF is currently under investigation for use in multiple sclerosis (MS) and a recently completed Phase III study demonstrated a significant benefit in suppressing relapses, disease progression and brain lesion inflammation (2011). Using the rodent model of MS, experimental allergic encephalomyelitis (EAE), it was shown that DMF reduces the recruitment of monocytes into areas of active demyelination in the brain (Schilling et al., 2006). In *in vitro* model systems, DMF has been shown to inhibit pro-inflammatory cytokine production and NF- $\kappa$ B signaling via inhibition of nuclear translocation (Loewe et al., 2002; Schilling et al., 2006; Seidel et al., 2009; Stoof et al., 2001). Furthermore, DMF induces the expression of Nrf2-driven antioxidant response genes, including heme oxygenase-1 (HO-1) and NAD(P)H quinone oxidoreductase 1 (NQO1) (Lehmann et al., 2007; Linker et al., 2011). Notably, induction of HO-1 expression in human monocytes by hemin has been associated with suppression of HIV-1 replication (Devadas and Dhawan, 2006).


Because HIV replication can be strongly driven by NF- $\kappa$ B activation and nuclear translocation, we hypothesized that DMF treatment of HIV-infected monocyte-derived macrophages (HIV/MDM)

would result in attenuation of HIV replication, immune activation and neurotoxin production. Our *in vitro* system models macrophage-mediated neurotoxicity during HIV infection by utilizing human MDM and rat cerebrocortical neuronal cultures. In this system, HIV infection of MDM results in the release of low molecular weight excitotoxins that injure neurons through excessive activation of N-methyl-D-aspartate (NMDA) receptors (Kaul et al., 2005; Lipton, 2004; O'Donnell et al., 2006). In this study we demonstrate that DMF attenuates HIV replication, nuclear translocation of NF- $\kappa$ B subunits and TNF $\alpha$  production in human MDM. Furthermore, supernatants from DMF and MMF-treated HIV/MDM cultures are markedly less neurotoxic to primary neurons than those from non-treated HIV/MDM cultures. Suppression of neurotoxin production is mediated by induction of HO-1 in HIV/MDM, and this suppression of neurotoxin production can occur even without suppression of HIV replication. Finally, DMF and MMF also reduce CCL2-induced chemotaxis in human monocytes. This study demonstrates that DMF inhibits key steps in HAND pathogenesis through distinct effects on HIV replication and macrophage-mediated neurotoxin production and DMF should be considered as an adjunctive therapeutic for ameliorating the neurological complications of HIV infection.

## Results


### *DMF and MMF inhibit HIV replication in human MDM*

Dimethyl fumarate (DMF), and its *in vivo* primary metabolite monomethyl fumarate (MMF), inhibit NF- $\kappa$ B signaling, suppress the production of inflammatory mediators and induce an antioxidant response in a variety of cell types (Lehmann et al., 2007; Loewe et al., 2002; Schilling et al., 2006; Seidel et al., 2009; Stoof et al., 2001; Vandermeeren et al., 1997; Vandermeeren et al., 2001). NF- $\kappa$ B signaling has been established as a major pathway of HIV transcriptional regulation, and recent studies have implicated the antioxidant response enzyme, HO-1, as a negative regulator of HIV replication in monocytes (Devadas and Dhawan, 2006; Devadas et al., 2010). Therefore, we hypothesized that DMF could modulate HIV replication in human macrophages through one or both of these mechanisms. Human MDM were treated with DMF or MMF and then examined for virus replication. As shown in Figure 4.1, exposure of MDM to DMF (A) or MMF (B) attenuated HIV replication in a dose-dependent manner, as determined by culture supernatant reverse transcriptase levels. Suppression of replication in MDM was seen with the R5 CSF HIV strain, Jago (Figure 4.1) and the prototypic R5/X4 strain, 89.6 (Figure 4.2). As shown in Table 4.1, HIV replication was inhibited by an average of approximately 30% at MMF concentrations achieved *in vivo* after single dose administration (4.4 $\mu$ M in CSF and 6.5 $\mu$ M in plasma) (Linker et al., 2011; Litjens et al., 2004). No drug toxicity was detected at concentrations up to 100 $\mu$ M in HIV-infected MDM (HIV/MDM) (Figure 4.1C, 1D) and non-infected MDM (data not shown). DMF demonstrated additive effects in attenuating HIV replication when used in combination with efavirenz, a non-nucleoside reverse transcriptase inhibitor (Figure 4.3A). There was no observed cellular toxicity when DMF was used in combination with efavirenz (Figure 4.3B).



**Figure 4.1 Dimethyl fumarate and monomethyl fumarate attenuate HIV replication in human MDM.** Human MDM infected with 50ng HIV (p24 ELISA, equivalent to  $1.82 \pm 0.22$  kcpm/ $\mu$ L by reverse transcriptase (RT) activity assay) were treated with DMF (**A**) or MMF (**B**) over the course of infection at the indicated concentrations (1-30 $\mu$ M) or with 20nM of the non-nucleoside reverse transcriptase inhibitor, efavirenz (EFZ). Culture supernatants were collected every 2-3 days, as indicated, and HIV replication was quantified by RT activity. **C**) DMF and **D**) MMF cause no cytotoxicity in HIV/MDM as assessed by LDH assay of supernatants harvested at day 14 post infection. Maximum (Max) LDH release represents the soluble LDH release following cell lysis. RT curves are representative of 3-4 independent experiments, with each replicate performed on cell preparations from different donors. LDH assays represent data averaged from 3-5 individual donors. All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing, \*\*\* $p < 0.001$  vs. EFZ.





**Figure 4.2 Dimethyl fumarate attenuates 89.6 HIV replication in human MDM.** Human MDM infected with HIV-89.6 were treated with DMF (A) or MMF (B) over the course of infection at the indicated concentrations (1- 30μM) or with the non-nucleoside reverse transcriptase inhibitor, efavirenz (EFZ). Culture supernatants were collected every 2-4 days, as indicated, and assessed for RT activity. Values indicate the mean  $\pm$  SEM. RT curves are representative of 2-3 independent experiments, with each replicate performed on cell preparations from different donors.


**Table 4.1.** Reduction in RT activity, relative to vehicle control, with increasing concentrations of DMF and MMF among individual donors.

<b>Donor #</b>	<b>5<math>\mu</math>M DMF</b>	<b>15<math>\mu</math>M DMF</b>	<b>30<math>\mu</math>M DMF</b>
<b>5</b>	16.7 $\pm$ 11.0% n =4	27.8 $\pm$ 9.8% n =3	64.0 $\pm$ 13.4% n =5
<b>36</b>	10.0 $\pm$ 13.1% n =3	68.0 $\pm$ 3.6% n =1	52.7 $\pm$ 15.4% n =3
<b>40</b>	17.1 $\pm$ 6.7% n =2	66.0 $\pm$ 5.0% n =1	73.4 $\pm$ 20.9% n =2
<b>7</b>	59.6 $\pm$ 5.0% n =2	65.2 $\pm$ 9.2% n =1	60.6 $\pm$ 5.7% n =2
<b>26</b>	81.5 $\pm$ 12.4% n =1	91.3 $\pm$ 6.7% n =1	51.2 $\pm$ 15.1% n =3
<b>38</b>	38.7 $\pm$ 4.3% n =1	42.3 $\pm$ 4.2% n =1	53.9 $\pm$ 6.0% n =2
<b>AVG</b>	<b>26.5 <math>\pm</math> 36.0%</b> <b>n =20 (12)</b>	<b>45.5 <math>\pm</math> 26.4%</b> <b>n =13 (11)</b>	<b>55.0 <math>\pm</math> 32.3%</b> <b>n =25 (13)</b>

<b>Donor #</b>	<b>5<math>\mu</math>M MMF</b>	<b>15<math>\mu</math>M MMF</b>	<b>30<math>\mu</math>M MMF</b>
<b>33</b>	29.5 $\pm$ 19.2% n =3	36.7 $\pm$ 14.6% n =3	66.1 $\pm$ 13.9% n =3
<b>40</b>	14.7 $\pm$ 21.5% n =2	57.0 $\pm$ 8.3% n =2	61.6 $\pm$ 8.3% n =2
<b>38</b>	29.5 $\pm$ 9.9% n =2	36.7 $\pm$ 2.9% n =1	66.1 $\pm$ 3.7% n =2
<b>29</b>	33.6 $\pm$ 93.9% n =2	76.8 $\pm$ 4.3% n =1	86.9 $\pm$ 1.7% n =1
<b>AVG</b>	<b>28.6 <math>\pm</math> 32.4%</b> <b>n =12 (7)</b>	<b>47.8 <math>\pm</math> 27.1%</b> <b>n =10 (7)</b>	<b>67.5 <math>\pm</math> 20.5%</b> <b>n =12 (7)</b>


Values were calculated from RT data from the peak of infection and are presented as mean  $\pm$  SD. For combined average (AVG) data, n = number of replicates averaged (number of unique donors). Only individual donors assayed more than once are included in the table.



**Figure 4.3 Dimethyl fumarate and efavirenz have additive effects on attenuation of HIV replication and neurotoxin production in human MDM.** **A)** DMF and efavirenz (EFZ), a non-nucleoside reverse transcriptase inhibitor, have additive effects on the attenuation of HIV replication in human MDM. MDM were infected with HIV-Jago and treated with DMF and/or EFZ over the course of infection. Culture supernatants were collected every 3 days and assessed for RT activity. Values indicate the mean  $\pm$  SEM. RT curves are representative of 2 independent experiments, with each replicate performed on cell preparations from a different donor. **B)** DMF and EFZ cause no cytotoxicity in HIV/MDM as assessed by LDH assay of supernatants harvested at day 15 post infection. Maximum (Max) LDH release represents the soluble LDH release following cell lysis. LDH assays represent data averaged from 2 individual donors. **C)** DMF and EFZ used in combination reduce macrophage-mediated neurotoxicity to a greater extent than either drug alone. Neuronal survival was assessed by MAP2 ELISA following exposure to supernatant of HIV-infected macrophages, which were treated with DMF and/or EFZ over the course of infection. MAP2 data is expressed as a percentage of untreated (UT) cultures ( $n = 6$ ). All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing, \*\* $p < 0.01$  and \*\*\* $p < 0.001$  vs. Vehicle.

#### *DMF and MMF reduce HIV/MDM-mediated neurotoxicity*

We and others have shown that HIV-infected MDM release potent neurotoxins that injure neurons through over-activation of N-methyl-D-aspartate receptors (NMDAR) and that this excitotoxicity is mediated by glutamate and other low molecular weight NMDAR agonists (Chen et al., 2002; Jiang et al., 2001; O'Donnell et al., 2006). Although the mechanisms underlying neurotoxin production in HIV/MDM are not fully understood, suppression of HIV replication in MDM generally suppresses such neurotoxicity, as demonstrated by treatment with efavirenz (Figure 4.4). Similarly, in addition to suppressing HIV replication (Figure 4.1), DMF (Figure 4.4A) and MMF (Figure 4.4B) also reduce HIV/MDM neurotoxin production in a dose-dependent manner, as assessed by neuronal survival in our *in vitro* HIV neurotoxicity model. Representative images of HIV/MDM-mediated neurotoxicity and the protective effects of DMF and MMF are shown (Figure 4.4C), where surviving neurons are labeled for MAP2 (microtubule-associated protein 2). DMF and EFZ used in combination resulted in additive effects on the suppression of macrophage-mediated neurotoxicity (Figure 4.3C), demonstrating that DMF may successfully reduce HIV replication and macrophage-mediated neurotoxicity that is not fully suppressed by ART. This neuroprotection is due to drug effects on the macrophages, as DMF and MMF do not prevent HIV/MDM mediated neurotoxicity when applied directly to the neurons prior to addition of HIV/MDM supernatants (data not shown).




**Figure 4.4 DMF and MMF reduce HIV/MDM mediated neurotoxicity.** Rat cerebrocortical cultures were exposed to supernatant from HIV-infected macrophages that were treated with DMF (**A**) or MMF (**B**) at the indicated concentrations (1-30 $\mu$ M) during the course of infection. Neuronal survival was assessed by MAP2 ELISA and expressed as a percentage of untreated (UT) cultures (n = 6; \*\*\*p<0.001 vs. Vehicle). **C**) Representative images of rat cerebrocortical cultures immunofluorescently stained for MAP2 (red) and Hoescht 33324 (blue) following 24 hours treatment with the indicated HIV/MDM supernatant. Scale bar represents 50 $\mu$ m. All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing.

#### *DMF inhibits NF- $\kappa$ B nuclear entry, DNA binding and TNF $\alpha$ production in human MDM*

NF- $\kappa$ B and TNF $\alpha$  are part of a positive feedback loop that regulates the transcriptional activity of the HIV long terminal repeat (LTR). In unstimulated cells, NF- $\kappa$ B is unable to bind DNA due to its association with inhibitory  $\kappa$ B (I $\kappa$ B) proteins, which sequester NF- $\kappa$ B in the cytoplasmic compartment (Baldwin, 1996; Ganchi et al., 1992; Henkel et al., 1992). Following exposure to an activating stimulus such as TNF $\alpha$ , NF- $\kappa$ B is rapidly freed from the inhibitory complex and translocates into the nucleus to induce transcriptional activation of viral and host genes. NF- $\kappa$ B proteins are major modulators of the HIV LTR and are among the most potent activators of

proinflammatory and inflammatory genes. Five members of the mammalian NF- $\kappa$ B/Rel family have been described, including c-Rel, NF- $\kappa$ B1 (p50/p105), NF- $\kappa$ B2 (p52/p100), RelA (p65), and RelB. Functional NF- $\kappa$ B complexes are composed of heterodimer complexes containing p65, c-Rel, or RelB bound to p50 or p52 (Baldwin, 1996; Kilariski et al., 2009; Neumann et al., 2000). Exposure to activating stimuli, such as TNF $\alpha$ , induces the nuclear accumulation of NF- $\kappa$ B proteins, DNA binding by NF- $\kappa$ B p50 and transcription from the HIV LTR (Duh et al., 1989).

To determine if DMF and MMF inhibit the nuclear translocation of NF- $\kappa$ B proteins in MDM, DMF and MMF-treated MDM were stimulated with TNF $\alpha$  and subjected to subcellular fractionations before detection of NF- $\kappa$ B subunits by Western blotting. DMF and MMF each inhibited TNF $\alpha$ -induced nuclear accumulation of RelB, p65 and p50 in a dose-dependent manner (Figure 4.5A, 3B). We also demonstrate that DMF inhibited the formation of the NF- $\kappa$ B p50-DNA complex, as assessed by EMSA (Figure 4.5C). Because NF- $\kappa$ B signaling also induces expression of inflammatory mediators, we assessed the effects of DMF treatment on TNF $\alpha$  release from MDM. In agreement with previous reports of DMF decreasing the release of inflammatory mediators from multiple cell types, including TNF $\alpha$ , IL-1 $\beta$  and IL-6, (Lehmann et al., 2007; Wilms et al., 2010) we found that DMF suppresses release of TNF $\alpha$  from PHA-activated MDM (Figure 4.5D). Furthermore, DMF also markedly suppressed HIV-induced TNF $\alpha$  release from MDM (Figure 4.5E). Thus, DMF and its primary metabolite, MMF, inhibit NF- $\kappa$ B translocation and signaling events that contribute to the positive feedback loop that modulates HIV transcription in infected and activated MDM.



**Figure 4.5 DMF inhibits NF-κB nuclear translocation, DNA binding and TNFα production in human MDM.** **A)** DMF and **B)** MMF inhibit the nuclear translocation of the NF-κB proteins RelB, p65 and p50 in human MDM in a dose-dependent manner. Cells were treated with DMF or MMF for 24 hours, exposed to TNFα (10 min), separated into cytoplasmic and nuclear fractions and analyzed by Western blotting. Results of densitometry analysis are presented numerically under each panel as the ratio of NF-κB protein to PARP, a nuclear marker and loading control. Blots are representative of 4-6 independent experiments, with each replicate performed on cell preparations from different donors. **C)** DMF inhibits nuclear NF-κB p50 binding to DNA in TNFα stimulated MDM, as assessed by EMSA. Results of densitometry analysis were normalized to vehicle. **D)** DMF inhibits the production of TNFα in MDM stimulated with PHA (10μg/mL). Values are expressed as percent TNFα production relative to Vehicle treated cells (227 ± 11.9 pg/mL TNFα in Vehicle). Data are expressed as mean ± SEM and represent data averaged from 4 different donors. **E)** TNFα production in HIV/MDM is inhibited by DMF treatment. HIV/MDM were treated with DMF (1-30μM) or 20nM efavirenz (EFZ) over the course of infection and culture supernatants from day 14-15 post infection were assayed for TNFα by ELISA. Values represent the mean ± SEM of data averaged from 5 different donors. All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing (\*p<0.05, \*\*p<0.01, \*\*\*p<0.01 vs. Vehicle).


*DMF restores the antioxidant response suppressed by HIV infection in MDM*

The antioxidant response is one of the cellular adaptive stress responses that can modulate virus replication and host cell survival, as shown in Hepatitis B and Dengue 2 infection models (Chen et al., 2011; Schaedler et al., 2010). The antioxidant response maintains redox balance and counteracts oxidative damage through induction of proteins that are involved in detoxification of reactive oxygen species (ROS). These proteins are produced by genes with a common promoter element, the antioxidant response element (ARE), and ARE transcription is mediated by Nrf2. Under conditions of low oxidative stress, Nrf2 is kept transcriptionally inactive by Kelch-like ECH-associated protein 1 (Keap1), which sequesters Nrf2 in the cytoplasmic compartment (Itoh et al., 1999). Following exposure to ROS or electrophiles, Keap1 is degraded by the proteasome and Nrf2 translocates to the nucleus to drive expression of numerous genes including, HO-1, NQO1, glutathione peroxidase 1 (GPX1), and genes responsible for glutathione synthesis (glutamate cysteine ligase modifier, glutamate cysteine ligase catalytic subunit and glutathione synthetase). HIV infection is associated with increased ROS production and depressed levels of glutathione, the major intracellular antioxidant (Dworkin et al., 1986). We observed a marked reduction in the level of HO-1 expression in HIV/MDM across multiple human donors, with a more modest but nonetheless consistent reduction in GPX1 levels (Figure 4.6A, 4B). The effects of DMF on Nrf2 and NQO1 levels were more variable among HIV/MDM cultures from different donors, but trended towards increased expression relative to uninfected Mock/MDM (Figure 4.6A, 4B).

Upon exposure of HIV/MDM to DMF, expression of Nrf2, HO-1, GPX1 and NQO1 increased with increasing doses of DMF (Figure 4.6C), suggesting a restoration of antioxidant responses in HIV-infected MDM. Both HIV infection and DMF increase total levels of Nrf2, suggesting that while HIV infection stabilizes or induces total cellular Nrf2 levels, this is not sufficient for the coordinated transcriptional activation of ARE-regulated genes, such as HO-1 and GPX1. DMF and MMF treatment activates transcription of these ARE-regulated genes in HIV-infected macrophages, possibly by disrupting inhibitory Nrf2-Keap1 interactions (Linker et al., 2011). DMF restores levels




of HO-1 and GPX1 to those observed in uninfected MDM, while NQO1, which is not suppressed during HIV infection, is induced to levels exceeding those in uninfected MDM (Figure 4.6D). We have also confirmed that MMF can induce the antioxidant response in HIV/MDM (Figure 4.6E, 4F) and that both DMF and MMF induce the antioxidant response in uninfected MDM (data not shown). DMF induction of antioxidant responses in MDM occurs independently of HIV infection, which is consistent with previous findings describing induction of the antioxidant response by DMF in multiple cell types, including glia and neurons (Lehmann et al., 2007; Linker et al., 2011).



**Figure 4.6 DMF restores the imbalance in the antioxidant response caused by HIV infection.** HIV infection of human MDM reduces HO-1 and GPX1 expression, as assessed by Western blotting (**A**) and quantified by densitometry analysis (**B**). Values indicate mean  $\pm$  SEM of 6 different donors. Statistical comparisons were made by two-tailed paired t-test (\* $p$ <0.05, \*\* $p$ <0.01 and \*\*\* $p$ <0.001 vs. Mock). **C**) DMF activates the Nrf2-dependent antioxidant response in HIV/MDM and restores HO-1 and GPX1 levels to that found in uninfected Mock cells, as quantified by densitometry analysis (**D**). **E**) MMF activates the Nrf2-dependent antioxidant response in HIV/MDM and restores HO-1 and GPX1 levels to that found in uninfected Mock cells, as quantified by densitometry analysis (**F**). Blots are representative of 3 independent experiments, with each replicate performed on cell preparations from different donors. Densitometry data are expressed as mean  $\pm$  SEM and represent data averaged from 3 different donors.

*DMF inhibition of HIV replication and NF- $\kappa$ B signaling is not mediated by HO-1*


HIV infection of human MDM results in alterations to the antioxidant response with a striking reduction in HO-1 levels (Figure 4.6A, 4.6B). Induction of HO-1 by hemin has been reported to decrease HIV replication in human monocytes, suggesting that DMF's induction of HO-1 may underlie its antiviral effects (Devadas and Dhawan, 2006). We used a pharmacologic inhibitor of HO-1 enzymatic activity, tin mesoporphyrin (SnMP), to determine the potential role for HO-1 in DMF-mediated suppression of HIV replication and NF- $\kappa$ B translocation. As shown in Figure 4.7A, SnMP had no effect on DMF-mediated HIV suppression, which suggests that DMF does not suppress HIV replication through enhanced HO-1 expression and activity. We found no effect of SnMP on DMF-mediated suppression of HIV replication regardless of donor, level of infection, DMF dose or timing of SnMP addition (data not shown). We also confirmed that SnMP does not inhibit DMF's suppression of TNF $\alpha$ -induced nuclear accumulation of NF- $\kappa$ B (Figure 4.7B). In addition, we show that an inducer of HO-1 expression, cobalt protoporphyrin (CoPP) had no effect on TNF $\alpha$ -induced nuclear accumulation of NF- $\kappa$ B (Figure 4.7B). These results suggest that DMF's induction of HO-1 does not directly suppress HIV replication or NF- $\kappa$ B signaling.



**Figure 4.7 HO-1 does not mediate the attenuation of HIV replication or NF-κB signaling induced by DMF. A)** SnMP, an inhibitor of HO-1 enzymatic activity, does not inhibit DMF-mediated attenuation of HIV replication. DMF and vehicle treated HIV/MDM were exposed to 10μM SnMP from day 6 through day 15 post infection. Culture supernatants were collected every 3 days and assessed for RT activity. RT curves are representative of 3 independent experiments, with each replicate performed on cell preparations from different donors. **B)** SnMP and CoPP, a specific inducer of HO-1, do not directly affect or alter DMF-mediated inhibition of the nuclear translocation of NF-κB proteins, as assessed by western blotting. Human MDM were treated with 10μM SnMP, 10μM CoPP and/or 100μM DMF for 24 hours before treatment with 1 ng/mL TNFα (10min) and subcellular fractionation. Western blot is representative of 3 independent experiments, with each replicate performed on cell preparations from different donors.

#### *Induction of HO-1 reduces neurotoxin production from HIV/MDM*


We sought to determine whether the suppression of HIV/MDM neurotoxin production by DMF (Figure 4.4) was associated with DMF's suppression of HIV replication and/or induction of HO-1 expression. Inhibiting HIV replication in HIV/MDM can suppress neurotoxin release *in vitro*, as demonstrated by efavirenz treatment (Figure 4.4), and similar effects of ART drugs *in vivo* are thought to account for their ability to limit the severity of HAND in ART-experienced cohorts. While previous studies found that increased HO-1 activity is associated with decreased HIV replication in MDM (Devadas and Dhawan, 2006; Devadas et al., 2010), we found that neither inhibition of HO-1 activity by SnMP treatment of MDM (Figure 4.8A) nor induction of HO-1 expression by CoPP (Figure 4.8D, 6F) altered HIV replication. Remarkably, however, SnMP treatment significantly increased the neurotoxicity of MDM supernatant (Figure 4.8B), even when HIV replication was low or absent (Figure 4.8C). The increase in MDM-mediated neurotoxicity was a consequence of inhibiting HO-1 activity in the macrophage since SnMP was not toxic when added directly onto neurons (data not shown). And while CoPP does not attenuate HIV replication or inhibit NF- $\kappa$ B signaling, supernatant from CoPP-treated HIV/MDM is significantly less neurotoxic than untreated controls with a similar level of HIV replication (Figure 4.8E). These studies demonstrate that HO-1 is a critical modulator of neurotoxin production in HIV/MDM and that HO-1 levels can modulate HIV/MDM neurotoxicity without affecting HIV replication.



**Figure 4.8 HO-1 induction reduces neurotoxin production in HIV/MDM without affecting HIV replication.** **A)** SnMP, an inhibitor of HO-1 enzymatic activity, does not directly affect HIV replication and supernatant from these SnMP treated HIV/MDM are significantly more neurotoxic (**B**), despite equal levels of HIV replication. **C)** Uninfected Mock/MDM and 20nM efaverinz (EFZ) treated HIV/MDM, which normally produce minimal neurotoxins, are significantly more neurotoxic when treated with 10μM SnMP. **D)** CoPP, a specific inducer of HO-1 expression, does not directly affect HIV replication and supernatant from these CoPP treated HIV/MDM are significantly less neurotoxic (**E**), despite high levels of virus replication. **F)** CoPP treatment exponentially increases HO-1 levels without greatly altering the other components of the antioxidant response, as assessed by Western blotting. For A and C, SnMP or CoPP was added at day 6 post infection onwards and culture supernatants were collected every 3 days and assessed for RT activity. RT curves are representative of 3 independent experiments, with each replicate performed on cell preparations from different donors. For neuronal survival assays, survival was assessed by MAP2 ELISA and expressed as a percentage of untreated (UT) cultures (n = 6; \*\*\*p<0.001 vs. vehicle treated paired-condition). Statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing. Western blot is representative of 3 independent experiments, with each replicate performed on cell preparations from different donors. Two film exposures (short and long) of HO-1 are presented to demonstrate the extent of HO-1 induction over basal levels.

*DMF and MMF inhibit CCL2 induced chemotaxis in human monocytes*

The recruitment of activated and infected monocytes to the CNS in response to CCL2 is a key step in the pathogenesis of HAND (Cinque et al., 1998; Zink et al., 2001). In a previous DMF study using the mouse EAE model, DMF reduced macrophage infiltration into the spinal cord in areas of active demyelination (Schilling et al., 2006). We hypothesized that DMF could inhibit chemotaxis of human monocytes in response to chemotactic cytokines, such as CCL2. We found that DMF and MMF inhibited chemotaxis in freshly isolated human monocytes in response to CCL2 in a dose-dependent manner (Figure 4.9A, 7B). Furthermore, we found that DMF reduced the expression of the CCL2 receptor, CCR2, in freshly isolated human CD11b<sup>+</sup>CD14<sup>+</sup> monocytes within 6 hours of treatment (Figure 4.9C, 7D), without causing death (Figure 4.9E). These results indicate that DMF and MMF can decrease monocyte chemotaxis in response to CCL2 and that this effect is associated with downregulation of CCR2 expression.



**Figure 4.9 DMF and MMF reduce CCL2 induced chemotaxis in human monocytes.** **A)** DMF and **B)** MMF inhibit CCL2-induced chemotaxis in freshly isolated human monocytes in a dose-dependent manner. Values are expressed as percent migration of unstimulated cells (US; 0 ng/mL CCL2) ( $n = 10-22$ ; \*\* $p < 0.01$  vs. Vehicle). **C)** DMF decreases CCR2 expression on CD11b<sup>+</sup>CD14<sup>+</sup> PBMCs following 6 hours of treatment, as quantified (**D**). **E)** DMF does not cause significant cell death over 6 hours of treatment in freshly isolated human monocytes, as measured by DAPI positivity in CD11b<sup>+</sup>CD14<sup>+</sup> gated PBMCs. For all experiments, values represent data averaged from 3 different donors. All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing. Results of post test for linear trend are also presented.



## Discussion

Monocytes and macrophages are major reservoirs for HIV in both the periphery and CNS, and they facilitate the spread of virus to target cells, allow for viral persistence and serve as major contributors to inflammation-mediated pathology. Despite current ART, latently infected monocytes and CD4<sup>+</sup> T-lymphocytes persist, resulting in inflammation in the periphery and in the CNS in up to 50% of patients on ART (Robertson et al., 2007; Sacktor et al., 2002). While ART will remain the mainstay of HIV therapy, effective adjunctive therapies that suppress inflammation, improve morbidity and improve long-term cognitive outcomes are greatly needed. The immunomodulator DMF, which is effective for the treatment of psoriasis and which shows promising results for multiple sclerosis treatment in recent clinical trials, is an attractive candidate as a safe adjunctive neuroprotectant against HIV. We have demonstrated that physiologically relevant doses of DMF and its primary metabolite, MMF, (Linker et al., 2011; Litjens et al., 2004) affect key steps in the pathogenesis of HAND in our *in vitro* model system by inhibiting HIV replication, neurotoxin production, NF- $\kappa$ B signaling and TNF $\alpha$  production in human MDM and reducing monocyte chemotaxis in response to CCL2. These results suggest that DMF could serve as an effective neuroprotectant in HAND and have beneficial effects on systemic HIV-disease progression as well.

We have shown that DMF and MMF attenuate macrophage-mediated neurotoxicity following HIV infection by simultaneously attenuating viral replication and inducing HO-1 expression. Furthermore, induction of HO-1 can significantly decrease macrophage-mediated neurotoxicity even without decreasing HIV replication. Consequently, DMF may be an especially relevant therapeutic in patients who have relatively good virologic control but still suffer from neurological complications of HIV. We have shown that HIV infection of MDM results in a dysregulation of the antioxidant response with an especially prominent reduction in HO-1 levels, associated with supernatant neurotoxicity, and that DMF treatment restores HO-1 levels and reduces neurotoxin production in macrophages. In activated microglia, an oxidative burst is required for the release of

excitotoxic glutamate (Barger et al., 2007), demonstrating that alterations to cellular oxidative state can mediate the production and/or release of MDM neurotoxins. DMF's ability to decrease HIV replication and neurotoxin production by distinct mechanisms makes it an especially attractive therapeutic candidate for HAND. Furthermore, macrophage- and microglia-mediated neurotoxicity contribute to many other neurological disorders including multiple sclerosis, Alzheimer's disease, amyotrophic lateral sclerosis, Huntington's disease and stroke/reperfusion injury, for which therapeutics for restoring oxidative balance resultant from the disease state have been investigated/prescribed (Uttara et al., 2009).

Numerous proinflammatory factors contribute to HIV disease pathogenesis in both the peripheral and CNS compartments. TNF $\alpha$ , IL-6, IL-1 $\beta$ , IFN- $\gamma$  and other proinflammatory cytokines are elevated in the blood and CSF of HIV-infected patients (Breen et al., 1990; Ciardi et al., 1994; Kobayashi S, 1990; Ownby et al., 2009). Among these, TNF $\alpha$  is the most potent mediator of inflammation and is induced early after HIV monocytic infection and its expression continues to increase over the course of infection (Esser et al., 1996; Folks et al., 1988; Koyanagi et al., 1988). It is well established that TNF $\alpha$  exposure upregulates HIV replication by initiating a signaling cascade that activates the nuclear translocation of NF- $\kappa$ B (Butera et al., 1993; Nabel and Baltimore, 1987; Poli et al., 1990; West et al., 2001). We have shown that DMF and MMF attenuate TNF $\alpha$ -mediated NF- $\kappa$ B signaling in human macrophages and reduce nuclear NF- $\kappa$ B levels, which are expected to decrease transcription from the HIV-LTR. However, the NF- $\kappa$ B and TNF $\alpha$  signaling loop may not entirely mediate DMF's antiviral activity. Attenuation of HIV replication occurs at low concentrations, as does induction of the ARE, while inhibition of NF- $\kappa$ B signaling may be more relevant at concentrations of 15 $\mu$ M and greater. Future studies are necessary to assess the role of the antioxidant response, including NQO1 and the cellular redox state, in mediating HIV infection and replication. DMF may alter the expression of the HIV co-receptors, CXCR4 and CCR5, similarly to the observed downregulation of cell surface CCR2. It has been reported that antioxidants decrease the stability of mRNA transcripts for CXCR4 and

CCR5 in human monocytes, suggesting that DMF treatment may directly reduce HIV entry into human monocytes (Saccani et al., 2000).

However, DMF's ability to inhibit NF- $\kappa$ B and TNF $\alpha$  signaling following both PHA stimulation and HIV infection has clear implications for the physiologic reduction of neuroinflammation and cytokine induced neuronal injury. Elevated TNF $\alpha$  levels increase monocyte entry into the brain, promote HIV replication, and drive inflammatory cascades, thereby enhancing the production of neurotoxins in the CNS from MDM, microglia and astrocytes (Fiala et al., 1997). Therefore, dampening TNF $\alpha$ -driven processes might also afford neuroprotection against HIV. Indeed, TNF $\alpha$  is linked to glutamine synthetase and glutamate import in macrophages (Porcheray et al., 2006), and DMF's inhibition of TNF $\alpha$ -driven processes may further decrease the release of excitatory neurotoxins, such as glutamate, in HIV/MDM. In human macrophages, we have shown that DMF is a potent suppressor of NF- $\kappa$ B nuclear translocation, subsequent binding to DNA and expression of NF- $\kappa$ B dependent genes. Therefore, DMF is a particularly good therapeutic candidate for pathological states characterized by macrophage driven inflammation and NF- $\kappa$ B signaling.

Although not directly dependent upon HO-1, DMF's antioxidant properties are likely mediating the inhibition of NF- $\kappa$ B activity. We hypothesize that such effects are due to DMF's modulation of the macrophage intracellular redox state as activation of the antioxidant response has been shown to block NF- $\kappa$ B activity and HIV transcription (Roederer et al., 1990; Schreck R, 1991; Staal FJ, 1993). Furthermore, classical ( $\alpha$  and  $\beta$ ), novel ( $\delta$ ) and atypical ( $\zeta$ ) PKC isotypes can modulate the nuclear translocation and transcriptional activity of NF- $\kappa$ B and PKC is activated by oxidative stress and inhibited by antioxidants (Asehnoune et al., 2005; Boscoboinik et al., 1991; Gopalakrishna et al., 1995; Gopalakrishna et al., 1997; Sun et al., 2000). In addition to potential effects on PKC, DMF may also affect the phosphorylation of I $\kappa$ B kinases (IKK) and subsequent phosphorylation and degradation of I $\kappa$ B proteins (Seidel et al., 2009; Vandermeeren et al., 2001).

Finally, DMF may affect NF- $\kappa$ B dependent transcription by modulating the preferred composition of NF- $\kappa$ B homo- and heterodimers that form after nuclear translocation has occurred. The intracellular oxidative state can affect levels of NF- $\kappa$ B p50 homodimers, which do not possess transactivation domains and are thought to act as transcriptional repressors of NF- $\kappa$ B heterodimer responsive genes (Cristofanon et al., 2009; Hoberg et al., 2006; Zhou et al., 2001). We are currently examining the role of DMF and MMF in modulating the activation state of the macrophage, which would affect the cell's relative sensitivity to pro-inflammatory signals and thereby contribute to decreased NF- $\kappa$ B signaling.

While other antioxidants have been considered as potential therapeutics for HAND, through direct effects on macrophages or neurons, DMF is unique in its ability to inhibit CCL2-induced monocyte chemotaxis. Monocyte transmigration across the blood-brain barrier is dependent upon production of chemokines, such as CCL2, in the CNS and the activation of monocytes in the periphery. Levels of CCL2 in the CSF correlate with CSF viral load and with the clinical severity of HAND (Cinque et al., 1998; Conant et al., 1998; Kelder et al., 1998; Letendre et al., 1999; Sozzani et al., 1997; Zink et al., 2001), and CCL2 is produced by brain macrophages, astrocytes and endothelial cells in response to inflammatory mediators and HIV proteins (Gu et al., 1997; Guillemin et al., 2003; Lehmann et al., 2006). Not only does DMF decrease TNF $\alpha$  production and NF- $\kappa$ B signaling in MDM, both of which have been implicated in CCL2 production, but DMF and MMF inhibit CCL2-driven monocyte chemotaxis, possibly by modulation of CCR2 expression. DMF and MMF may modulate the cell surface expression of CCR2 by inducing the antioxidant response and consequently altering the redox state of the cell. It has been demonstrated that direct antioxidants are capable of reducing the transcript stability of CCR2, which has been linked to decreased cell surface expression and CCL2-induced chemotaxis in human monocytes (Saccani et al., 2000). These findings in our *in vitro* model system predict suppression of transendothelial migration of monocytes into the CNS during HIV infection. Furthermore, it has been reported that DMF modulates adhesion molecule expression in human endothelial cells by


inhibiting TNF $\alpha$ -induced expression of ICAM-1, VCAM-1 and E-selectin (Vandermeeren et al., 1997). Expression of each of these adhesion molecules has been linked to monocyte entry into the CNS after HIV infection and down-regulation by DMF is expected to further inhibit monocyte entry into the CNS. Given these findings, DMF should be considered as a potential therapeutic for other neuroinflammatory diseases associated with CCL2-induced recruitment of leukocytes to the CNS.

With this study, we identify dimethyl fumarate as a candidate adjunctive therapy and potential neuroprotectant against HIV. To our knowledge, we are the first to demonstrate that HIV infection dysregulates components of the antioxidant response in human macrophages and that restoration of HO-1 levels, specifically, can reduce macrophage-mediated neurotoxicity. DMF is the first proposed neuroprotectant that reduces CCL2-mediated monocyte chemotaxis as a component of its mechanism of action. Furthermore, we have shown that DMF attenuates HIV replication associated with decreased TNF $\alpha$  and NF- $\kappa$ B signaling. Given these findings, we propose that DMF should be considered a relevant therapeutic candidate for neurological disorders and other complications of HIV-infection mediated by monocyte and macrophage inflammation.

## Work in Progress

### *Induction of the antioxidant response with tBHQ attenuates HIV replication, macrophage-mediated neurotoxicity and NF- $\kappa$ B signaling*

We have demonstrated that DMF and MMF treatment attenuate macrophage-mediated neurotoxicity in a dose-dependent manner and that these effects are due to induction of the antioxidant response, specifically restoration of HO-1 levels (Cross et al., 2011). We also hypothesized that induction of the antioxidant response and alteration of the cellular redox state is responsible for DMF and MMF's inhibition of NF- $\kappa$ B nuclear translocation following TNF $\alpha$  exposure and PHA-induced TNF $\alpha$  production in uninfected macrophages. tBHQ (*tert*-butylhydroquinone) is a well characterized and potent inducer of the antioxidant response. tBHQ possesses an oxidizable 1,4 diphenolic structure that dissociates the Keap1-Nrf2 complex, inducing Nrf2 translocation to the nucleus and transcription of ARE-regulated genes (De Long et al., 1987; Talalay, 1989; van Ommen et al., 1992). As shown in Figure 4.10, tBHQ treatment results in a robust increase in HO-1 protein levels, with a more moderate but significant increase in NQO1. And similar to DMF/MMF treatment, tBHQ treatment also results in a dose-dependent decrease in HIV replication in human macrophages with concomitant decreases macrophage-mediated neurotoxicity. Furthermore, tBHQ inhibits the nuclear translocation of NF- $\kappa$ B proteins following TNF $\alpha$  exposure and TNF $\alpha$  release following PHA stimulation (Figure 4.10D and E). tBHQ, a classic activator of the antioxidant response, has the same antiviral and anti-inflammatory properties as DMF and MMF treatment, supporting the hypothesis that alterations to the cellular redox state via activation of the antioxidant response are mediating the observed effects. These results also suggest that other pharmacologics that restore the dysregulation of the antioxidant response in HIV-infected macrophages should be considered as adjunctive therapies for HAND. Ongoing experiments are investigating the effect of tBHQ treatment on CCR2 expression and CCL2-induced monocyte chemotaxis.




**Figure 4.10 tBHQ, an inducer of the antioxidant response, attenuates HIV replication and macrophage-mediated neurotoxicity.** **A)** tBHQ attenuates HIV replication in a dose-dependent manner. Human MDM infected with HIV-Jago were treated with tBHQ over the course of infection. Culture supernatants were collected every 3 days and assessed for RT activity. RT curves are representative of 2 independent experiments, with each replicate performed on cell preparations from a different donor. **B)** tBHQ reduces macrophage-mediated neurotoxicity following HIV infection. Neuronal survival was assessed by MAP2 ELISA following exposure of rat cerebrocortical neuronal cultures to supernatant of HIV-infected macrophages treated with tBHQ over the course of infection. MAP2 data is expressed as a percentage of untreated (UT) cultures (n = 6). **C)** tBHQ induces the antioxidant response in uninfected human MDM as assessed by Western blotting. **D)** Similarly to DMF and MMF, tBHQ inhibits the nuclear translocation of NF-κB proteins RelB, p65 and p50 in human MDM. Cells were treated with 100 μM tBHQ for 24 hours, exposed to TNFα (10 min), separated into nuclear fractions and analyzed by Western blotting. Blots are representative of 3 independent experiments, with each replicate performed on cell preparations from different donors. **E)** tBHQ inhibits the production of TNFα in MDM stimulated with PHA (10 μg/mL) for 24 hours. Values represent data averaged from 3 different donors. Data are expressed as mean ± SEM. All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing, \*\*\*p<0.001 vs. Vehicle.

*Induction of HO-1 expression with hemin attenuates macrophage-mediated neurotoxicity without altering HIV replication*

Previous reports have associated hemin treatment, an inducer of HO-1, with attenuation of HIV replication (Devadas and Dhawan, 2006). However, we were unable to replicate these previously published results. As shown in Figure 4.11, hemin treatment of HIV-infected macrophages had no effect on viral replication, although it did induce levels of HO-1 above those seen in EFZ-treated controls. However, the extent of HO-1 induction following 20 $\mu$ M hemin treatment was several-fold less than that achieved with 100 $\mu$ M DMF treatment (Figure 4.11B). Hemin treatment also resulted in a small increase in NQO1 levels in HIV/MDM and previously published studies suggest that hemin is capable of activating several ARE-regulated genes, in addition to HO-1, through activation of Nrf2 (Iwasaki et al., 2006). However in our *in vitro* system, the effect of hemin in macrophages is similar to that seen with CoPP treatment, with preferential activation of HO-1 and no effect on HIV replication. And consistent with previous findings using CoPP (Figure 4.8), induction of HO-1 with hemin reduced neurotoxin production in HIV-infected macrophages, even in the absence of alterations to viral replication (Figure 4.11C). As previously demonstrated in Figure 4.8, treatment of HIV-infected macrophages with SnMP, which inhibits HO-1 enzymatic activity despite increasing HO-1 expression, had no effect on HIV replication but enhanced neurotoxin production. While robust induction of the antioxidant response attenuates HIV replication and NF- $\kappa$ B driven inflammation, these effects are mediated by induction of ARE-regulated genes other than HO-1. However, the enzymatic activity of HO-1 is sufficient to alter neurotoxin production pathways, possibly by increasing iron, biliverdin or carbon monoxide (CO) levels, the catalytic byproducts following heme catabolism by HO-1. Future experiments will explore the role of HO-1 enzymatic activity in modulating known neurotoxin production pathways in macrophages.






**Figure 4.11 Hemin, an inducer of HO-1 and the antioxidant response, attenuates macrophage-mediated neurotoxicity without altering HIV replication.** **A)** Human MDM were infected with HIV-Jago and treated with 20µM hemin, 10µM SnMP or the non-nucleoside reverse transcriptase inhibitor, efavirenz (EFZ), over the course of infection. Culture supernatants were collected every 3 days and assessed for RT activity. Values indicate the mean  $\pm$  SEM. **B)** 50µM hemin increases total levels of HO-1 and modestly increases NQO1 levels, as assessed by Western blotting in HIV-infected human macrophages. **C)** Hemin reduces macrophage-mediated neurotoxicity following HIV infection. Neuronal survival was assessed by MAP2 ELISA following exposure of rat cerebrocortical neuronal cultures to supernatant of HIV-infected macrophages treated with hemin over the course of infection. MAP2 data is expressed as a percentage of untreated (UT) cultures (n = 6). All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing, \*\*\*p<0.001.

*DMF treatment reduces levels of the HIV co-receptors, CCR5 and CXCR4, on human monocytes*

We have demonstrated that treatment of macrophages with DMF or MMF over the course of infection inhibits HIV replication (Figure 4.1). We hypothesized that DMF and MMF could inhibit HIV infection by decreasing levels of the HIV co-receptors, CCR5 and CXCR4, via alterations of the cellular redox state (Cross et al., 2011). As shown in Figure 4.12, CD11b<sup>+</sup>CD14<sup>+</sup> PBMCs had significantly less cell surface expression of CXCR4 after 6 hours and CCR5 after 36 hours of DMF treatment. DMF treatment of CD11b<sup>+</sup>CD14<sup>+</sup> PBMCs transiently increased cell surface levels of CD4 after 6 hours of treatment. However, CD4 cell surface expression was equal to or less than the vehicle control after 36 hours of DMF treatment. DMF's effect on receptor expression was not global, as levels of CD14 did not significantly change with treatment. These results suggest that DMF and MMF treatment may attenuate HIV replication in macrophages by inhibiting receptor-mediated viral entry. Furthermore, they suggest that DMF treatment can inhibit infection of human monocytes by both CCR5- and CXCR4-tropic viruses

We are currently investigating the effects of DMF treatment on early stages of viral replication through the use of virus pseudotyping. A pseudotyped virus is an enveloped virus particle, in this case HIV, which has been assembled with a foreign viral glycoprotein, such as vesicular stomatitis virus glycoprotein (VSV-G). VSV-G pseudotyped viruses can infect macrophages independently of the traditional HIV co-receptors. Luciferase assays, detecting transcription from the *nef* gene of the HIV genome, will be performed on infected macrophages, in the presence and absence of DMF, to assess if DMF treatment alters early events in viral infection (reverse transcription, nuclear translocation, incorporation into the genome and transcription/translation) in addition to viral entry.



**Figure 4.12 DMF decreases the cell surface expression of CXCR4 and CCR5 on human monocytes<sup>3</sup>.** **A)** DMF significantly decreases CXCR4 cell surface expression on CD11b<sup>+</sup>CD14<sup>+</sup> human PBMCs following 6 hours of treatment, **B)** but has little effect following 36 hours. **C)** DMF has no significant effect on CCR5 cell surface expression on CD11b<sup>+</sup>CD14<sup>+</sup> human PBMCs after 6 hours of treatment, but **D)** significantly decreases CCR5 expression following 36 hours. **E)** DMF significantly increases CD4 cell surface expression on CD11b<sup>+</sup>CD14<sup>+</sup> human PBMCs following 6 hours of treatment, **F)** but decreases CD4 expression following 36 hours of treatment. DMF has no significant effect on CD14 cell surface expression on CD11b<sup>+</sup>CD14<sup>+</sup> human PBMCs after **G)** 6 hours or **H)** 36 hours of treatment. For all experiments, values represent data averaged from 6-8 different donors. All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing, \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001 vs. Vehicle.

<sup>3</sup> All of the data contained in this figure was collected and analyzed by Anthony W.S. Chi, PhD

*DMF treatment of HIV-infected macrophages has no effect on viral replication after infection is established; although DMF, but not efavirenz, decreases macrophage-mediated neurotoxicity*


If DMF treatment decreases HIV infection by reducing the cell surface expression of the HIV co-receptors, then addition of DMF to an established infection would have little effect on HIV replication. Figure 4.13 demonstrates that as expected, the NNRTI, efavirenz, does not alter HIV replication when added to macrophages at day 9 post infection. Since efavirenz attenuates HIV replication by inhibiting the reverse transcription of the HIV RNA genome immediately following HIV entry, efavirenz application to an established infection would have no antiviral activity.

Efavirenz treatment of HIV-infected macrophages supporting high levels of viral replication does not attenuate neurotoxin production (Figure 4.13B). When added to an established infection, DMF treatment has no effect on HIV replication (Figure 4.13C). This supports the hypothesis that DMF attenuates viral replication by inhibiting cell surface CCR5 and CXCR4 levels and thereby viral entry. However, when DMF was added to macrophages supporting high levels of HIV replication, there was significantly less neurotoxin production, as compared to vehicle control (Figure 4.13D). This finding is in agreement with previous studies demonstrating that induction of HO-1 reduces neurotoxin production in HIV-infected macrophages, even without attenuation of HIV replication.

These experimental results also suggest that DMF treatment has little effect on HIV replication kinetics after HIV has entered the cell and incorporated into the host genome. We had proposed that DMF's antiviral effects were a consequence of decreased transcription from the HIV LTR due to attenuation of NF- $\kappa$ B and TNF $\alpha$  signaling (Cross et al., 2011). However, if this hypothesis were true, addition of DMF to HIV-infected macrophages should have decreased viral transcription.

DMF-mediated inhibition of NF- $\kappa$ B and TNF $\alpha$  signaling has important implications for inflammation mediated pathology of HAND, but our data suggests that this is not a major mechanism underlying DMF's antiviral effects. Rather, DMF treatment prevents the infection of macrophages and would limit viral spread of both CXCR4 and CCR5 tropic viruses. Furthermore,

DMF is an attractive adjunctive therapy for HAND, since treatment of infected macrophages attenuates neurotoxin production, whereas traditional antiretrovirals would have little effect.



**Figure 4.13 DMF, but not efavirenz, added to an established HIV infection decreases macrophage-mediated neurotoxicity without altering HIV replication.** **A)** MDM infected with HIV-Jago were treated with 20nM of the non-nucleoside reverse transcriptase inhibitor, efavirenz (EFZ), over the course of infection or were left untreated until day 6 post infection, and then treated with EFZ for the remainder of the time course (+EFZ). **B)** HIV/MDM treated with EFZ over the course of infection do not produce neurotoxins, whereas HIV/MDM treated with EFZ from day 6 through day 12 (+EFZ) cause as much neurotoxicity as vehicle controls. **C)** MDM infected with HIV-Jago were left untreated until day 6 post infection, and then treated with 15uM or 30uM DMF until day 9 post infection. **D)** HIV/MDM treated with 15uM and 30uM DMF from day 6 through day 9 post infection cause significantly less neurotoxicity than vehicle control, despite having equivalent levels of RT activity. For panels A and C, culture supernatants were collected every 3 days and assessed for RT activity. Values indicate the mean  $\pm$  SEM. RT curves are representative of 2 independent experiments, with each replicate performed on cell preparations from a different donor. For panels B and D, neuronal survival was assessed by MAP2 ELISA following exposure of rat cerebrocortical neuronal cultures to supernatant of HIV-infected macrophages from the end of the time course (day 12 in panel B and day 9 in panel D). MAP2 data is expressed as a percentage of untreated (UT) cultures ( $n = 6$ ). All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing, \*\*\* $p < 0.001$  vs. Vehicle.

## **CHAPTER 5**

### **THE UNFOLDED PROTEIN RESPONSE ATTENUATES HIV REPLICATION AND MACROPHAGE-MEDIATED NEUROTOXICITY**

## Abstract

Viral infection can result in the activation of host cell adaptive stress responses, including the antioxidant response and the unfolded protein response (UPR). Virus replication and assembly can strain the capacity of the endoplasmic reticulum (ER) and result in the accumulation of misfolded proteins and activation of the UPR. The UPR is designed to eliminate misfolded proteins and promote cellular recovery by attenuating translation and upregulating the expression of chaperones, degradation factors, and regulators of metabolic and redox states. Several viruses, including hepatitis C virus and herpes simplex virus, regulate antiviral components of the UPR in order to promote efficient viral replication. However, the effect of HIV infection on the UPR has not yet been investigated. We hypothesized that HIV infection would activate and modulate the UPR in infected macrophages. Using our *in vitro* model system, we found that HIV infection increases levels of phosphorylated eIF2 $\alpha$ , which inhibits protein translational during states of cellular stress. We also found that the chaperone BiP, considered the master regulator of the UPR, was increased in the macrophages of HAND frontal cortex. While HIV infection activates components of the UPR in macrophages, pharmacological induction of the UPR attenuates HIV replication. This suggests that induction of phosphorylated eIF2 $\alpha$  during HIV replication has some antiviral effect, although it is not sufficient to considerably attenuate HIV replication. Interestingly, pharmacological induction of the UPR, which attenuates viral replication, is associated with increased macrophage-mediated neurotoxicity. This finding has important implications for the development of adjunctive therapies for HAND. Therapeutics or processes that induce the UPR in macrophages, regardless of the effect on HIV replication, could enhance neurotoxin production and contribute to the pathological processes underlying HAND.

As we have previously described the effect of HIV replication on the antioxidant response, we assessed how restoration of the antioxidant response altered the UPR in HIV-infected macrophages. Activation of the antioxidant response with DMF and MMF attenuated HIV replication and decreased levels of phosphorylated eIF2 $\alpha$  and BiP in HIV/MDM. Reductions in

phosphorylated eIF2 $\alpha$  trend with increased levels of HO-1, but do not depend upon HO-1 enzymatic activity. In contrast, BiP levels decrease with induction of the antioxidant response, but these alterations are independent of HO-1. These findings suggest that while dysregulation of the antioxidant response reflects levels of HIV replication and HO-1 levels modulate neurotoxin production, levels of phosphorylated eIF2 $\alpha$  and BiP do not predict HIV replication or extent of macrophage-mediated neurotoxicity. Ongoing studies are assessing the link between UPR activation, inflammation pathways and the antioxidant response as they relate to HIV infection and macrophage-mediated neurotoxicity.

HIV infection alters several adaptive stress pathways in the macrophage. The suppression of the antioxidant response and induction of the UPR both associate with high levels of viral replication and neurotoxin production. Pharmacological induction of either stress pathway attenuates HIV replication, however, induction of the antioxidant response, but not the UPR, decreases neurotoxin production. Therefore, activation of the UPR by inflammation, oxidative stress or ART is predicted to enhance neurotoxin production in macrophages, regardless of the level of HIV replication. Understanding how HIV infection affects adaptive stress responses and neurotoxin production pathways in the macrophage will continue to improve our ability to develop effective adjunctive therapies for HAND.



## Introduction


Accumulation of misfolded proteins in the endoplasmic reticulum (ER) lumen results in the activation of an adaptive stress response known as the unfolded protein response (UPR). The UPR is a quality control mechanism that can be activated during physiological stress (e.g., glucose deprivation, oxidative stress) and viral infection. The UPR reduces the load of misfolded proteins in the ER by attenuating new protein translation via phosphorylated-eIF2 $\alpha$  and upregulating chaperones that mediate protein folding, such as BiP. During viral infection, the cell can experience an excess flux of viral proteins passing through the ER. Several viruses, including human cytomegalovirus (CMV) and herpes simplex virus (HSV), are capable of regulating cellular responses to infection, including the UPR, in order to promote host cell survival while promoting efficient viral replication. The effect of HIV infection on the UPR has not yet been investigated.

### *The Unfolded Protein Response*

The endoplasmic reticulum facilitates protein folding, is part of the protein-sorting pathway and is the site of post-translational modifications such as N-linked glycosylation and disulfide bond arrangement. These ER processes are tightly monitored in order to prevent improperly folded or modified proteins from being released into the cell. Under normal conditions, approximately 30% of newly synthesized proteins fail to achieve native structure due to errors in transcription, translation, post-translational modifications or protein folding (Schubert et al., 2000). If not managed, misfolded proteins accumulate in the ER and can result in toxic aggregates that interfere with the function of normal proteins (Romisch, 2004). In response to increased levels of misfolded proteins in the ER lumen, a signaling cascade known as the Unfolded Protein Response (UPR) or Integrated Stress Response (ISR) is activated. This signaling cascade limits the translation of new proteins, which decreases the incoming burden on the ER machinery, enhances the degradation of improperly folded proteins and upregulates levels of chaperones, which assist in protein folding/refolding. ER stress and induction of the UPR can occur under physiological conditions known to increase levels of misfolded protein, such as glucose

deprivation (Kaufman et al., 2002), heat shock (Kasuya et al., 1999; Matsumoto et al., 2005), oxidative stress (Holtz et al., 2006; Yan et al., 2008), hypoxia (Feldman et al., 2005; Koumenis and Wouters, 2006), increase demand of secretory function (B-cell antibody production) (Gass et al., 2002), and viral infection (Ambrose and Mackenzie, 2011; He et al., 1997; Isler et al., 2005; Yoshida et al., 2001).

Activation of the UPR attempts to eliminate misfolded proteins in the ER lumen through two mechanisms; 1) through attenuation of protein translation, in order to reduce the flux of proteins entering into the ER and 2) through the induction of chaperone proteins and degradation factors to refold and/or eliminate misfolded proteins. Three proteins have been identified as sensors of ER stress: protein kinase RNA-like ER kinase (PERK), activating transcription factor 6 (ATF6), and inositol-requiring enzyme 1 (IRE1). These proteins are transmembrane proteins with a sensor domain in the ER lumen and a kinase effector domain in the cytosolic compartment. Under normal conditions, the ER chaperone immunoglobulin heavy-chain-binding protein (BiP), also known as glucose regulated protein-78 (GRP78), is bound to the sensor domain of PERK, ATF6 and IRE1 in the ER lumen. When misfolded proteins accumulate in the ER, BiP is sequestered away from these sensors to bind to misfolded proteins (Bertolotti et al., 2000). BiP release from PERK and IRE1 results in the homodimerization of their luminal domains, causing autophosphorylation and activation of downstream signaling cascades (Figure 5.1). In contrast, BiP release from ATF6 unmasks a Golgi localization signal, resulting in the relocation of ATF6 to the Golgi, where it is cleaved into an activated form (Shen et al., 2002).



**Figure 5.1 The signaling pathways comprising the Unfolded Protein Response (UPR).**

Accumulation of misfolded protein in the ER causes ER stress and results in the activation of the UPR, an adaptive stress pathway. PERK, IRE1 and ATF6 have sensor domains that monitor levels of misfolded protein in the ER lumen. Following activation, each of these pathways initiates a signaling cascade that results in the transcriptional upregulation of genes that will help the cell reduce, manage or recover from ER stress. Phosphorylation of eIF2 $\alpha$  by activated PERK results in the attenuation of all cap-dependent protein translation, in order to reduce the incoming burden of new proteins into the ER. Increased levels of phosphorylated eIF2 $\alpha$  also result in the translational upregulation of ATF4, a transcription factor that increases amino acid response element (AARE)-regulated genes in order to modulate cellular metabolism and redox state during cellular recovery from ER stress. ATF4 also increases levels of GADD34, which restores protein translation by mediating eIF2 $\alpha$  dephosphorylation as part of a negative feedback loop. In addition to PERK, three other kinases (PKR, GCN2 and HRI) can phosphorylate eIF2 $\alpha$  and initiate the UPR, although instead of misfolded protein levels, they respond to double stranded RNA, amino acid limitation and heme levels, respectively. ATF6, upon activation, translocates to the Golgi where it is cleaved into an active form, which also functions as a transcriptional regulator of ER stress-element (ERSE)-regulated genes. ERSEs regulate genes encoding for a variety of cellular chaperones, which promote the proper folding of misfolded proteins in the ER, including BiP and GRP94. Activation of IRE1 results in its auto-phosphorylation and the subsequent activation, by a splicing event, of XBP1 mRNA. Spliced XBP1 (sXBP1) encodes a transcription factor capable of upregulating genes with ERSEs or UPR elements (UPREs). Transcriptional upregulation of UPRE-regulated genes results in increased levels of EDEM, which enhances the degradation of misfolded proteins in the ER by ER-associated degradation (ERAD). If the UPR is unable to manage the stress to the ER, the cell will undergo CHOP-mediated apoptosis.

After activation, PERK initiates a signaling cascade that results in the inhibition of new protein translation and the induction of genes important for cellular recovery from ER stress. PERK directly phosphorylates the eukaryotic initiation factor eIF2 $\alpha$  at serine 51 (Harding et al., 2000b; Harding et al., 1999). When phosphorylated, eIF2 $\alpha$  inhibits cap-dependent protein translation by tightly binding eIF2B. This prevents eIF2B from catalyzing the GDP-GTP exchange and thereby stops the binding of the initiator Met-tRNA to the ribosome and initiation of translation (Krishnamoorthy et al., 2001). While inhibiting the majority of cellular protein translation, phosphorylated-eIF2 $\alpha$  specifically increases the translation of select stress-responsive proteins, including activating transcription factor 4 (ATF4) (Harding et al., 2000b). ATF4 mRNA is efficiently translated due to small upstream open-reading frames (uORF) within the 5' untranslated region, resulting in a "uORF bypass scanning system" (Lu et al., 2004). ATF4 binds to the promoter regions of an array of different genes, including metabolism and redox regulatory factors involved in mediating recovery from ER stress. ATF4 deficiency results in increased sensitivity to cell death with a variety of stresses, including oxidative stress and amino acid deprivation (Harding et al., 2003). ATF4 also activates an inhibitory feedback loop to restore cellular protein translation by inducing levels of growth arrest- and DNA damage-inducible gene 34 (GADD34), which interacts with protein phosphatase 1 (PP1) and increases the dephosphorylation of eIF2 $\alpha$  (Brush et al., 2003; Ma and Hendershot, 2003).

When BiP is sequestered away from ATF6 during ER stress, the inactive 90-kDa ATF6 precursor translocates to the Golgi where it is cleaved by site-1 and site-2 proteases into the transcriptionally active 50-kDa protein (Ye et al., 2000). ATF6, along with the constitutively expressed transcriptional regulator, nuclear factor Y (NF-Y), activate genes carrying ER stress response elements (ERSEs) in their transcriptional promoters [Mori, Cell, 2000]. ERSEs are found in genes encoding for cellular chaperones including BiP, GRP94, protein-disulfide isomerase (PDI) and calreticulin (Harding et al., 2003). Consequently, ATF6 activation results in the upregulation of ER chaperones and other proteins that are needed to promote proper protein

folding in an effort to relieve ER stress. Increased levels of BiP dampen PERK, IRE1 and ATF6 activation when the misfolded protein burden in the ER is managed. Furthermore, ATF6 can increase the amount of X box binding protein 1 (XBP-1) mRNA, providing a link between the ATF6 and IRE1 pathways (Lee et al., 2002; Yoshida et al., 2001).

Phosphorylation of IRE1, which has intrinsic endonuclease activity, results in the removal of a 26-nucleotide intron from the XBP-1 transcript (Lee et al., 2002). The spliced form of XBP-1 (sXBP-1) mRNA encodes an active transcription factor, capable of inducing genes regulated by the ERSE (Lee et al., 2003). In addition, sXBP-1 can also activate the transcription of genes containing UPR elements (UPRE), which are responsive to ER stress and are distinct from the ERSE (Lee et al., 2002; Yamamoto et al., 2004; Yoshida et al., 2001). One consequence of sXBP1 activation of the UPRE is the enhanced transcription of ER degradation enhancing  $\alpha$ -mannosidase-like protein (EDEM). EDEM increases the capacity of ER-stressed cells to degrade irrevocably misfolded proteins via ER-associated protein degradation (ERAD) (Hosokawa et al., 2001; Yoshida et al., 2003). EDEM promotes the release of terminally misfolded proteins from the ER-resident chaperone calnexin, thereby making them available for transport to the cytoplasm for degradation by the ubiquitin proteasome system (UPS) (Kopito, 1997; Oda et al., 2003). Therefore, while the ATF6 pathway is important for protein refolding, the IRE1-XBP1 pathway mediates protein degradation via ERAD (Yoshida et al., 2003).

Chronic activation of the UPR, which occurs when the cell cannot manage or recover from ER stress, results in the activation of signaling pathways that will commit the cell to apoptosis (Ferri and Kroemer, 2001; Oyadomari et al., 2002a). At least three apoptosis pathways have been implicated in this apoptotic event. The first is transcriptional activation of C/EBP homologous protein (CHOP, also known as GADD153), primarily by ATF4 (Fawcett et al., 1999; Harding et al., 2000a; Ma et al., 2002), although the *CHOP* promoter can also be regulated by IRE1 and ATF6 (Wang et al., 1998; Yoshida et al., 2000). Induction of CHOP compromises cell viability

(Friedman, 1996; McCullough et al., 2001; Zhan et al., 1994) and targeted disruption of *CHOP* significantly protects cells from apoptosis following sustained ER stress (Oyadomari et al., 2002b; Zinszner et al., 1998). The second is activation of the cJUN NH<sub>2</sub>-terminal kinase (JNK) pathway, mediated by a complex between IRE1, TNF receptor-associated factor 2 (TRAF2) and apoptosis signal-regulating kinase1 (ASK1) (Nishitoh et al., 2002; Urano et al., 2000). The third pathway involved in initiating apoptosis in ER-stressed cells is mediated by caspase-12. Caspase-12, which is localized to the ER, is activated by specifically ER stress and not by apoptotic signals originating from the membrane or mitochondria (Nakagawa et al., 2000). All three of these apoptosis pathways ultimately lead to the activation of caspase-3 with BAX (Bcl-2-associated X protein) and BAK (Bcl-2 homologous antagonist/killer) mediating the final steps of ER stress-mediated apoptosis (Wei et al., 2001).

#### *Viral infection activates the UPR*

In some cases, viral infection induces ER stress and activates components of the three branches of the UPR. BiP, considered the master regulator of the UPR, is induced in cells infected with respiratory syncytial virus (RSV) (Bitko and Barik, 2001), hanta viruses (Li et al., 2005), hepatitis C virus (HCV) (Liberman et al., 1999) and several members of the flavivirus family (Jordan et al., 2002; Su et al., 2002; Tardif et al., 2002). Activation of PERK has been reported in cells infected with herpes simplex virus (HSV) (Cheng et al., 2005; Mulvey et al., 2007) and cytomegalovirus (CMV) (Isler et al., 2005). Furthermore, the IRE1 and XBP-1 signaling pathway is activated with dengue infection (Yu et al., 2006) and the ATF6 pathway is activated in HCV infected cells (Tardif et al., 2004). Enveloped viruses utilize the ER as the primary site of envelope glycoprotein biogenesis and several viruses undergo genomic replication and particle assembly in the ER compartment. Increasing the burden on the ER, especially at times of high viral production, would easily increase the accumulation of misfolded proteins in the ER lumen and activate the UPR. However, while activation of the UPR is essential for host cell survival during viral infection, some of the consequences of UPR activation would be detrimental to viral replication. Consequently,

many viruses have evolved mechanisms to regulate the UPR and promote efficient viral replication.

Some of the consequences of UPR activation would be detrimental to efficient viral replication, while others would promote viral protein folding and assembly. Activation of the PERK pathway and phosphorylation of eIF2 $\alpha$  results in the attenuation of protein translation for most transcripts in the host cell and viral genome dependent upon cap-dependent protein translation. Thus, a virus that causes phosphorylation of eIF2 $\alpha$  must escape translational inhibition in order to express viral and cellular gene products necessary for viral replication. However, activation of ATF4 results in the transcriptional upregulation of genes involved in reestablishing cellular metabolism (as part of the UPR recovery effort), translation and buffering oxidative stress (Harding et al., 2003), and may therefore be beneficial to viral replication. ATF6 activation typically results in the activation of molecular chaperones, whose expression would benefit virus replication by promoting the proper folding of viral proteins. Activation of XBP-1, however, increases the transcription of EDEM and thereby enhances the degradation of misfolded protein by ERAD. Activation of IRE1 and XBP-1 could hamper viral replication by degrading viral proteins that are not efficiently folded within the ER lumen.

#### *Viruses can modulate components of the UPR*

Several viruses that are known to induce ER stress and activate components of the UPR have adapted strategies to avoid or shutdown signaling pathways that would attenuate viral replication. Infection by herpes simplex virus (HSV) results in the activation of PERK and phosphorylation of eIF2 $\alpha$  (Wylie et al., 2009), however, the virus is able to escape from the resultant translational inhibition (He et al., 1997). The  $\gamma_134.5$  gene of HSV has homology to GADD34, whose expression level modulates the rate of eIF2 $\alpha$  dephosphorylation by protein phosphatase 1 (PP1). HSV infection results in the transcription and translation of  $\gamma_134.5$ . As HSV induces ER stress and phosphorylation of eIF2 $\alpha$ ,  $\gamma_134.5$  associates with PP1 and mediates the dephosphorylation of

eIF2 $\alpha$ . HSV, which has a large genome, encodes a viral protein to modulate the UPR. For many other viruses, the mechanisms by which they modulate and avoid the consequences of the UPR are not as well understood.

Cytomegalovirus (CMV) infection results in the activation of PERK, phosphorylation of eIF2 $\alpha$  and increased translation of ATF4. Despite this strong activation of the PERK-eIF2 $\alpha$  pathway, CMV infection does not attenuate protein translation. The mechanism by which CMV overcomes the translational inhibition normally mediated by phosphorylated eIF2 $\alpha$  is unknown (Isler et al., 2005).

Infection with dengue virus also results in increased levels of phosphorylated eIF2 $\alpha$ , although this induction does not attenuate protein translation of either host or viral transcripts (Edgil et al., 2006). During infection, dengue virus may sustain host and viral cell protein translation by simultaneously inducing the phosphorylation of eIF2 $\alpha$  and increasing total levels of eIF2 $\alpha$  (Umareddy et al., 2007), thereby reducing the activated fraction of total eIF2. Interestingly, pharmacological induction of ER stress in dengue-infected cells inhibits approximately 50% of cellular protein synthesis, while translation of dengue viral proteins is only inhibited 0-20% (Edgil et al., 2006). This finding suggests that translation of dengue viral proteins, as compared to most host cell proteins, can occur efficiently under conditions of impaired cap-dependent protein translation. While the exact mechanism is still unknown, dengue virus can replicate by a novel noncanonical, cap-independent mechanism that is independent of eIF4E (Edgil et al., 2006). eIF4E, the rate limiting component of cap-dependent translation, directs ribosomes to the cap structure of mRNA transcripts.

Many viruses actively antagonize the host translational machinery in order to shut off host cell protein translation, thereby preventing the activation of host-mediated antiviral responses.

Picornavirus and poliovirus infection results in the cleavage of eIF4G isoforms by viral proteases and results in host translational shutdown (Gradi et al., 1998). In order for viral protein translation



to continue unimpaired, viruses such as poliovirus and hepatitis C virus (HCV) utilize internal ribosome entry sites (IRES) (Pelletier and Sonenberg, 1988). These are RNA elements that directly recruit ribosomes in a cap-independent manner, thereby requiring only a subset of the canonical eIFs. While the cellular mechanisms underlying IRES-dependent translation are still under investigation, the use of IRES by viruses has been more intensively studied. The HCV IRES directly bind the 40S ribosomal subunit in such a way that the HCV initiator codons enter the ribosomal P-site without mRNA scanning. Consequently, HCV IRES-dependent translation requires eIF2 and eIF3, but not eIF1 or eIF4 isoforms (Fraser and Doudna, 2007). In contrast, the IRES of picornaviruses does not attract the 40S ribosomal subunit directly, rather, a high-affinity binding site for eIF4G brings viral mRNA into contact with the ribosome. Translation from the IRES is dependent upon all of the standard eukaryotic translation initiation factors, except for the cap-binding protein eIF4E. However, picornavirus transcripts recruit several RNA-binding proteins that are not usually involved in translation, such as La protein and the polypyrimidine tract-binding protein (PTB) (Hunt and Jackson, 1999), to stimulate their IRES-mediated translation. How these recruited proteins and other IRES *trans*-acting factors (ITAFs) aid in IRES-dependent protein translation is still under active investigation.

In addition to modulation of the PERK-eIF2 $\alpha$  pathway and translation initiation, viruses can also modulate the rate of ERAD in order to prevent viral protein degradation. HCV infection results in increased splicing of the uXBP1 transcript and elevation of sXBP1 protein levels (Tardif et al., 2004). However, the transcriptional upregulation of UPR genes and ERAD activity is repressed in HCV-infected cells, suggesting that HCV dampens the transcriptional activity of sXBP1 (Tardif et al., 2004). This defect in IRE1-XBP1 pathway signaling may ultimately result in the translation of more viral protein, since functional knockdown of IRE1 results in increased IRES-dependent translation (Tardif et al., 2004). Based upon these studies, it is predicted that a host cell experiencing ER stress following HCV infection may promote viral translation over host cell protein translation due to HCV-mediated modulation of the UPR.

While activation of the UPR is primarily an adaptive stress response, it can also be responsible for virus-induced cellular pathology. For example, respiratory syncytial virus (RSV), a lytic virus, primarily spreads by inducing cellular apoptosis via caspase-12, which is specifically activated with sustained ER-stress. Functional knockdown of caspase-12 protects RSV-infected cells from apoptosis, demonstrating that activation of the UPR is part of the viral life cycle (Bitko and Barik, 2001). Moloney murine leukemia virus (MoMLV), a retrovirus that causes a non-inflammatory spongiform neurodegenerative disease, induces the UPR as a consequence of viral envelope misfolding in the ER (Dimcheff et al., 2003). Interestingly, the severity of neurological disease correlates to the folding instability of the viral envelope, suggesting that the observed spongiform neurodegeneration is the consequence of a virus-induced protein folding disorder (Portis et al., 2009). Viruses that induce ER-stress and the UPR have evolved many mechanisms to promote efficient replication and avoid the antiviral consequences of host-cell adaptive stress responses. The effect of HIV on the UPR and the role of the UPR in HIV replication and neurotoxicity have not yet been explored.

#### *HIV proteins are processed in the ER and Golgi*

The genome of HIV consists of 9.8 kilobases of single-stranded RNA that encode nine viral genes. The HIV genome encodes for three structural genes (*env*, *gag* and *pol*) and six regulatory genes, which control viral replication (*vif*, *rev*, *nef*, *tat*, *vpu* and *vpr*). The extensive glycosylation of the HIV-1 envelope glycoproteins (gp), gp120 and gp41, plays an important role in evasion of the host immune system and is required for infectivity (Wei et al., 2003). First synthesized in the endoplasmic reticulum, the 160kD envelope precursor protein (cleaved into gp120 and gp41 in later processing stages) undergoes extensive processing with the addition of 25-30 N-linked glycosylations (Leonard et al., 1990). In fact, approximately 50% of the molecular weight of envelope glycoprotein is due to oligosaccharide modifications. In addition to requiring extensive folding and glycosylation in the ER, HIV proteins also require the host cell's Golgi and endosome

network in order to complete the viral lifecycle. Newly synthesized envelope proteins are sent to the plasma membrane, where they must be endocytosed and routed back to the plasma membrane via the *trans*-Golgi-network (TGN) (Rowell et al., 1995). The TGN is a critical site for HIV biogenesis, as it where envelope glycoproteins incorporate into infectious viral particles. In HIV-infected macrophages, virus release then proceeds through the subsequent release of mature viruses by exocytosis (Nguyen et al., 2003; Pelchen-Matthews et al., 2003). Given the extent of the involvement of the ER and Golgi in HIV replication, HIV infection and may induce the accumulation of misfolded proteins in the ER and activate the UPR, especially during times of high viral replication. Interestingly, it has been reported that HIV's envelope protein directly interacts with BiP (Earl et al., 1991).

#### *HIV infection may induce and modulate the UPR*

Given the extent of involvement of the ER and Golgi in HIV biogenesis, we hypothesize that HIV infection induces the UPR in macrophages, especially at times of high viral replication. However, since HIV replicates efficiently without causing cellular apoptosis in macrophages, HIV may also modulate the UPR in order to prevent the inhibition of protein translation and upregulation of CHOP. Furthermore, macrophages experiencing ER stress may be more prone to producing neurotoxins and/or inflammatory mediators that drive HAND pathology. In order to address these questions, we will use our *in vitro* model system and pharmacologic manipulations of the UPR in infected and non-infected human macrophages. We will assess the impact of HIV infection on the UPR, the role of the UPR in modulating HIV infection and the consequences of UPR induction on neurotoxin production.

## Results

### *HIV infection induces phosphorylation of eIF2 $\alpha$ in human MDM*


It has become increasingly clear that viruses that activate adaptive stress responses in the host cell have adapted to modulate or take advantage of these responses in order to promote efficient viral replication. We hypothesized that HIV infection of human macrophages would result in the activation of the UPR, especially during periods of high viral replication. As shown in Figure 5.2B, HIV-infected monocyte derived macrophages (HIV/MDM) demonstrate a biphasic induction of phosphorylated eIF2 $\alpha$ , while total eIF2, BiP and GRP94 (probe not shown) levels remain steady. Phosphorylated eIF2 $\alpha$  is highest following the 24hours of infection and at times of high viral load. This increase in phosphorylated eIF2 $\alpha$  is a consequence of active HIV replication, since HIV-infected macrophages treated with the non-nucleoside reverse transcriptase inhibitor, efavirenz (EFZ), do not show elevations in phosphorylated eIF2 $\alpha$  at either 24hours post infection (HPI) or at times of high viral reverse transcriptase (RT) activity (Figure 5.2D). EFZ treated HIV/MDM are exposed to viral inoculum and have normal levels of viral entry but the HIV-genome is not reverse transcribed, incorporated into the host genome or actively replicated. This suggests that HIV infection of the host cell increases levels of phosphorylated eIF2 $\alpha$  via viral post-entry and post reverse transcription mechanisms. Furthermore, the biphasic induction of phosphorylated eIF2 $\alpha$  may represent two separate signaling processes with two different time courses, both of which result in enhanced phosphorylation of eIF2 $\alpha$ . Currently, we do not know the kinase responsible for phosphorylating eIF2 $\alpha$  in HIV-infected macrophages.

In addition to PERK, the classic kinase mediating eIF2 $\alpha$  phosphorylation in the UPR, there are three other kinases capable of phosphorylating eIF2 $\alpha$ . These are protein kinase RNA regulated kinase (PKR), heme-regulated inhibitor kinase (HRI) and general control nonderepressible-2 kinase (GCN2). All of these kinases inhibit protein synthesis by phosphorylation of eIF2 $\alpha$ , although their tissue distributions and activator domains make each kinase uniquely suited to detect different stress conditions. PKR is induced by interferon and double-stranded RNA

(dsRNA) and is a principal mediator of host-cell antiviral responses since dsRNA rarely occurs under normal conditions within the cell. PKR activation has been reported during human T-lymphotropic virus type I (HTLV-1) (Mordechai et al., 1995), Epstein-Barr virus (EBV) (Elia et al., 1996), Hepatitis D (Circle et al., 1997; Robertson et al., 1996) and HIV infection (Maitra et al., 1994; Roy et al., 1991), among others. GCN2, which is highly expressed in the liver and brain, is activated under conditions of nutritional deprivation and inhibits protein translation via phosphorylated eIF2 $\alpha$  in order to conserve amino acids for essential metabolic processes (Sood et al., 2000). However, stressors unrelated to nutritional deprivation, such as UV irradiation, proteasomal inhibition and viral infection can also activate GCN2 (Jiang and Wek, 2005). Sindbis virus (SV) genomic RNA directly binds to GCN2 and functional GCN2 decreases the permissiveness of cells to both SV and vesicular stomatitis virus (VSV) infection. These findings suggest that GCN2 contributes to the cellular antiviral response to RNA viruses (Berlanga et al., 2006). HRI, which is primarily expressed in cells of the erythroid lineage, attenuates protein synthesis in heme-deficient states, coordinating hemoglobin synthesis with iron availability (Chen and London, 1995). HRI protein is also present in macrophages and HRI regulates iron homeostasis and macrophage maturation (Liu et al., 2007). Less is known about the role of HRI in viral infection, although HRI can be transcriptionally and translationally upregulated in response to heat shock, treatment with a dsRNA mimic (Polyinosinic:polycytidylic acid; poly I:C) and infection with the aquareovirus, grass carp hemorrhagic virus (GCHV) (Zhu et al., 2006). While each of the four eIF2 $\alpha$  kinases has a stimulus that they are particularly well suited to detect, it is clear that they can be broadly activated by several stresses and that any or all of these kinases could be activated during viral infection.

HIV infection in human macrophages induces the biphasic induction of phosphorylated eIF2 $\alpha$  without altering total eIF2 $\alpha$  or BiP levels (Figure 5.2). The increase in phosphorylated eIF2 $\alpha$  observed at either 24HPI or at times of high viral replication may be mediated by any of the four known eIF2 $\alpha$  kinases. At periods of high viral replication, the enhanced burden on the processing

capacity of the ER and Golgi by HIV glycoproteins could result in the accumulation of misfolded proteins and activation of PERK. The transactivation-responsive (TAR) element of HIV forms an elaborate RNA tertiary structure, including a 23-basepair hairpin with three bulges capable of dimerization, which is capable of activating PKR (Heinicke et al., 2009). GCN2 could be activated during periods of high viral replication if the rate of viral protein synthesis exceeded the availability of amino acids needed for basal host cell translation. Cellular iron homeostasis has been linked to rates of HIV transcription, suggesting that HRI activation and activity may impact HIV replication in macrophages (Debebe et al., 2007). In order to determine the kinase(s) responsible for eIF2 $\alpha$  phosphorylation in HIV-infected macrophages, genetic manipulation of the human macrophage is necessary. With typical transfection protocols, only 2-5% of primary human macrophages will express the inserted gene of interest. However, recent work with Minivector DNA (a non-viral, supercoiled vector incorporating short hairpin RNA) has been reported to work well in primary cells such as macrophages and dendritic cells (Zhao et al., 2011). This highly biostable vector produces high cell transfection efficiency and gene silencing capacity, which would be required in order to detect effects on HIV replication levels (Zhao et al., 2011). Using the Minivector DNA system, we could knockdown each of the eIF2 $\alpha$  kinases to determine the role of each in modulating phosphorylated eIF2 $\alpha$  levels in basal conditions and during HIV infection.



**Figure 5.2 HIV infection induces phosphorylation of eIF2α in human MDM.** **A)** MDM were infected with HIV-Jago (0-24 HPI) and culture supernatants were collected, as indicated, and assessed for reverse transcriptase (RT) activity. Values indicate the mean  $\pm$  SEM. **B)** HIV infection results in a biphasic induction of phosphorylated eIF2α without an increase of BiP, as assessed by Western blotting. Maximum levels of phosphorylated eIF2α occur at the end of the 24-hour infection and at times of high viral production/release. Blot is representative of 3 independent experiments, with each replicate performed on cell preparations from different donors. **C)** MDM were infected with HIV-Jago (day -1 to day 0) and culture supernatants were collected every 2-3 days, as indicated, and assessed for RT activity. Viral replication is effectively inhibited when MDM are treated with the non-nucleoside reverse transcriptase inhibitor, efavirenz (EFZ, 20nM), over the course of infection. Values indicate the mean  $\pm$  SEM. **D)** HIV infection results in increased levels of phosphorylated eIF2α at 24-hours post infection (24 HPI) and at times of high viral production/release (day 8 post infection), relative to EFZ-treated MDM, as assessed by Western blotting. Blot is representative of 3 independent experiments, with each replicate performed on cell preparations from different donors.


*HIV infection increases levels of macrophage specific BiP in the human frontal cortex of HAND*

Our *in vitro* model system of HIV-induced macrophage-mediated neurotoxicity utilizes primary human macrophages and rat cerebrocortical cultures. Following HIV-infection of human macrophages, cell culture supernatants are collected over the course of infection. This supernatant is applied to cerebrocortical cultures (in a 1:10 to 1:50 dilution) and neuronal survival is then assessed. While this model system utilizes primary cells and recapitulates the fundamental processes mediating HAND neuropathogenesis, many aspects of the human disease cannot be modeled. *In vitro*, the majority of our macrophages are infected with HIV over the course of the infection. In contrast, it is estimated that less than 0.1% of peripheral circulating monocytes are infected with HIV (Crowe et al., 2003). The proportion of HIV-infected CNS-resident macrophages is likely less than 5% during ART, although this is expected to be highly variable and dependent upon the CNS-penetrance of ART regimen, relative immune-activation state of the individual and extent of CNS neuroinvasion and inflammation, for which the susceptibility factors are not well understood. Since the overwhelming majority of cells in the CNS are astrocytes and neurons, techniques that selectively parse out effects in CNS-resident macrophages are required to assess the role of the macrophage in the neuropathology of HAND. We analyzed human frontal cortex for brain-resident macrophages by indirect immunofluorescence (IFA) in order to characterize the state of UPR activation in HIV-associated neurocognitive disorder (HAND).


IFA staining of BiP (Figure 5.3) and phosphorylated-eIF2 $\alpha$  (Figure 5.4) in human frontal cortex from HIV-, HIV+ and HAND cases was performed. IFA results demonstrate that HIV infection associated with HAND significantly increases total macrophage specific BiP, relative to uninfected controls (HIV-) (Figure 5.3D). Cognitively normal, HIV+ cases had macrophage specific BiP levels in between the uninfected controls and the HAND cases. This increase in macrophage specific BiP in HAND cases occurred in the absence of increased macrophage area (Figure 5.3A and 5.3B), suggesting that BiP expression is increasing on the individual cell level. Phosphorylated-



eIF2 $\alpha$  staining has proven more difficult. The signal for phosphorylated-eIF2 $\alpha$  in macrophages has thus far been weak and highly variable, resulting in quantifications with large error bars and no statistical significance between conditions (Figure 5.4). At this point, we have no evidence that levels of macrophage-specific phosphorylated eIF2 $\alpha$  are altered with either HIV infection or neurocognitive status.



**Figure 5.3 HIV infection increases levels of macrophage-specific BiP in the human frontal cortex of HAND cases.** **A)** Frontal cortex from cognitively normal HIV- and HIV+ cases and neurocognitively impaired HIV+ (HAND) cases were labeled with BiP, HLA-DR (macrophage marker, CCR3/43 antibody) and DAPI (DNA marker, blue). Quantification of Macrophage/HLA-DR pixel area (**B**) shows no increase of HLA-DR-positive cells in HIV+ or HAND cases. Quantification of integrated pixel intensity of BiP colocalizing with HLA-DR (**C**, **D**) shows increased cell type-specific BiP expression with HIV infection and neurocognitive impairment (HAND). Statistical comparisons were made by two-tailed *t*-test in C and one-way ANOVA plus Newman-Keuls post hoc testing in D, \**p*<0.05. Data is expressed as mean  $\pm$  SEM for *n* = 2 HIV-, *n* = 3 HIV+, cognitively normal and *n* = 9 HAND. For panels A and C, HIV+ represents all HIV+ cases (*n* = 12). Table 3.1, in the Methods section, provides further information on case history and tissue collection of individual samples.



**Figure 5.4 HIV infection does not alter levels of macrophage specific phosphorylated-eIF2α in the human frontal cortex of HAND cases.** A) Frontal cortex from cognitively normal HIV- and HIV+ cases and neurocognitively impaired HIV+ (HAND) cases was labeled for phosphorylated eIF2α and HAM56, a macrophage/microglia marker. Quantification of HAM56 pixel area (**A**, **B**) and integrated pixel intensity of HAM56 (**C**, **D**) shows no change in the staining of HAM56-positive cells in HIV+ or HAND cases, as compared to HIV- cases. Quantification of phosphorylated eIF2α pixel area (**E**, **F**) and integrated pixel intensity of phosphorylated eIF2α (**G**, **H**) shows no change in the staining of phosphorylated eIF2α-positive cells in HIV+ or HAND cases, as compared to HIV- cases. Quantification of integrated pixel intensity of phosphorylated eIF2α colocalizing with HAM56 (**I**, **J**) and integrated pixel intensity of phosphorylated eIF2α colocalizing with HAM56, normalized to GFAP area (**K**, **L**), shows no increased macrophage-specific phosphorylated eIF2α expression with HIV infection or neurocognitive impairment. Statistical comparisons were made by two-tailed *t*-test or one-way ANOVA plus Newman-Keuls post hoc testing and no significant differences were found (significance was defined as  $p < 0.05$ ). Data is expressed as mean  $\pm$  SEM for  $n = 4$  HIV-,  $n = 2$  HIV+, cognitively normal and  $n = 13$  HAND. For panels comparing only HIV- and HIV+, HIV+ represents all HIV+ cases ( $n = 15$ ). Table 3.2, in the Methods section, provides further information on case history and tissue collection of individual samples.

*Pharmacologic induction of the UPR by thapsigargin attenuates HIV replication and enhances neurotoxin production in human MDM*


Using our *in vitro* system, we found that HIV infection resulted in the activation of phosphorylated eIF2 $\alpha$  at times of high viral replication (Figure 5.2). We have also demonstrated that the UPR is activated, with increased BiP expression, in CNS-resident macrophages of the frontal cortex in HAND (Figure 5.3). While the extent of UPR activation following HIV infection has not yet been fully characterized, initial studies suggest that HIV infection activates components of the UPR in macrophages and that the extent of this activation may correlate with the clinical appearance of HAND. To investigate the role of UPR activation in modulating HIV replication and neurotoxin production in infected macrophages, we used pharmacological inducers of the UPR in HIV/MDM in our *in vitro* model system.

Pharmacological agents that disrupt calcium homeostasis in the ER, such as thapsigargin, induce the UPR. Thapsigargin is a non-competitive inhibitor of sarco/endoplasmic reticulum Ca<sup>2+</sup>-ATPase (SERCA), and is considered a classic activator of the UPR. Thapsigargin treatment decreases ER calcium levels, decreases the activity of calcium-dependent chaperones and induces the accumulation of misfolded proteins in the ER lumen. However, inhibition of SERCA can result in the secondary activation of plasma membrane calcium channels, resulting in increased cytosolic calcium levels and alterations in the regulatory functions of Ca<sup>2+</sup>-dependent kinases. While thapsigargin is considered a strong activator of the UPR, the signaling pathways induced by thapsigargin treatment vary according to cell type, concentration and duration of treatment and need to be empirically defined for each model system.

HIV-infected macrophages were pulsed with thapsigargin for 6 hours after infection was established and viral replication was robust. As shown in Figure 5.5, thapsigargin treatment drastically reduces HIV replication in the 48 hours following drug exposure. This reduction in viral replication is not a consequence of cell death, since HIV replication began to recover in the 2-4

days following thapsigargin exposure (Figure 5.5A). Thapsigargin treatment induced a robust increase in phosphorylated eIF2 $\alpha$  and a moderate increase in levels of BiP following the 6 hour treatment (Figure 5.5B). And, these markers of the UPR were not altered in vehicle treated HIV/MDM. This pharmacological induction of the UPR was transient and could no longer be observed 6 days after thapsigargin treatment. These findings demonstrate that HIV-infection does not prevent the induction of the UPR and that infected macrophages are capable of responding to robust ER stress by increasing levels of ERSE regulated genes, such as BiP. Furthermore, robust activation of the UPR is capable of inhibiting viral replication, suggesting that HIV cannot efficiently escape the antiviral effects of the UPR, including translational inhibition and/or induction of ERAD. Since inhibition of virus replication has traditionally resulted in less neurotoxin production in HIV-infected macrophages (Figures 4.1 and 4.2), we assessed the consequence of the inhibition of viral replication due to thapsigargin treatment on macrophage-mediated neurotoxicity.

As previously discussed, attenuation of HIV replication with the nonnucleoside reverse transcriptase inhibitor (NNRTI), efavirenz (EFZ), significantly abrogates neurotoxin production in HIV/MDM (Figure 4.2 and Figure 5.5C). Interestingly, activation of the UPR and attenuation of HIV replication with thapsigargin treatment enhances neurotoxin production in HIV/MDM (Figure 5.5C). This demonstrates that the inhibition of virus replication does not always produce a concomitant decrease in neurotoxin production in HIV/MDM; an important implication for the development of adjunctive therapies for HAND. And as shown in Figure 5.5D, uninfected macrophages, which normally do not produce neurotoxins, cause significant neurotoxicity when treated with thapsigargin. These findings suggest that ER stress can modulate neurotoxin production in macrophages and that the elevations of phosphorylated eIF2 $\alpha$  with HIV infection (Figure 5.2) and increased BiP levels in HAND (Figure 5.3) may contribute to HIV-induced macrophage-mediated neurotoxicity.




**Figure 5.5 Thapsigargin attenuates HIV replication and enhances neurotoxin production in human MDM.** **A)** Thapsigargin (Thap) attenuates HIV replication in human MDM. MDM were infected with HIV-Jago and then treated for 6 hours on day 6 post infection with 0.5μM Thapsigargin. Following drug exposure, cells were washed and put into fresh drug-free media. Culture supernatants were collected every 2-3 days, as indicated, and assessed for RT activity. Values indicate the mean ± SEM. RT curves are representative of 3 independent experiments, with each replicate performed on cell preparations from a different donor. **B)** Thapsigargin treatment results in increased levels of phosphorylated eIF2α and BiP, as assessed by Western blotting of lysates collected after the 6 hour thapsigargin treatment on day 6 post infection (6+6). Thapsigargin treatment enhances neurotoxin production in **C)** HIV-infected and **D)** uninfected human macrophages. For C, culture supernatants from day 8 post infection were assayed. For D, uninfected macrophages were treated with 0.5μM of thapsigargin for 24hours followed by drug washout and media replacement. Culture supernatant from 72 hours after thapsigargin exposure was then assayed. Neuronal survival was assessed by MAP2 ELISA and data is expressed as a percentage of untreated (UT) cultures (n = 6). All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing, \*p<0.05, \*\*p<0.01 and \*\*\*p<0.001.

*Phosphorylated eIF2 $\alpha$  attenuates HIV replication and enhances neurotoxin production*

Thapsigargin treatment results in multiple cellular effects, including robust activation of the UPR signaling pathways that respond to the accumulated misfolded proteins in the ER. However, thapsigargin treatment also results in the depletion of ER Ca<sup>2+</sup> stores and potential alterations to Ca<sup>2+</sup> dependent kinases in the cytosol. In order to better define the role of the UPR in affecting HIV replication and macrophage-mediated neurotoxicity, more specific pharmacological modulators were used. Clotrimazole (CLT) disrupts ER calcium stores by inhibiting calcium-activated potassium (K<sub>Ca2+</sub>) channels, resulting in calcium release from intracellular stores of the ER. In contrast to thapsigargin, however, clotrimazole treatment of human macrophages does not induce the UPR, as assessed by Western blotting for phosphorylated eIF2 $\alpha$  and BiP (Figure 5.6C). Sal003, a derivative of salubrinal with enhanced cell permeability, indirectly inhibits the dephosphorylation of eIF2, resulting in robust activation of the UPR (Figure 5.5C) without directly affecting ER calcium levels or Ca<sup>2+</sup> dependent kinases. As shown in Figure 5.6A, clotrimazole has no major effect on HIV replication when added to HIV/MDM for 24 hours during an established infection. In contrast, Sal003 treatment inhibited HIV replication by approximately 25%, as compared to vehicle control. This suggests that HIV replication is moderately sensitive to phosphorylated eIF2 $\alpha$  levels, but not to disruptions in ER calcium homeostasis. While Sal003 most likely mediates the attenuation of HIV replication by inhibiting translation via induction of phosphorylated eIF2 $\alpha$ , further experiments are required to confirm the mechanism of action.

While clotrimazole treatment had minimal impact on HIV replication, supernatant collected from macrophages that were pulsed with clotrimazole 48 hours previously were more neurotoxic than the vehicle treated control (Figure 5.6B). Sal003 treatment, which attenuated HIV replication, also caused HIV-infected macrophages to produce more neurotoxins (Figure 5.6B). These results confirm that attenuation of HIV infection does not necessarily result in a concomitant decrease in neurotoxin production by macrophages. Furthermore, both the activation of the UPR via induction

of phosphorylated eIF2 $\alpha$  and disruption of ER calcium homeostasis are capable of enhancing neurotoxin production in HIV-infected macrophages.



**Figure 5.6 Pharmacological induction of phosphorylated eIF2 $\alpha$  and disruption of Ca<sup>2+</sup> stores in the endoplasmic reticulum enhances neurotoxin release in human MDM. A)** Sal003, a specific inhibitor of eIF2 $\alpha$  dephosphorylation, but not clotrimazole (CLT), an inducer of Ca<sup>2+</sup> release from the endoplasmic reticulum, attenuates HIV replication in human macrophages. MDM infected with HIV-Jago were pulse-treated with Sal003 or clotrimazole for 24 hours (day 6-7 post infection). Following drug exposure, cells were washed and put into fresh drug-free media. Culture supernatants were collected every 2-3 days, as indicated, and assessed for RT activity. Values indicate the mean  $\pm$  SEM. **B)** Sal003 and clotrimazole enhance neurotoxin production in HIV-infected human macrophages. Neuronal survival was assessed by MAP2 ELISA following exposure to macrophage supernatants from day 9-post infection, which had been treated with Sal003 or clotrimazole 48-hours previously. MAP2 data is expressed in arbitrary fluorescent units (n = 6). All statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing, \*\*p<0.01 and \*\*\*p<0.001. **C)** 0.5 $\mu$ M thapsigargin and 50 $\mu$ M Sal003 treatment (6 or 24-hours) results in increased levels of phosphorylated eIF2 $\alpha$  and BiP, while 10 $\mu$ M clotrimazole has little effect on these markers of the unfolded protein response (UPR), as assessed by Western blotting in uninfected human macrophages.


*Activation of the antioxidant response reduces levels of phosphorylated eIF2 $\alpha$  in HIV-infected macrophages*

In addition to misfolded protein accumulation in the ER, the UPR can also respond to oxidative stress (Holtz et al., 2006; Yan et al., 2008). Reactive oxygen species (ROS) can target ER calcium channels and protein folding enzymes to exacerbate ER calcium release and ER stress. As a consequence of ROS, unfolded or misfolded proteins can accumulate in the ER and induce the activation of the classic UPR pathways (Ron and Walter, 2007; Zhang and Kaufman, 2008). We have previously demonstrated that HIV infection results in the dysregulation of components of the antioxidant response in macrophages and that specifically, reductions in HO-1 levels contribute to macrophage-mediated neurotoxicity (Cross et al., 2011). We first sought to determine how restoration of the antioxidant response, through DMF and MMF treatment, affected levels of phosphorylated eIF2 $\alpha$  in HIV-infected macrophages. As shown in Figure 5.7, DMF and MMF treatment inhibited HIV replication and induced components of the antioxidant response (HO-1 and NQO1) in a dose-dependent manner. DMF and MMF treatment caused substantial decreases in phosphorylated eIF2 $\alpha$  in a dose-dependent manner, approaching levels seen in uninfected (mock) MDM with 30 $\mu$ M of DMF/MMF (Figure 5.7E and F). Additionally, DMF and MMF treatment also resulted in moderate decreases in BiP and GRP94, while total levels of eIF2 $\alpha$  remained unchanged. Since DMF/MMF treatment results in both the restoration of the antioxidant response (Figure 4.4) and attenuation of HIV replication in HIV/MDM, the observed decreases in phosphorylated eIF2 $\alpha$  could be a consequence of either effect.


In uninfected macrophages, MMF treatment results in the activation of the antioxidant response, as detected by increased levels of Nrf2 and NQO1 (Figure 5.8A). MMF treatment also decreases levels of GRP94 and BiP, without substantial alterations to levels of phosphorylated eIF2 $\alpha$  or total eIF2 $\alpha$  (Figure 5.8B). These findings suggest that DMF and MMF activate the antioxidant response and cause the concomitant down-regulation in expression of ERSE regulated genes, including BiP and GRP94, without significantly affecting levels of phosphorylated eIF2 $\alpha$ .



Therefore, in HIV-infected macrophages, DMF/MMF treatment likely decreases levels of phosphorylated eIF2 $\alpha$  secondary to attenuation of HIV replication. However, further experiments are necessary to demonstrate that phosphorylated eIF2 $\alpha$  levels are not varying with the redox state of the cell, since induction of HO-1 and NQO1 can correlate with a substantial decrease in phosphorylated eIF2 $\alpha$  levels without significant changes in the level of HIV replication (5 $\mu$ M versus 15 $\mu$ M DMF treatment in Figure 5.7).



**Figure 5.7 DMF and MMF activate the antioxidant response and decrease levels of phosphorylated eIF2 $\alpha$  in HIV-infected human macrophages.** **A)** DMF and **B)** MMF attenuate HIV replication in human MDM. MDM were infected with HIV-Jago and treated with DMF, MMF or the non-nucleoside reverse transcriptase inhibitor, efavirenz (EFZ), over the course of infection. Culture supernatants were collected every 2-3 days, as indicated, and assessed for reverse transcriptase (RT) activity. Values indicate the mean  $\pm$  SEM. RT curves are representative of 3 independent experiments, with each replicate performed on cell preparations from a different donor. **C)** DMF and **D)** MMF activate the Nrf2-dependent antioxidant response in HIV/MDM and restore HO-1 and GPX1 levels to that found in uninfected Mock cells. **E)** DMF and **F)** MMF decrease levels of phosphorylated eIF2 $\alpha$ , which is induced with HIV replication, as assessed by Western blotting. DMF and MMF treatment also decreases Grp94 and BiP levels in a dose-dependent manner. Blots are representative of 2-4 independent experiments, with each replicate performed on cell preparations from different donors.




**Figure 5.8 MMF activates the antioxidant response and decreases levels of Grp94 and BiP in uninfected human macrophages.** **A)** MMF activates the Nrf2-dependent antioxidant response in uninfected human macrophages, as demonstrated by increased levels of NQO1 and Nrf2. **B)** MMF treatment decreases Grp94 and BiP levels in a dose-dependent manner in uninfected human macrophages, without consistently affecting levels of phosphorylated eIF2 $\alpha$ . Blots are representative of 2 independent experiments, with each replicate performed on cell preparations from different donors.

*Levels of phosphorylated eIF2 $\alpha$  do not predict the extent of HIV replication or neurotoxin production in HIV-infected macrophages*

Components of the UPR and antioxidant response are both altered during HIV infection of macrophages. As we have previously demonstrated that decreased expression of HO-1 is associated with high levels of HIV replication and neurotoxin production in HIV/MDM (Cross et al., 2011), we examined levels of phosphorylated eIF2 $\alpha$  in conditions of high viral replication but altered neurotoxin production. As shown in Figure 5.9, restoration of HO-1 levels with CoPP, an inducer of HO-1 expression, decreases macrophage-mediated neurotoxicity without altering levels of replication. In contrast, enzymatic inhibition of HO-1 with SnMP results in enhanced neurotoxin production in the absence of alterations to HIV replication. Interestingly, both SnMP and CoPP treatment decreased levels of phosphorylated eIF2 $\alpha$ , relative to vehicle controls, without causing significant alterations to GRP94, BiP or total eIF2 $\alpha$  levels (Figure 5.9D). This

suggests that phosphorylated eIF2 $\alpha$  levels do not necessarily correlate with HIV replication or with neurotoxin production in HIV-infected macrophages. Furthermore, these results suggest that pharmacological modulation of the antioxidant response, even in the setting of high viral replication and enhanced neurotoxin production, is sufficient to decrease levels of phosphorylated eIF2 $\alpha$ . CoPP and SnMP are relatively specific for their actions on HO-1, however, both drugs also moderately increase NQO1 levels, suggesting that they have off-target effects on other components of the antioxidant response element (ARE). SnMP treatment also causes a reflexive increase in total HO-1 protein levels (Figure 5.9E), and while the enzymatic activity of HO-1 is inhibited, HO-1 may alter the redox state of the cell or dampen activation of phosphorylated eIF2 $\alpha$  via a non-enzymatic mechanism. In fact, total HO-1 levels may specifically correlate with phosphorylated eIF2 $\alpha$  since unlike DMF and MMF treatment, CoPP and SnMP had no effect on levels of the ERSE regulated chaperones, BiP and GRP94. The relationship between the antioxidant response and the UPR is complex and has not been fully elucidated for any cell type or paradigm. However, these experiments suggest that induction of the antioxidant response dampens the UPR in HIV-infected macrophages, although the underlying mechanisms and consequences of this interaction require more study.



**Figure 5.9 Levels of phosphorylated eIF2α do not correlate with HIV replication or neurotoxin production in HIV-infected human macrophages.** **A)** SnMP, an enzymatic inhibitor of HO-1, and CoPP, an inducer of HO-1, do not alter levels of HIV replication. MDM were infected with HIV-Jago and then treated with SnMP or CoPP from day 6 post infection onwards and culture supernatants were collected every 3 days and assessed for RT activity. RT curves are representative of 3 independent experiments, with each replicate performed on cell preparations from different donors. **B)** SnMP enhances neurotoxin production while **(C)** CoPP decreases neurotoxin production in HIV-infected macrophages. For neuronal survival assays, survival was assessed by MAP2 ELISA and expressed as a percentage of untreated (UT) cultures (n = 6; \*\*\*p<0.001 vs. Vehicle). Statistical comparisons were made by one-way ANOVA plus Newman-Keuls post hoc testing. **D)** SnMP and CoPP decrease levels of phosphorylated eIF2α, as assessed by Western blotting. **E)** SnMP inhibits HO-1 enzymatic activity but exponentially increases total levels of HO-1 and modestly increases NQO1, as assessed by Western blotting. **F)** CoPP treatment exponentially increases HO-1 levels and modestly increases NQO1 levels, as assessed by Western blotting. Western blots are representative of 2-3 independent experiments, with each replicate performed on cell preparations from different donors. Two film exposures (short and long) of HO-1 are presented in order to demonstrate the extent of HO-1 induction over basal levels.

## Discussion

Viral infection can induce the activation of host cell survival pathways, which may have direct antiviral effects and/or promote cell survival during sustained ER stress, alterations to the cellular redox state, and viral infection. We have demonstrated that HIV infection of human macrophages results in the activation of phosphorylated eIF2 $\alpha$  and BiP, two critical components of the UPR, and that activation of the UPR correlates with macrophage-mediated neurotoxicity. In our *in vitro* model system, HIV infected macrophages have elevated levels of phosphorylated eIF2 $\alpha$  during times of high viral replication but expression of the ERSE regulated genes, BiP and GRP94, is not increased. However, we found a significant increase in the amount of macrophagic-BiP in HAND frontal cortex, demonstrating that the UPR and ERSE regulated genes are activated in HIV infected brain tissue, suggesting an association with human pathology. Both our *in vitro* and *in vivo* data demonstrate activation of the UPR during HIV infection of the macrophage. Importantly, in our *in vitro* model system the majority of macrophages are robustly infected with HIV, while less than 5% of macrophages in the CNS are estimated to be infected during HAND. Furthermore, HAND pathology is mediated by both infected and non-infected, activated macrophages with both populations driving neuroinflammation and producing excitatory neurotoxins. Therefore, primary HIV infection of the macrophage may increase levels of phosphorylated eIF2 $\alpha$  without activation of ERSE regulated genes. In contrast, immune activation of non-infected (or chronic, low-level replicating) macrophages may induce expression of ERSE genes, which may be more sensitive to upregulation in inflammatory states. In order to address these hypotheses, we plan to assess for the activation of phosphorylated eIF2 $\alpha$  and ERSE regulated genes in uninfected macrophages that have been exposed to virus-free HIV/MDM supernatant in our *in vitro* model system. In this way, we can determine how the UPR is modulated in uninfected macrophages exposed to a pro-inflammatory environment induced by HIV-infection. We are also continuing to evaluate how the three signaling pathways comprising the UPR (PERK, IRE1 and ATF6) respond to HIV infection and correlate to macrophage-mediated

neurotoxicity. Given limitations to available antibodies and the difficulty of performing efficient genetic manipulations in primary human macrophages, future studies will utilize qPCR.

Importantly, we have demonstrated that conditions that induce robust activation of the UPR, with alterations in ERSE gene expression, are associated with the enhancement of neurotoxin production in macrophages (Figure 5.5 and 5.6). This suggests that activation of ER stress pathways can augment neurotoxin production pathways, even in uninfected macrophages. Interestingly, we have noted that in our *in vitro* model system, HIV-infected macrophages produce neurotoxins late in the course of infection. The extent of neurotoxin production varies with the donor and as previously noted, increases with robust HIV replication. However, macrophage-mediated neurotoxicity also correlates with the appearance of increased phosphorylated eIF2 $\alpha$  and decreased levels of HO-1, as assessed by Western blotting. While this may be a coincidental observation, it is consistent with the hypothesis that neurotoxin production in the macrophage is enhanced under states of extreme cellular stress, as indicated by dysregulation of the antioxidant response and activation of the UPR. Furthermore, these findings have important implications for the development of adjunctive therapeutics for HAND. Reduction of HIV replication has been the principal target for ameliorating the pathological consequences of HIV-infection. And while ART has greatly improved clinical outcomes for patients infected with HIV, our data demonstrate that drugs that activate the UPR may attenuate HIV replication, but will also enhance macrophage-mediated neurotoxicity. In addition to physiological stressors like viral infection and inflammation, there have been reports that ART, specifically protease inhibitors, can activate the UPR in macrophages (Zhou et al., 2005). The role of ART-induced activation of the UPR in macrophages and the consequence to macrophage-mediated neurotoxicity has not yet been studied and remain a focus for future research.

Activation of the UPR, especially phosphorylated eIF2 $\alpha$ , during viral infection may be part of a cellular antiviral defense that serves to limit virus replication in the host cell. We have

demonstrated that HIV infection activates components of the UPR and that pharmacological induction of the UPR can attenuate HIV replication. This suggests that during HIV infection, the induction of phosphorylated eIF2 $\alpha$  likely has some antiviral effect, but that HIV replication is relatively efficient despite this. In fact, selective induction of phosphorylated eIF2 $\alpha$  with the pharmacological agent Sal003 (Figure 5.6) resulted in only a 25% inhibition of viral replication when applied for 24 hours. In contrast, a 6 hour treatment with thapsigargin resulted in a 70% inhibition of viral replication. These results suggest that high levels of phosphorylated eIF2 $\alpha$  can attenuate HIV replication, to an extent, but that robust activation of all components of the UPR has a more substantial effect on viral replication. Perhaps, thapsigargin treatment induces the IRE1-XBP1 signaling pathway to upregulate EDEM and ERAD, thereby inducing an antiviral component of the UPR that HIV infection does not. It also remains possible that HIV is capable of escaping the translational inhibition normally imposed by activation of phosphorylated eIF2 $\alpha$ .

In order to understand how induction of the UPR alters HIV replication, there are several key experiments that need to be undertaken. One of the outstanding questions for this work is whether the level of phosphorylated eIF2 $\alpha$  induced with HIV infection is sufficient to attenuate host cell protein translation. And, to determine how sensitive uninfected macrophages are to phosphorylated eIF2 $\alpha$ -mediated inhibition of protein translation. In HIV-infected macrophages, we have seen no evidence of protein instability by Western blotting to suggest that the levels of phosphorylated eIF2 $\alpha$  induced with infection have a substantial effect on protein translation. However, we must determine if this is due to viral modulation of the UPR and/or cellular translational machinery or an intrinsic property of the macrophage. To address this question, we are planning to perform metabolic labeling experiments ( $^{35}\text{S}$ -methionine and  $^{35}\text{S}$ -cysteine) on uninfected and HIV-infected macrophages, both in the presence and absence of pharmacological modulators of the UPR (thapsigargin, Sal003 and cyclohexamide). We will also use qPCR to determine if HIV-induced induction of phosphorylated eIF2 $\alpha$  results in the transcriptional upregulation of ATF4, and if ATF4 transcriptional targets and CHOP are upregulated.



If we find that HIV-infected macrophages are less sensitive to the inhibition of protein translation as mediated by phosphorylated eIF2 $\alpha$ , we must then consider the mechanism by which HIV modulates this component of the UPR. The 5' leader of the genomic RNA of HIV has an IRES (Brasey et al., 2003), and it is possible that HIV replication is capable of efficient translation even in conditions of phosphorylated eIF2 $\alpha$ -mediated inhibition of protein translation, in a mechanism similar to other stress-responsive mRNAs (Fernandez et al., 2002). Independent of these findings, the relationship between UPR-activation following HIV infection and neurotoxin production in the macrophage will also be explored.

Following HIV infection of the CNS, infected and non-infected/activated macrophages and glia release neurotoxins that result in neuronal dendritic damage, apoptosis and necrosis (Adamson et al., 1996; Dawson et al., 1993; Gelbard et al., 1995; Kaul et al., 2001; O'Donnell et al., 2006; Petit and Roberts, 1995). These neurotoxins include HIV viral proteins, proinflammatory cytokines, chemokines, and excitatory amino acids (Chen et al., 2002; Dawson et al., 1993; Kaul et al., 2001; Power et al., 1998; Tardieu et al., 1992). Excitatory neurotoxins that mediate neuronal damage via NMDA receptor activation are the major mediators of HIV-induced neurotoxicity. These low-molecular weight excitatory neurotoxins include glutamate, quinolinic acid, platelet activating factor, reactive oxygen species and NTox. In our *in vitro* system, glutamate has been determined to mediate a significant component of HIV/MDM-mediated neurotoxicity, with other low-molecular weight heat resistant species also contributing (O'Donnell et al., 2006). This correlates to *in vivo* findings, where HIV-infected patients have been found to have significantly higher concentrations of glutamate in their plasma and CSF, compared to uninfected controls (Droge et al., 1987; Ferrarese et al., 2001; Ollenschlager et al., 1988). Glutaminase, which is localized to the inner membrane of the mitochondria, is the primary enzyme involved in converting glutamine into glutamate by macrophages in the CNS (Curthoys and Watford, 1995; Holcomb et al., 2000; Nicklas et al., 1987; Wurdig and Kugler, 1991). HIV

infection results in the upregulation of the GAC isoform of glutaminase and its release into the cytosol of macrophages, presumably due to HIV-induced destabilization of the mitochondrial membrane (Erdmann et al., 2009). Interestingly, sustained activation of the UPR results in caspase-12 activation and cellular apoptosis mediated by mitochondrial outer membrane permeabilization via BAX and/or BAK. Therefore, activation of the UPR by HIV-infection or pharmacological treatment may result in enhanced macrophage-mediated neurotoxicity by compromising the mitochondria membrane and increasing glutaminase-mediated glutamate production. It is also possible for ER stress to directly affect the mitochondria through calcium and reactive oxygen species, thereby altering mitochondrial processes without inducing apoptosis pathways (Zhang and Kaufman, 2008).

In addition to investigating glutamate production during UPR activation in HIV-infected macrophages, we will also examine quinolinic acid (QUIN). QUIN is a neurotoxin derived from tryptophan through the kynurenine metabolic pathway. HIV infection increases levels and the activity of indoleamine-2,3-dioxygenase (IDO), the rate-limiting enzyme of the kynurenine pathway, which leads to increased metabolism of tryptophan and the enhanced generation of neurotoxins such as QUIN (Guillemin et al., 2003; Sardar et al., 1995). Generation of QUIN by IDO is limited to activated macrophages and microglia in the CNS and IDO is induced by exposure to interferon- $\gamma$  or during infection in order to deplete cellular tryptophan and limit metabolic processes (Carlin et al., 1989). Furthermore, TNF $\alpha$  levels can alter IDO activity, suggesting that any process that alters the inflammatory state and TNF $\alpha$  production can ultimately modulate IDO and QUIN production in the macrophage (Werner-Felmayer et al., 1989). UPR stress can directly increase inflammation and the production of proinflammatory mediators (Zhang and Kaufman, 2008). Activation of IRE1 can induce a signaling cascade involving JNK and the transcription factor AP1, resulting in the transcription of genes involved in the inflammatory response. Additionally, phosphorylated eIF2 $\alpha$ -mediated inhibition of protein

translation can result in the destabilization of I $\kappa$ B, promoting the nuclear translocation of NF- $\kappa$ B proteins and inducing the expression of TNF $\alpha$  and other pro-inflammatory genes.

Interactions between ER stress, oxidative stress and the inflammatory response are necessary and crucial for cell types with high metabolic and secretory demands, such as macrophages (Gregor and Hotamisligil, 2007; Schenk et al., 2008; Zhang and Kaufman, 2008), in order to cope with the cellular consequences of high levels of protein translation. Numerous stimuli such as glucose, lipids, and cytokines can activate both the UPR and the inflammatory pathways through calcium, reactive oxygen species, and/or nitric oxide mediators. We have demonstrated that HIV-infection of human macrophages results in the suppression of components of the antioxidant response and activation of the UPR and that alterations to these pathways affect neurotoxin production following HIV infection. Interestingly, restoration of the antioxidant response attenuates HIV replication, decreases neurotoxin production and dampens induction of phosphorylated eIF2 $\alpha$  (Figure 5.7). However, decreased neurotoxin production is associated with restoration of HO-1 levels and not to levels of phosphorylated eIF2 $\alpha$  or the ERSE regulated genes, Bip and GRP94. Phosphorylated eIF2 $\alpha$  levels can be dampened by restoring the antioxidant response in HIV-infected macrophages, even in the setting of robust HIV replication. These data suggest that reducing phosphorylated eIF2 $\alpha$  will not necessarily attenuate HIV replication or neurotoxin production. In fact, phosphorylated eIF2 $\alpha$  and activation of the UPR in HIV-infected macrophages may better correlate to overall cellular health, redox state or levels of inflammatory signaling. In HIV-infected macrophages, activation of the UPR likely enhances neurotoxin production, but our studies suggest that treatments that dampen activation of phosphorylated eIF2 $\alpha$  will not be effective as adjunctive therapies in HAND. However, more study is needed to understand the role of phosphorylated eIF2 $\alpha$  during HIV infection, especially as it intersects with inflammatory pathways and neurotoxin production in macrophages.

## **CHAPTER 6**

### **DISCUSSION**

### *Summary of Work*

Despite antiretroviral therapy (ART), HIV infection of the central nervous system (CNS) promotes cognitive dysfunction and neurodegeneration through persistent inflammation and the release of neurotoxic factors from infected and/or activated macrophages/microglia. The persistence of HAND in individuals effectively controlled for systemic viral replication is incompletely explained, although prolonged inflammation in both the CNS and periphery may be responsible (Ancuta et al., 2008; Brenchley et al., 2006b; Eden et al., 2007). Because markers of inflammation and immune activation correlate with disease progression and morbidity in ART-treated individuals, drugs targeting these pathological processes are needed for effective, adjunctive therapy. This work begins to address the roles of two adaptive stress pathways, the antioxidant response and the unfolded protein response (UPR), in modulating macrophage-mediated neurotoxicity following HIV infection.

Using our *in vitro* model of HIV-mediated neurotoxicity, we have shown that HIV infection dysregulates the macrophage antioxidant response, with profound reductions in levels of heme oxygenase-1 (HO-1). Induction of the antioxidant response with DMF, MMF or tBHQ attenuates HIV replication in a dose-dependent manner. Our data suggest that decreased levels of the HIV co-receptors, CXCR4 and CCR5, mediate attenuation of viral replication following the induction of the antioxidant response. We have also demonstrated that activation of the antioxidant response and restoration of HO-1 expression, specifically, reduces HIV-induced macrophage-mediated neurotoxicity. Importantly, we demonstrate that therapeutics that increase HO-1 levels, even during times of robust HIV replication, can suppress neurotoxin production. In addition to effects on macrophage-mediated neurotoxicity, inducers of the antioxidant response can also modulate inflammatory pathways and may decrease the immune activation state of HIV-infected macrophages. We have demonstrated that induction of the antioxidant response with DMF, MMF and tBHQ inhibits of NF- $\kappa$ B nuclear translocation, signaling and the production of the proinflammatory mediator, TNF $\alpha$ . Finally, we found that DMF and MMF treatment attenuates

CCL2-induced monocyte chemotaxis, suggesting that therapeutic inducers of the antioxidant response could decrease recruitment of activated monocytes to the CNS in response to inflammatory mediators. We propose that dysregulation of the antioxidant response during HIV infection drives macrophage-mediated neurotoxicity and that inducers of the antioxidant response, such as DMF, could serve as adjunctive neuroprotectants and HIV disease modifiers in ART-treated individuals.

In addition to investigating the antioxidant response in HIV-infected macrophage, we also assessed the consequence of HIV infection on the unfolded protein response (UPR). Using our *in vitro* model system, we found that HIV infection increases levels of phosphorylated eIF2 $\alpha$  in human macrophages. We also demonstrate that the chaperone BiP, considered the master regulator of the UPR, is increased in the macrophages of HAND frontal cortex. HIV-infected macrophages are capable of responding to pharmacological induction of the UPR and induction of the UPR attenuates HIV replication. These results suggest that induction of phosphorylated eIF2 $\alpha$  during HIV replication has antiviral consequences, although activation of the UPR it is not sufficient to considerably attenuate HIV replication during the normal course of infection in macrophages. Interestingly, pharmacological induction of the UPR, which attenuates viral replication, is associated with increased macrophage-mediated neurotoxicity. This finding has important implications for the development of adjunctive therapies for HAND. Therapeutics or processes that induce the UPR in macrophages, regardless of the effect on HIV replication, could enhance neurotoxin production and contribute to the pathological processes underlying HAND.

As we have also described the effect of HIV replication on the antioxidant response, we assessed how restoration of the antioxidant response altered the UPR in HIV-infected macrophages. Activation of the antioxidant response with DMF and MMF attenuated HIV replication and decreased levels of phosphorylated eIF2 $\alpha$  and BiP in HIV/MDM. Reductions in phosphorylated eIF2 $\alpha$  trend with increased levels of HO-1, but do not depend upon HO-1 enzymatic activity. In

contrast, BiP levels decreased with induction of the antioxidant response, but these alterations were independent of HO-1. These findings suggest that while dysregulation of the antioxidant response reflects levels of HIV replication and HO-1 levels modulate neurotoxin production, levels of phosphorylated eIF2 $\alpha$  and BiP do not predict the extent of HIV replication or macrophage-mediated neurotoxicity. Ongoing studies are assessing the link between UPR activation, inflammation pathways and the antioxidant response as they relate to HIV infection and macrophage-mediated neurotoxicity.

Developing adjunctive therapies for HAND requires an understanding of the pathological processes underlying the disease. The macrophage is a critical mediator of HAND neuropathology and is a critical mediator of inflammation and neurotoxin production within the CNS. This body of work focused on the roles of the antioxidant response and the UPR in modulating HIV replication, macrophage-mediated neurotoxicity and inflammatory processes. These experiments have demonstrated that induction of the antioxidant response, but not the UPR, should be considered as a potential target for adjunctive therapies for HAND. Our lab is continuing its focus on the role of the antioxidant response in modulating the neurological consequences of HIV infection and is planning to apply this work to animal model systems and hopefully, human clinical trials in the future.

### *Implications*

Despite highly active antiretroviral therapies (ART) and effective virological control, approximately 50% of HIV-infected individuals will develop neurological consequences of HIV infection. In order to develop effective adjunctive therapies for HAND, we must consider and target the pathological consequences of HIV infection that persist in the CNS despite ART. Chronic immune activation, inflammation and viral persistence all contribute to neuronal damage and one of the principal cell types mediating these processes is the macrophage. While previous studies have explored the pathways responsible for neurotoxin production, little is known about the mechanisms linking HIV infection to these pathways. We hypothesized that HIV infection alters many components of cellular metabolism and signaling within the infected macrophage. Specifically, we focused on adaptive stress pathways, which have been implicated in the modulation of viral replication and viral-mediated pathology in recent years. We concentrated on the antioxidant response, due to clinical data demonstrating alterations to the oxidative state in HIV-infected patients, and the unfolded protein response (UPR), because of its clear link to inflammation pathways and its involvement during infection by many other viruses. The effect of HIV infection on the antioxidant response, the UPR and the interactions between these adaptive stress responses and inflammation had not yet been explored.

This work has begun to address the roles of the antioxidant response and UPR in modulating HIV-induced macrophage mediated neurotoxicity and has implications for the development of adjunctive therapy for HAND. Namely, we demonstrate that restoration of the antioxidant response in HIV-infected macrophages can attenuate many of the key pathways contributing to the neuropathology of HAND including HIV replication, neurotoxin production, inflammatory signaling and chemotaxis. In order to develop more targeted therapies, we are continuing to explore how the antioxidant response affects neurotoxin production pathways and are concentrating on the role of HO-1 and its enzymatic activity. HO-1 reduces neurotoxin production in macrophages, even when viral replication is robust, suggesting that it may have a direct effect



on or modulate a critical component of neurotoxin production pathways. There are also reports that HO-1 plays a role in determining the immune activation state of cells, although this is still at a rather speculative stage and will require more experimentation. However, when considering how the antioxidant response, inflammation and neurotoxin production pathways modulate and are influenced by one other, we will need to consider the immune activation state, or profile, of the HIV-infected macrophages in our *in vitro* system.

The microbial and cytokine milieu drives macrophages, T-cells, and other cell types of the immune system to express specialized and polarized functional properties. Interestingly, it has been reported that MMF shifts CD4<sup>+</sup> T-cells towards a Th2 phenotype, which inhibits proinflammatory signaling (de Jong et al., 1996; Litjens et al., 2004). DMF and MMF likely also influence the polarization of activated and HIV-infected macrophages. TNF $\alpha$  exposure elicits a classic M1 form of macrophage activation, which results in the secretion of high levels of proinflammatory cytokines (TNF $\alpha$ , IL-1 and IL-6), and reactive oxygen and nitrogen intermediates (Mantovani, 2006). In contrast, M2 is a generic name for various forms of macrophage activation, which encompass all but M1-activated cells. The various versions of M2 cells generally dampen proinflammatory cytokine levels (IL-4R $\alpha$ , IL-10, IL-1 receptor antagonist<sup>high</sup>, decoy IL-1 type II receptor<sup>high</sup>) and have high levels of scavenger, mannose, and galactose-type receptors. DMF may promote a shift from M1 activated macrophages towards a M2 phenotype. M2 HIV-infected macrophages would produce less TNF $\alpha$ , have less NF- $\kappa$ B signaling and may have decreased transcription from the HIV-LTR. It is also highly likely that this immune activation state and alterations to inflammation pathways and cell surface receptors will influence signaling events related to neurotoxin production pathways. The mechanism by which DMF and MMF shift the activation state of immune cells towards an anti-inflammatory state and the impact of activation state on neurotoxin production remains to be determined.

With this body of work, we have also demonstrated the antioxidant response is a better therapeutic target in regards to reducing macrophage-mediated neurotoxicity than targeting the UPR. It has been well established that activation of the UPR can drive inflammatory pathways and the production of proinflammatory mediators (Ron and Walter, 2007; Zhang and Kaufman, 2008). And although pharmacological activation of the UPR attenuates HIV replication, it also enhances neurotoxin production, making it a poor therapeutic target for HAND. These data have implications for how we think about continued inflammation and macrophage-mediated neurotoxicity in patients experiencing neurological complications of HIV. Patients who are well controlled for viral replication through ART can still have neuronal damage and neurological impairment. We demonstrate that HAND patients have evidence of UPR activation in the frontal cortex, as evidenced by elevated levels of macrophagic BiP. And as we have demonstrated that UPR activation enhances neurotoxin production, even if HIV replication is low or absent, these results suggest that activation of the UPR may be driving neurotoxin production in macrophages even in the setting of ART. Even more importantly, our data demonstrate that simply controlling HIV replication does not guarantee a concomitant decrease in macrophage-mediated neurotoxicity, a concept that has not been previously proposed. Furthermore, we need to consider how potential therapies impact the UPR of the macrophage when we consider adjunctive therapies for HAND, since controlling viral replication is not sufficient to prevent neurological sequelae or ensure a reduction in neurotoxicity.

This body of work has proposed many future avenues of research in regards to the role of adaptive stress responses in macrophage-mediated neurotoxicity. One of the key questions that remains in this work is how the antioxidant response, UPR and neurotoxin pathways intersect. It has been demonstrated that intracellular calcium signals and free radicals, such as reactive oxygen species (ROS) and nitric oxide (NO), act as messengers in coordinating adaptive stress responses. However, we have yet to determine if HIV infection directly impacts the antioxidant response or UPR, or if the markers of modulation/activation that we detect are an indirect

consequence of infection or mediated by other signaling pathways. It is possible that the antioxidant response drives the activation of the UPR, increases inflammation and directly modulates neurotoxin production pathways. However, it is also a possibility that activation of the UPR or inflammation drives the antioxidant response and neurotoxin production. In order to develop targeted and effective adjunctive therapies for HAND, we must continue to explore these pathways and identify critical mediators of the pathology.

## BIBLIOGRAPHY

(2011). Trial watch: Phase III success for Biogen's oral multiple sclerosis therapy. *Nat Rev Drug Discov* 10, 404.

Achim, C.L., Heyes, M.P., and Wiley, C.A. (1993). Quantitation of human immunodeficiency virus, immune activation factors, and quinolinic acid in AIDS brains. *J Clin Invest* 91, 2769-2775.

Adamson, D.C., Wildemann, B., Sasaki, M., Glass, J.D., McArthur, J.C., Christov, V.I., Dawson, T.M., and Dawson, V.L. (1996). Immunologic NO synthase: elevation in severe AIDS dementia and induction by HIV-1 gp41. *Science* 274, 1917-1921.

Adle-Biassette, H., Chretien, F., Wingertsmann, L., Hery, C., Ereau, T., Scaravilli, F., Tardieu, M., and Gray, F. (1999). Neuronal apoptosis does not correlate with dementia in HIV infection but is related to microglial activation and axonal damage. *Neuropathol Appl Neurobiol* 25, 123-133.

Alirezaei, M., Kiosses, W.B., Flynn, C.T., Brady, N.R., and Fox, H.S. (2008). Disruption of neuronal autophagy by infected microglia results in neurodegeneration. *PLoS One* 3, e2906.

Ambrose, R.L., and Mackenzie, J.M. (2011). West Nile virus differentially modulates the unfolded protein response to facilitate replication and immune evasion. *J Virol* 85, 2723-2732.

Ances, B.M., Letendre, S.L., Alexander, T., and Ellis, R.J. (2008). Role of psychiatric medications as adjunct therapy in the treatment of HIV associated neurocognitive disorders. *Int Rev Psychiatry* 20, 89-93.

Ancuta, P., Kamat, A., Kunstman, K.J., Kim, E.Y., Autissier, P., Wurcel, A., Zaman, T., Stone, D., Mefford, M., Morgello, S., *et al.* (2008). Microbial translocation is associated with increased monocyte activation and dementia in AIDS patients. *PLoS One* 3, e2516.

Ancuta, P., Rao, R., Moses, A., Mehle, A., Shaw, S.K., Luscinskas, F.W., and Gabuzda, D. (2003). Fractalkine preferentially mediates arrest and migration of CD16+ monocytes. *J Exp Med* 197, 1701-1707.

Ancuta, P., Wang, J., and Gabuzda, D. (2006). CD16+ monocytes produce IL-6, CCL2, and matrix metalloproteinase-9 upon interaction with CX3CL1-expressing endothelial cells. *J Leukoc Biol* 80, 1156-1164.

Anderson, E., Zink, W., Xiong, H., and Gendelman, H.E. (2002). HIV-1-associated dementia: a metabolic encephalopathy perpetrated by virus-infected and immune-competent mononuclear phagocytes. *J Acquir Immune Defic Syndr* 31 Suppl 2, S43-54.

Anderson, E.R., Gendelman, H.E., and Xiong, H. (2004). Memantine protects hippocampal neuronal function in murine human immunodeficiency virus type 1 encephalitis. *J Neurosci* 24, 7194-7198.

Anthony, I.C., and Bell, J.E. (2008). The Neuropathology of HIV/AIDS. International review of psychiatry 20, 15-24.

- Anthony, I.C., Ramage, S.N., Carnie, F.W., Simmonds, P., and Bell, J.E. (2005). Influence of HAART on HIV-related CNS disease and neuroinflammation. *J Neuropathol Exp Neurol* **64**, 529-536.
- Antinori, A., Arendt, G., Becker, J.T., Brew, B.J., Byrd, D.A., Cherner, M., Clifford, D.B., Cinque, P., Epstein, L.G., Goodkin, K., *et al.* (2007). Updated research nosology for HIV-associated neurocognitive disorders. *Neurology* **69**, 1789-1799.
- Asehnoune, K., Strassheim, D., Mitra, S., Yeol Kim, J., and Abraham, E. (2005). Involvement of PKC[alpha]/[beta] in TLR4 and TLR2 dependent activation of NF- $\kappa$ B. *Cellular Signalling* **17**, 385-394.
- Asensio, V.C., Maier, J., Milner, R., Boztug, K., Kincaid, C., Moulard, M., Phillipson, C., Lindsley, K., Krucker, T., Fox, H.S., *et al.* (2001). Interferon-independent, human immunodeficiency virus type 1 gp120-mediated induction of CXCL10/IP-10 gene expression by astrocytes in vivo and in vitro. *J Virol* **75**, 7067-7077.
- Bajetto, A., Bonavia, R., Barbero, S., Florio, T., and Schettini, G. (2001). Chemokines and their receptors in the central nervous system. *Front Neuroendocrinol* **22**, 147-184.
- Bajetto, A., Bonavia, R., Barbero, S., Piccioli, P., Costa, A., Florio, T., and Schettini, G. (1999). Glial and neuronal cells express functional chemokine receptor CXCR4 and its natural ligand stromal cell-derived factor 1. *J Neurochem* **73**, 2348-2357.
- Baldwin, A.S. (1996). The NF- $\kappa$ B and I $\kappa$ B Proteins: New Discoveries and Insights. *Annual Review of Immunology* **14**, 649-681.
- Bandaru, V.V., McArthur, J.C., Sacktor, N., Cutler, R.G., Knapp, E.L., Mattson, M.P., and Haughey, N.J. (2007). Associative and predictive biomarkers of dementia in HIV-1-infected patients. *Neurology* **68**, 1481-1487.
- Banks, W.A., Ercal, N., and Price, T.O. (2006). The blood-brain barrier in neuroAIDS. *Curr HIV Res* **4**, 259-266.
- Barger, S.W., Goodwin, M.E., Porter, M.M., and Beggs, M.L. (2007). Glutamate release from activated microglia requires the oxidative burst and lipid peroxidation. *J Neurochem* **101**, 1205-1213.
- Bartosik-Psujek, H., and Stelmasiak, Z. (2005). Correlations between IL-4, IL-12 levels and CCL2, CCL5 levels in serum and cerebrospinal fluid of multiple sclerosis patients. *J Neural Transm* **112**, 797-803.
- Bell, J.E. (2004). An update on the neuropathology of HIV in the HAART era. *Histopathology* **45**, 549-559.
- Berlanga, J.J., Ventoso, I., Harding, H.P., Deng, J., Ron, D., Sonenberg, N., Carrasco, L., and de Haro, C. (2006). Antiviral effect of the mammalian translation initiation factor 2alpha kinase GCN2 against RNA viruses. *Eur Mol Biol Org J* **25**, 1730-1740.
- Bertolotti, A., Zhang, Y., Hendershot, L.M., Harding, H.P., and Ron, D. (2000). Dynamic interaction of BiP and ER stress transducers in the unfolded-protein response. *Nat Cell Biol* **2**, 326-332.

Bezzi, P., Domercq, M., Brambilla, L., Galli, R., Schols, D., De Clercq, E., Vescovi, A., Bagetta, G., Kollias, G., Meldolesi, J., *et al.* (2001). CXCR4-activated astrocyte glutamate release via TNF $\alpha$ : amplification by microglia triggers neurotoxicity. *Nat Neurosci* 4, 702-710.

Biddison, W.E., Kubota, R., Kawanishi, T., Taub, D.D., Cruikshank, W.W., Center, D.M., Connor, E.W., Utz, U., and Jacobson, S. (1997). Human T cell leukemia virus type I (HTLV-I)-specific CD8<sup>+</sup> CTL clones from patients with HTLV-I-associated neurologic disease secrete proinflammatory cytokines, chemokines, and matrix metalloproteinase. *J Immunol* 159, 2018-2025.

Bitko, V., and Barik, S. (2001). An endoplasmic reticulum-specific stress-activated caspase (caspase-12) is implicated in the apoptosis of A549 epithelial cells by respiratory syncytial virus. *J Cell Biochem* 80, 441-454.

Block, M.L., and Hong, J.S. (2005). Microglia and inflammation-mediated neurodegeneration: multiple triggers with a common mechanism. *Prog Neurobiol* 76, 77-98.

Bodner, A., Maroney, A.C., Finn, J.P., Ghadge, G., Roos, R., and Miller, R.J. (2002). Mixed lineage kinase 3 mediates gp120IIIB-induced neurotoxicity. *J Neurochem* 82, 1424-1434.

Boisse, L., Gill, M.J., and Power, C. (2008). HIV infection of the central nervous system: clinical features and neuropathogenesis. *Neurol Clin* 26, 799-819, x.

Boscoboinik, D., Szewczyk, A., Hensey, C., and Azzi, A. (1991). Inhibition of cell proliferation by alpha-tocopherol. Role of protein kinase C. *J Biol Chem* 266, 6188-6194.

Boven, L.A., Gomes, L., Hery, C., Gray, F., Verhoef, J., Portegies, P., Tardieu, M., and Nottet, H.S. (1999). Increased peroxynitrite activity in AIDS dementia complex: implications for the neuropathogenesis of HIV-1 infection. *J Immunol* 162, 4319-4327.

Brasey, A., Lopez-Lastra, M., Ohlmann, T., Beerens, N., Berkhout, B., Darlix, J.L., and Sonenberg, N. (2003). The leader of human immunodeficiency virus type 1 genomic RNA harbors an internal ribosome entry segment that is active during the G2/M phase of the cell cycle. *J Virol* 77, 3939-3949.

Breen, E., Rezai, A., Nakajima, K., Beall, G., Mitsuyasu, R., Hirano, T., Kishimoto, T., and Martinez-Maza, O. (1990). Infection with HIV is associated with elevated IL-6 levels and production. *J Immunol* 144, 480-484.

Breitner, J.C., Welsh, K.A., Helms, M.J., Gaskell, P.C., Gau, B.A., Roses, A.D., Pericak-Vance, M.A., and Saunders, A.M. (1995). Delayed onset of Alzheimer's disease with nonsteroidal anti-inflammatory and histamine H2 blocking drugs. *Neurobiol Aging* 16, 523-530.

Brenchley, J.M., and Douek, D.C. (2008). HIV infection and the gastrointestinal immune system. *Mucosal Immunol* 1, 23-30.

Brenchley, J.M., Price, D.A., and Douek, D.C. (2006a). HIV disease: fallout from a mucosal catastrophe? *Nat Immunol* 7, 235-239.

Brenchley, J.M., Price, D.A., Schacker, T.W., Asher, T.E., Silvestri, G., Rao, S., Kazzaz, Z., Bornstein, E., Lambotte, O., Altmann, D., *et al.* (2006b). Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med* 12, 1365-1371.

- Brenneman, D.E., Westbrook, G.L., Fitzgerald, S.P., Ennist, D.L., Elkins, K.L., Ruff, M.R., and Pert, C.B. (1988). Neuronal cell killing by the envelope protein of HIV and its prevention by vasoactive intestinal peptide. *Nature* 335, 639-642.
- Brew, B.J. (2004). Evidence for a change in AIDS dementia complex in the era of highly active antiretroviral therapy and the possibility of new forms of AIDS dementia complex. *AIDS* 18 Suppl 1, S75-78.
- Brew, B.J., Bhalla, R.B., Paul, M., Sidtis, J.J., Keilp, J.J., Sadler, A.E., Gallardo, H., McArthur, J.C., Schwartz, M.K., and Price, R.W. (1992). Cerebrospinal fluid beta 2-microglobulin in patients with AIDS dementia complex: an expanded series including response to zidovudine treatment. *AIDS* 6, 461-465.
- Brew, B.J., Corbeil, J., Pemberton, L., Evans, L., Saito, K., Penny, R., Cooper, D.A., and Heyes, M.P. (1995). Quinolinic acid production is related to macrophage tropic isolates of HIV-1. *J Neurovirol* 1, 369-374.
- Brew, B.J., Crowe, S.M., Landay, A., Cysique, L.A., and Guillemin, G. (2009). Neurodegeneration and ageing in the HAART era. *J Neuroimmune Pharmacol* 4, 163-174.
- Brew, B.J., Dunbar, N., Pemberton, L., and Kaldor, J. (1996). Predictive markers of AIDS dementia complex: CD4 cell count and cerebrospinal fluid concentrations of beta 2-microglobulin and neopterin. *J Infect Dis* 174, 294-298.
- Brew, B.J., Halman, M., Catalan, J., Sacktor, N., Price, R.W., Brown, S., Atkinson, H., Clifford, D.B., Simpson, D., Torres, G., *et al.* (2007). Factors in AIDS dementia complex trial design: results and lessons from the abacavir trial. *PLoS Clin Trials* 2, e13.
- Brew, B.J., Pemberton, L., Cunningham, P., and Law, M.G. (1997). Levels of human immunodeficiency virus type 1 RNA in cerebrospinal fluid correlate with AIDS dementia stage. *J Infect Dis* 175, 963-966.
- Bruno, V., Copani, A., Besong, G., Scoto, G., and Nicoletti, F. (2000). Neuroprotective activity of chemokines against N-methyl-D-aspartate or beta-amyloid-induced toxicity in culture. *Eur J Pharmacol* 399, 117-121.
- Brush, M.H., Weiser, D.C., and Shenolikar, S. (2003). Growth arrest and DNA damage-inducible protein GADD34 targets protein phosphatase 1 alpha to the endoplasmic reticulum and promotes dephosphorylation of the alpha subunit of eukaryotic translation initiation factor 2. *Mol Cell Biol* 23, 1292-1303.
- Burdo, T.H., Ellis, R.J., and Fox, H.S. (2008). Osteopontin is increased in HIV-associated dementia. *J Infect Dis* 198, 715-722.
- Butera, S., Roberts, B., and Folks, T. (1993). Regulation of HIV-1 expression by cytokine networks in a CD4+ model of chronic infection. *J Immunol* 150, 625-634.
- Cannella, B., and Raine, C.S. (1995). The adhesion molecule and cytokine profile of multiple sclerosis lesions. *Ann Neurol* 37, 424-435.
- Cardona, A.E., Pioro, E.P., Sasse, M.E., Kostenko, V., Cardona, S.M., Dijkstra, I.M., Huang, D., Kidd, G., Dombrowski, S., Dutta, R., *et al.* (2006). Control of microglial neurotoxicity by the fractalkine receptor. *Nat Neurosci* 9, 917-924.

- Carlin, J.M., Borden, E.C., and Byrne, G.I. (1989). Interferon-induced indoleamine 2,3-dioxygenase activity inhibits *Chlamydia psittaci* replication in human macrophages. *J Interferon Res* 9, 329-337.
- Casoli, T., Di Stefano, G., Baliotti, M., Solazzi, M., Giorgetti, B., and Fattoretti, P. (2010). Peripheral inflammatory biomarkers of Alzheimer's disease: the role of platelets. *Biogerontology*.
- Chalovich, E.M., Koike, M.A., Aras, M.A., Murphey-Corb, M., Wiley, C.A., and Jordan-Sciutto, K.L. (2005). Pocket proteins p107 and p130 exhibit increased expression in macrophages during SIV encephalitis. *Neuropathology* 25, 315-325.
- Chang, L., Ernst, T., St Hillaire, C., and Conant, K. (2004). Antiretroviral treatment alters relationship between MCP-1 and neurometabolites in HIV patients. *Antivir Ther* 9, 431-440.
- Chang, L., Wong, V., Nakama, H., Watters, M., Ramones, D., Miller, E.N., Cloak, C., and Ernst, T. (2008). Greater than age-related changes in brain diffusion of HIV patients after 1 year. *J Neuroimmune Pharmacol* 3, 265-274.
- Chapman, G.A., Moores, K., Harrison, D., Campbell, C.A., Stewart, B.R., and Strijbos, P.J. (2000a). Fractalkine cleavage from neuronal membranes represents an acute event in the inflammatory response to excitotoxic brain damage. *J Neurosci* 20, RC87.
- Chapman, G.A., Moores, K.E., Gohil, J., Berkhout, T.A., Patel, L., Green, P., Macphee, C.H., and Stewart, B.R. (2000b). The role of fractalkine in the recruitment of monocytes to the endothelium. *Eur J Pharmacol* 392, 189-195.
- Chen, H., Zhang, S.M., Hernan, M.A., Schwarzschild, M.A., Willett, W.C., Colditz, G.A., Speizer, F.E., and Ascherio, A. (2003). Nonsteroidal anti-inflammatory drugs and the risk of Parkinson disease. *Arch Neurol* 60, 1059-1064.
- Chen, J.J., and London, I.M. (1995). Regulation of protein synthesis by heme-regulated eIF-2 alpha kinase. *Trends Biochem Sci* 20, 105-108.
- Chen, T.H., Tang, P., Yang, C.F., Kao, L.H., Lo, Y.P., Chuang, C.K., Shih, Y.T., and Chen, W.J. (2011). Antioxidant defense is one of the mechanisms by which mosquito cells survive dengue 2 viral infection. *Virology* 410, 410-417.
- Chen, W., Sulcove, J., Frank, I., Jaffer, S., Ozdener, H., and Kolson, D.L. (2002). Development of a human neuronal cell model for human immunodeficiency virus (HIV)-infected macrophage-induced neurotoxicity: apoptosis induced by HIV type 1 primary isolates and evidence for involvement of the Bcl-2/Bcl-xL-sensitive intrinsic apoptosis pathway. *J Virol* 76, 9407-9419.
- Cheng, G., Feng, Z., and He, B. (2005). Herpes simplex virus 1 infection activates the endoplasmic reticulum resident kinase PERK and mediates eIF-2alpha dephosphorylation by the gamma(1)34.5 protein. *J Virol* 79, 1379-1388.
- Childs, E.A., Lyles, R.H., Selnes, O.A., Chen, B., Miller, E.N., Cohen, B.A., Becker, J.T., Mellors, J., and McArthur, J.C. (1999). Plasma viral load and CD4 lymphocytes predict HIV-associated dementia and sensory neuropathy. *Neurology* 52, 607-613.
- Chiou, W.F., Ko, H.C., and Wei, B.L. (2011). Evodia rutaecarpa and Three Major Alkaloids Abrogate Influenza A Virus (H1N1)-Induced Chemokines Production and Cell Migration. *Evid Based Complement Alternat Med* 2011, 750513.



- Choe, W., Volsky, D.J., and Potash, M.J. (2001). Induction of rapid and extensive beta-chemokine synthesis in macrophages by human immunodeficiency virus type 1 and gp120, independently of their coreceptor phenotype. *J Virol* 75, 10738-10745.
- Ciardi, M., Sharief, M.K., Thompson, E.J., Salotti, A., Vullo, V., Sorice, F., and Cirelli, A. (1994). High cerebrospinal fluid and serum levels of tumor necrosis factor-[alpha] in asymptomatic HIV-1 seropositive individuals. Correlation with interleukin-2 and soluble IL-2 receptor. *J Neurol Sci* 125, 175-179.
- Cinque, P., Bestetti, A., Marenzi, R., Sala, S., Gisslen, M., Hagberg, L., and Price, R.W. (2005). Cerebrospinal fluid interferon-gamma-inducible protein 10 (IP-10, CXCL10) in HIV-1 infection. *J Neuroimmunol* 168, 154-163.
- Cinque, P., Vago, L., Mengozzi, M., Torri, V., Ceresa, D., Vicenzi, E., Transidico, P., Vagani, A., Sozzani, S., Mantovani, A., *et al.* (1998). Elevated cerebrospinal fluid levels of monocyte chemoattractant protein-1 correlate with HIV-1 encephalitis and local viral replication. *AIDS* 12, 1327-1332.
- Circle, D.A., Neel, O.D., Robertson, H.D., Clarke, P.A., and Mathews, M.B. (1997). Surprising specificity of PKR binding to delta agent genomic RNA. *Rna* 3, 438-448.
- Clifford, D.B., Fagan, A.M., Holtzman, D.M., Morris, J.C., Teshome, M., Shah, A.R., and Kauwe, J.S. (2009). CSF biomarkers of Alzheimer disease in HIV-associated neurologic disease. *Neurology* 73, 1982-1987.
- Cocchi, F., DeVico, A.L., Garzino-Demo, A., Arya, S.K., Gallo, R.C., and Lusso, P. (1995). Identification of RANTES, MIP-1 alpha, and MIP-1 beta as the major HIV-suppressive factors produced by CD8+ T cells. *Science* 270, 1811-1815.
- Collman, R.G., and Yi, Y. (1999). Cofactors for human immunodeficiency virus entry into primary macrophages. *J Infect Dis* 179 Suppl 3, S422-426.
- Colovic, M., and Caccia, S. (2003). Liquid chromatographic determination of minocycline in brain-to-plasma distribution studies in the rat. *J Chromatogr B Analyt Technol Biomed Life Sci* 791, 337-343.
- Conant, K., Garzino-Demo, A., Nath, A., McArthur, J.C., Halliday, W., Power, C., Gallo, R.C., and Major, E.O. (1998). Induction of monocyte chemoattractant protein-1 in HIV-1 Tat-stimulated astrocytes and elevation in AIDS dementia. *Proc Natl Acad Sci USA* 95, 3117-3121.
- Copeland, K.F., and Brooks, J.I. (2010). A novel use for an old drug: the potential for minocycline as anti-HIV adjuvant therapy. *J Infect Dis* 201, 1115-1117.
- Cota, M., Kleinschmidt, A., Ceccherini-Silberstein, F., Aloisi, F., Mengozzi, M., Mantovani, A., Brack-Werner, R., and Poli, G. (2000). Upregulated expression of interleukin-8, RANTES and chemokine receptors in human astrocytic cells infected with HIV-1. *J Neurovirol* 6, 75-83.
- Cotter, R., Williams, C., Ryan, L., Erichsen, D., Lopez, A., Peng, H., and Zheng, J. (2002). Fractalkine (CX3CL1) and brain inflammation: Implications for HIV-1-associated dementia. *J Neurovirol* 8, 585-598.
- Craft, J.M., Watterson, D.M., and Van Eldik, L.J. (2006). Human amyloid beta-induced neuroinflammation is an early event in neurodegeneration. *Glia* 53, 484-490.

- Cristofanon, S., Morceau, F., Scovassi, A.I., Dicato, M., Ghibelli, L., and Diederich, M. (2009). Oxidative, multistep activation of the noncanonical NF- $\kappa$ B pathway via disulfide Bcl-3/p50 complex. *Fed Am Soc Exp Biol J* 23, 45-57.
- Cross, S., Cook, D., Chi, A., Vance, P., Kolson, L., Wong, B., Jordan-Sciutto, K., and Kolson, D. (2011). Dimethyl fumarate, an immune modulator and inducer of the antioxidant response, suppresses HIV replication and macrophage-mediated neurotoxicity; a novel candidate for HIV-neuroprotection. *J Immunol* 187.
- Crowe, S., Zhu, T., and Muller, W.A. (2003). The contribution of monocyte infection and trafficking to viral persistence, and maintenance of the viral reservoir in HIV infection. *J Leukocyte Biol* 74, 635-641.
- Curthoys, N.P., and Watford, M. (1995). Regulation of glutaminase activity and glutamine metabolism. *Annu Rev Nutr* 15, 133-159.
- D'Aversa, T.G., Yu, K.O., and Berman, J.W. (2004). Expression of chemokines by human fetal microglia after treatment with the human immunodeficiency virus type 1 protein Tat. *J Neurovirol* 10, 86-97.
- Dallasta, L.M., Pisarov, L.A., Esplen, J.E., Werley, J.V., Moses, A.V., Nelson, J.A., and Achim, C.L. (1999). Blood-brain barrier tight junction disruption in human immunodeficiency virus-1 encephalitis. *American J Pathol* 155, 1915-1927.
- Dawson, V.L., Dawson, T.M., Uhl, G.R., and Snyder, S.H. (1993). Human immunodeficiency virus type 1 coat protein neurotoxicity mediated by nitric oxide in primary cortical cultures. *Proc Natl Acad Sci USA* 90, 3256-3259.
- De Long, M.J., Santamaria, A.B., and Talalay, P. (1987). Role of cytochrome P1-450 in the induction of NAD(P)H:quinone reductase in a murine hepatoma cell line and its mutants. *Carcinogenesis* 8, 1549-1553.
- Debebe, Z., Ammosova, T., Jerebtsova, M., Kurantsin-Mills, J., Niu, X., Charles, S., Richardson, D.R., Ray, P.E., Gordeuk, V.R., and Nekhai, S. (2007). Iron chelators ICL670 and 311 inhibit HIV-1 transcription. *Virology* 367, 324-333.
- Deiva, K., Geeraerts, T., Salim, H., Leclerc, P., Hery, C., Hugel, B., Freyssinet, J.M., and Tardieu, M. (2004). Fractalkine reduces N-methyl-D-aspartate-induced calcium flux and apoptosis in human neurons through extracellular signal-regulated kinase activation. *Eur J Neurosci* 20, 3222-3232.
- Devadas, K., and Dhawan, S. (2006). Hemin activation ameliorates HIV-1 infection via heme oxygenase-1 induction. *J Immunol* 176, 4252-4257.
- Devadas, K., Hewlett, I.K., and Dhawan, S. (2010). Lipopolysaccharide suppresses HIV-1 replication in human monocytes by protein kinase C-dependent heme oxygenase-1 induction. *J Leukocyte Biol* 87, 915-924.
- Dhillon, N., Zhu, X., Peng, F., Yao, H., Williams, R., Qiu, J., Callen, S., Ladner, A.O., and Buch, S. (2008a). Molecular mechanism(s) involved in the synergistic induction of CXCL10 by human immunodeficiency virus type 1 Tat and interferon-gamma in macrophages. *J Neurovirol* 14, 196-204.

- Dhillon, N.K., Williams, R., Callen, S., Zien, C., Narayan, O., and Buch, S. (2008b). Roles of MCP-1 in development of HIV-dementia. *Front Biosci* 13, 3913-3918.
- Dimcheff, D.E., Askovic, S., Baker, A.H., Johnson-Fowler, C., and Portis, J.L. (2003). Endoplasmic reticulum stress is a determinant of retrovirus-induced spongiform neurodegeneration. *J Virol* 77, 12617-12629.
- Doms, R.W. (2000). Beyond receptor expression: the influence of receptor conformation, density, and affinity in HIV-1 infection. *Virology* 276, 229-237.
- Dore-Duffy, P., Newman, W., Balabanov, R., Lisak, R.P., Mainolfi, E., Rothlein, R., and Peterson, M. (1995). Circulating, soluble adhesion proteins in cerebrospinal fluid and serum of patients with multiple sclerosis: correlation with clinical activity. *Ann Neurol* 37, 55-62.
- Dou, H., Birusingh, K., Faraci, J., Gorantla, S., Poluektova, L.Y., Maggirwar, S.B., Dewhurst, S., Gelbard, H.A., and Gendelman, H.E. (2003). Neuroprotective activities of sodium valproate in a murine model of human immunodeficiency virus-1 encephalitis. *J Neurosci* 23, 9162-9170.
- Douek, D. (2007). HIV disease progression: immune activation, microbes, and a leaky gut. *Top HIV Med* 15, 114-117.
- Droge, W., Eck, H.P., Betzler, M., and Naher, H. (1987). Elevated plasma glutamate levels in colorectal carcinoma patients and in patients with acquired immunodeficiency syndrome (AIDS). *Immunobiology* 174, 473-479.
- Drulovic, J., Mostarica-Stojkovic, M., Levic, Z., Mesaros, S., Stojavljevic, N., Popadic, D., and Pravica, V. (1998). Serum interleukin-12 levels in patients with multiple sclerosis. *Neurosci Lett* 251, 129-132.
- Drulovic, J., Mostarica-Stojkovic, M., Levic, Z., Stojavljevic, N., Pravica, V., and Mesaros, S. (1997). Interleukin-12 and tumor necrosis factor-alpha levels in cerebrospinal fluid of multiple sclerosis patients. *J Neurol Sci* 147, 145-150.
- Duh, E.J., Maury, W.J., Folks, T.M., Fauci, A.S., and Rabson, A.B. (1989). Tumor necrosis factor alpha activates human immunodeficiency virus type 1 through induction of nuclear factor binding to the NF-kappa B sites in the long terminal repeat. *Proc Natl Acad Sci USA* 86, 5974-5978.
- Dunfee, R., Thomas, E.R., Gorry, P.R., Wang, J., Ancuta, P., and Gabuzda, D. (2006). Mechanisms of HIV-1 neurotropism. *Curr HIV Res* 4, 267-278.
- Dworkin, B.M., Rosenthal, W.S., Wormser, G.P., and Weiss, L. (1986). Selenium deficiency in the acquired immunodeficiency syndrome. *J Parenter Enteral Nutr* 10, 405-407.
- Earl, P.L., Moss, B., and Doms, R.W. (1991). Folding, interaction with GRP78-BiP, assembly, and transport of the human immunodeficiency virus type 1 envelope protein. *J Virol* 65, 2047-2055.
- Eden, A., Price, R.W., Spudich, S., Fuchs, D., Hagberg, L., and Gisslen, M. (2007). Immune activation of the central nervous system is still present after >4 years of effective highly active antiretroviral therapy. *J Infectious Dis* 196, 1779-1783.
- Edgil, D., Polacek, C., and Harris, E. (2006). Dengue virus utilizes a novel strategy for translation initiation when cap-dependent translation is inhibited. *J Virol* 80, 2976-2986.

- Eggert, D., Dash, P.K., Gorantla, S., Dou, H., Schifitto, G., Maggirwar, S.B., Dewhurst, S., Poluektova, L., Gelbard, H.A., and Gendelman, H.E. (2009a). Neuroprotective Activities of CEP-1347 in Models of NeuroAIDS. *J Immunol*.
- Eggert, D., Dash, P.K., Serradji, N., Dong, C.Z., Clayette, P., Heymans, F., Dou, H., Gorantla, S., Gelbard, H.A., Poluektova, L., *et al.* (2009b). Development of a platelet-activating factor antagonist for HIV-1 associated neurocognitive disorders. *J Neuroimmunol* 213, 47-59.
- Eikelenboom, P., Bate, C., Van Gool, W.A., Hoozemans, J.J., Rozemuller, J.M., Veerhuis, R., and Williams, A. (2002). Neuroinflammation in Alzheimer's disease and prion disease. *Glia* 40, 232-239.
- El-Hage, N., Gurwell, J.A., Singh, I.N., Knapp, P.E., Nath, A., and Hauser, K.F. (2005). Synergistic increases in intracellular Ca<sup>2+</sup>, and the release of MCP-1, RANTES, and IL-6 by astrocytes treated with opiates and HIV-1 Tat. *Glia* 50, 91-106.
- Elia, A., Laing, K.G., Schofield, A., Tilleray, V.J., and Clemens, M.J. (1996). Regulation of the double-stranded RNA-dependent protein kinase PKR by RNAs encoded by a repeated sequence in the Epstein-Barr virus genome. *Nucleic Acids Res* 24, 4471-4478.
- Ellery, P.J., Tippet, E., Chiu, Y.L., Paukovics, G., Cameron, P.U., Solomon, A., Lewin, S.R., Gorry, P.R., Jaworowski, A., Greene, W.C., *et al.* (2007). The CD16<sup>+</sup> monocyte subset is more permissive to infection and preferentially harbors HIV-1 in vivo. *J Immunol* 178, 6581-6589.
- Ellis, R., Langford, D., and Masliah, E. (2007). HIV and antiretroviral therapy in the brain: neuronal injury and repair. *Nat Rev Neurosci* 8, 33-44.
- Ellis, R.J., Gamst, A.C., Capparelli, E., Spector, S.A., Hsia, K., Wolfson, T., Abramson, I., Grant, I., and McCutchan, J.A. (2000). Cerebrospinal fluid HIV RNA originates from both local CNS and systemic sources. *Neurology* 54, 927-936.
- Ellis, R.J., Hsia, K., Spector, S.A., Nelson, J.A., Heaton, R.K., Wallace, M.R., Abramson, I., Atkinson, J.H., Grant, I., and McCutchan, J.A. (1997). Cerebrospinal fluid human immunodeficiency virus type 1 RNA levels are elevated in neurocognitively impaired individuals with acquired immunodeficiency syndrome. HIV Neurobehavioral Research Center Group. *Ann Neurol* 42, 679-688.
- Ellis, R.J., Moore, D.J., Childers, M.E., Letendre, S., McCutchan, J.A., Wolfson, T., Spector, S.A., Hsia, K., Heaton, R.K., and Grant, I. (2002). Progression to neuropsychological impairment in human immunodeficiency virus infection predicted by elevated cerebrospinal fluid levels of human immunodeficiency virus RNA. *Arch Neurol* 59, 923-928.
- Elovaara, I., Koenig, S., Brewah, A.Y., Woods, R.M., Lehky, T., and Jacobson, S. (1993). High human T cell lymphotropic virus type 1 (HTLV-1)-specific precursor cytotoxic T lymphocyte frequencies in patients with HTLV-1-associated neurological disease. *J Exp Med* 177, 1567-1573.
- Enose-Akahata, Y., Matsuura, E., Oh, U., and Jacobson, S. (2009). High expression of CD244 and SAP regulated CD8 T cell responses of patients with HTLV-I associated neurologic disease. *PLoS Pathog* 5, e1000682.
- Enting, R.H., Foudraine, N.A., Lange, J.M., Jurriaans, S., van der Poll, T., Weverling, G.J., and Portegies, P. (2000). Cerebrospinal fluid beta2-microglobulin, monocyte chemotactic protein-1,

and soluble tumour necrosis factor alpha receptors before and after treatment with lamivudine plus zidovudine or stavudine. *J Neuroimmunol* 102, 216-221.

Erdmann, N., Tian, C., Huang, Y., Zhao, J., Herek, S., Curthoys, N., and Zheng, J. (2009). In vitro glutaminase regulation and mechanisms of glutamate generation in HIV-1-infected macrophage. *J Neurochem* 109, 551-561.

Erichsen, D., Lopez, A.L., Peng, H., Niemann, D., Williams, C., Bauer, M., Morgello, S., Cotter, R.L., Ryan, L.A., Ghorpade, A., *et al.* (2003). Neuronal injury regulates fractalkine: relevance for HIV-1 associated dementia. *J Neuroimmunol* 138, 144-155.

Esiri, M.M., Biddolph, S.C., and Morris, C.S. (1998). Prevalence of Alzheimer plaques in AIDS. *J Neurol Neurosurg Psychiatry* 65, 29-33.

Esser, R., Glienke, W., von Briesen, H., Rubsamen-Waigmann, H., and Andreesen, R. (1996). Differential regulation of proinflammatory and hematopoietic cytokines in human macrophages after infection with human immunodeficiency virus. *Blood* 88, 3474-3481.

Eugenin, E.A., D'Aversa, T.G., Lopez, L., Calderon, T.M., and Berman, J.W. (2003). MCP-1 (CCL2) protects human neurons and astrocytes from NMDA or HIV-tat-induced apoptosis. *J Neurochem* 85, 1299-1311.

Eugenin, E.A., Osiecki, K., Lopez, L., Goldstein, H., Calderon, T.M., and Berman, J.W. (2006). CCL2/monocyte chemoattractant protein-1 mediates enhanced transmigration of human immunodeficiency virus (HIV)-infected leukocytes across the blood-brain barrier: a potential mechanism of HIV-CNS invasion and NeuroAIDS. *J Neurosci* 26, 1098-1106.

Everall, I.P., Bell, C., Mallory, M., Langford, D., Adame, A., Rockenstein, E., and Masliah, E. (2002). Lithium ameliorates HIV-gp120-mediated neurotoxicity. *Mol Cell Neurosci* 21, 493-501.

Everall, I.P., Luthert, P.J., and Lantos, P.L. (1991). Neuronal loss in the frontal cortex in HIV infection. *Lancet* 337, 1119-1121.

Fawcett, T.W., Martindale, J.L., Guyton, K.Z., Hai, T., and Holbrook, N.J. (1999). Complexes containing activating transcription factor (ATF)/cAMP-responsive-element-binding protein (CREB) interact with the CCAAT/enhancer-binding protein (C/EBP)-ATF composite site to regulate Gadd153 expression during the stress response. *Biochem J* 339 (Pt 1), 135-141.

Feldman, D.E., Chauhan, V., and Koong, A.C. (2005). The unfolded protein response: a novel component of the hypoxic stress response in tumors. *Mol Cancer Res* 3, 597-605.

Fernandez, J., Bode, B., Koromilas, A., Diehl, J.A., Krukovets, I., Snider, M.D., and Hatzoglou, M. (2002). Translation mediated by the internal ribosome entry site of the cat-1 mRNA is regulated by glucose availability in a PERK kinase-dependent manner. *J Biol Chem* 277, 11780-11787.

Ferrarese, C., Aliprandi, A., Tremolizzo, L., Stanzani, L., De Micheli, A., Dolaro, A., and Frattola, L. (2001). Increased glutamate in CSF and plasma of patients with HIV dementia. *Neurology* 57, 671-675.

Ferri, K.F., and Kroemer, G. (2001). Organelle-specific initiation of cell death pathways. *Nat Cell Biol* 3, E255-263.

Fiala, M., Looney, D.J., Stins, M., Way, D.D., Zhang, L., Gan, X., Chiappelli, F., Schweitzer, E.S., Shapshak, P., Weinand, M., *et al.* (1997). TNF-alpha opens a paracellular route for HIV-1 invasion across the blood-brain barrier. *Mol Med* 3, 553-564.

Fischer-Smith, T., Croul, S., Adeniyi, A., Rybicka, K., Morgello, S., Khalili, K., and Rappaport, J. (2004). Macrophage/microglial accumulation and proliferating cell nuclear antigen expression in the central nervous system in human immunodeficiency virus encephalopathy. *Am J Pathol* 164, 2089-2099.

Foli, A., Saville, M.W., May, L.T., Webb, D.S., and Yarchoan, R. (1997). Effects of human immunodeficiency virus and colony-stimulating factors on the production of interleukin 6 and tumor necrosis factor alpha by monocyte/macrophages. *AIDS Res Hum Retroviruses* 13, 829-839.

Folks, T., Kessler, S., Orenstein, J., Justement, J., Jaffe, E., and Fauci, A. (1988). Infection and replication of HIV-1 in purified progenitor cells of normal human bone marrow. *Science* 242, 919-922.

Fraser, C.S., and Doudna, J.A. (2007). Structural and mechanistic insights into hepatitis C viral translation initiation. *Nat Rev Microbiol* 5, 29-38.

Frei, K., Fredrikson, S., Fontana, A., and Link, H. (1991). Interleukin-6 is elevated in plasma in multiple sclerosis. *J Neuroimmunol* 31, 147-153.

Friedman, A.D. (1996). GADD153/CHOP, a DNA damage-inducible protein, reduced CAAT/enhancer binding protein activities and increased apoptosis in 32D c13 myeloid cells. *Cancer Res* 56, 3250-3256.

Frohman, E.M., Racke, M.K., and Raine, C.S. (2006). Multiple sclerosis--the plaque and its pathogenesis. *N Engl J Med* 354, 942-955.

Ganchi, P.A., Sun, S.C., Greene, W.C., and Ballard, D.W. (1992). I kappa B/MAD-3 masks the nuclear localization signal of NF-kappa B p65 and requires the transactivation domain to inhibit NF-kappa B p65 DNA binding. *Mol Biol Cell* 3, 1339-1352.

Gartner, S. (2000). HIV infection and dementia. *Science* 287, 602-604.

Gass, J.N., Gifford, N.M., and Brewer, J.W. (2002). Activation of an unfolded protein response during differentiation of antibody-secreting B cells. *J Biol Chem* 277, 49047-49054.

Gelbard, H.A., Dzenko, K.A., DiLoreto, D., del Cerro, C., del Cerro, M., and Epstein, L.G. (1993). Neurotoxic effects of tumor necrosis factor alpha in primary human neuronal cultures are mediated by activation of the glutamate AMPA receptor subtype: implications for AIDS neuropathogenesis. *Dev Neurosci* 15, 417-422.

Gelbard, H.A., James, H.J., Sharer, L.R., Perry, S.W., Saito, Y., Kazee, A.M., Blumberg, B.M., and Epstein, L.G. (1995). Apoptotic neurons in brains from paediatric patients with HIV-1 encephalitis and progressive encephalopathy. *Neuropathology and applied neurobiology* 21, 208-217.

Gelbard, H.A., Nottet, H.S.L.M., Swindells, S., Jett, M., Dzenko, K.A., Genis, P., White, R., Wang, L., Choi, Y.-B., Zhang, D., *et al.* (1994). Platelet-Activating Factor: a candidate human immunodeficiency virus type 1-induced neurotoxin. *J Virol* 68, 4628-4635.

- Gendelman, H.E., Lipton, S.A., Tardieu, M., Bukrinsky, M.I., and Nottet, H.S. (1994). The neuropathogenesis of HIV-1 infection. *J Leukoc Biol* 56, 389-398.
- Gendelman, H.E., Zheng, J., Coulter, C.L., Ghorpade, A., Che, M., Thylin, M., Rubocki, R., Persidsky, Y., Hahn, F., Reinhard, J., Jr., *et al.* (1998). Suppression of inflammatory neurotoxins by highly active antiretroviral therapy in human immunodeficiency virus-associated dementia. *J Infectious Dis* 178, 1000-1007.
- Genis, P., Jett, M., Bernton, E.W., Boyle, T., Gelbard, H.A., Dzenko, K., Keane, R.W., Resnick, L., Mizrachi, Y., Volsky, D.J., *et al.* (1992). Cytokines and arachidonic metabolites produced during human immunodeficiency virus (HIV)-infected macrophage-astroglia interactions: implications for the neuropathogenesis of HIV disease. *J Exp Med* 176, 1703-1718.
- Glass, J.D., Fedor, H., Wesselingh, S.L., and McArthur, J.C. (1995). Immunocytochemical quantitation of human immunodeficiency virus in the brain: correlations with dementia. *Ann Neurol* 38, 755-762.
- Gonzalez-Scarano, F., and Martin-Garcia, J. (2005). The neuropathogenesis of AIDS. *Nat Rev Immunol* 5, 69-81.
- Gopalakrishna, R., Chen, Z.-H., and Gundimeda, U. (1995). Modifications of cysteine-rich regions in protein kinase C induced by oxidant tumor promoters and enzyme-specific inhibitors. In *Methods in Enzymology*, P. Lester, ed. (Academic Press), pp. 132-146.
- Gopalakrishna, R., Gundimeda, U., and Chen, Z.-H. (1997). Cancer-Preventive Selenocompounds Induce a Specific Redox Modification of Cysteine-Rich Regions in Ca<sup>2+</sup>-Dependent Isoenzymes of Protein Kinase C. *Archives Biochem Biophysics* 348, 25-36.
- Gorantla, S., Liu, J., Sneller, H., Dou, H., Holguin, A., Smith, L., Ikezu, T., Volsky, D.J., Poluektova, L., and Gendelman, H.E. (2007). Copolymer-1 induces adaptive immune anti-inflammatory glial and neuroprotective responses in a murine model of HIV-1 encephalitis. *J Immunol* 179, 4345-4356.
- Gorantla, S., Liu, J., Wang, T., Holguin, A., Sneller, H.M., Dou, H., Kipnis, J., Poluektova, L., and Gendelman, H.E. (2008). Modulation of innate immunity by copolymer-1 leads to neuroprotection in murine HIV-1 encephalitis. *Glia* 56, 223-232.
- Gradi, A., Svitkin, Y.V., Imataka, H., and Sonenberg, N. (1998). Proteolysis of human eukaryotic translation initiation factor eIF4GII, but not eIF4GI, coincides with the shutoff of host protein synthesis after poliovirus infection. *Proc Natl Acad Sci USA* 95, 11089-11094.
- Grant, C., Barmak, K., Alefantis, T., Yao, J., Jacobson, S., and Wigdahl, B. (2002). Human T cell leukemia virus type I and neurologic disease: events in bone marrow, peripheral blood, and central nervous system during normal immune surveillance and neuroinflammation. *J Cell Physiol* 190, 133-159.
- Grant, I. (2008). Neurocognitive disturbances in HIV. *Int Rev Psychiatry* 20, 33-47.
- Gray, F., Chretien, F., Vallat-Decouvelaere, A.V., and Scaravilli, F. (2003). The changing pattern of HIV neuropathology in the HAART era. *J Neuropathol Exp Neurol* 62, 429-440.
- Gregor, M.F., and Hotamisligil, G.S. (2007). Thematic review series: Adipocyte Biology. Adipocyte stress: the endoplasmic reticulum and metabolic disease. *J Lipid Res* 48, 1905-1914.

- Greten, T.F., Slansky, J.E., Kubota, R., Soldan, S.S., Jaffee, E.M., Leist, T.P., Pardoll, D.M., Jacobson, S., and Schneck, J.P. (1998). Direct visualization of antigen-specific T cells: HTLV-1 Tax11-19- specific CD8(+) T cells are activated in peripheral blood and accumulate in cerebrospinal fluid from HAM/TSP patients. *Proc Natl Acad Sci USA* 95, 7568-7573.
- Griffin, D.E., Wesselingh, S.L., and McArthur, J.C. (1994). Elevated central nervous system prostaglandins in human immunodeficiency virus-associated dementia. *Ann Neurol* 35, 592-597.
- Gu, L., Rutledge, B., Fiorillo, J., Ernst, C., Grewal, I., Flavell, R., Gladue, R., and Rollins, B. (1997). In vivo properties of monocyte chemoattractant protein-1. *J Leukoc Biol* 62, 577-580.
- Guillemin, G.J., Croitoru-Lamoury, J., Dormont, D., Armati, P.J., and Brew, B.J. (2003). Quinolinic acid upregulates chemokine production and chemokine receptor expression in astrocytes. *Glia* 41, 371-381.
- Habata, Y., Fujii, R., Hosoya, M., Fukusumi, S., Kawamata, Y., Hinuma, S., Kitada, C., Nishizawa, N., Murosaki, S., Kurokawa, T., *et al.* (1999). Apelin, the natural ligand of the orphan receptor APJ, is abundantly secreted in the colostrum. *Biochim Biophys Acta* 1452, 25-35.
- Hakansson, A., Westberg, L., Nilsson, S., Buervenich, S., Carmine, A., Holmberg, B., Sydow, O., Olson, L., Johnels, B., Eriksson, E., *et al.* (2005a). Interaction of polymorphisms in the genes encoding interleukin-6 and estrogen receptor beta on the susceptibility to Parkinson's disease. *Am J Med Genet B Neuropsychiatr Genet* 133B, 88-92.
- Hakansson, A., Westberg, L., Nilsson, S., Buervenich, S., Carmine, A., Holmberg, B., Sydow, O., Olson, L., Johnels, B., Eriksson, E., *et al.* (2005b). Investigation of genes coding for inflammatory components in Parkinson's disease. *Mov Disord* 20, 569-573.
- Harding, H.P., Novoa, I., Zhang, Y., Zeng, H., Wek, R., Schapira, M., and Ron, D. (2000a). Regulated translation initiation controls stress-induced gene expression in mammalian cells. *Mol Cell* 6, 1099-1108.
- Harding, H.P., Zhang, Y., Bertolotti, A., Zeng, H., and Ron, D. (2000b). Perk is essential for translational regulation and cell survival during the unfolded protein response. *Mol Cell* 5, 897-904.
- Harding, H.P., Zhang, Y., and Ron, D. (1999). Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase. *Nature* 397, 271-274.
- Harding, H.P., Zhang, Y., Zeng, H., Novoa, I., Lu, P.D., Calton, M., Sadri, N., Yun, C., Popko, B., Paules, R., *et al.* (2003). An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell* 11, 619-633.
- Harris, V.K., and Sadiq, S.A. (2009). Disease biomarkers in multiple sclerosis: potential for use in therapeutic decision making. *Mol Diagn Ther* 13, 225-244.
- Harrison, J.K., Jiang, Y., Chen, S., Xia, Y., Maciejewski, D., McNamara, R.K., Streit, W.J., Salafranca, M.N., Adhikari, S., Thompson, D.A., *et al.* (1998). Role for neuronally derived fractalkine in mediating interactions between neurons and CX3CR1-expressing microglia. *Proc Natl Acad Sci USA* 95, 10896-10901.



Haughey, N.J., Cutler, R.G., Tamara, A., McArthur, J.C., Vargas, D.L., Pardo, C.A., Turchan, J., Nath, A., and Mattson, M.P. (2004). Perturbation of sphingolipid metabolism and ceramide production in HIV-dementia. *Ann Neurol* 55, 257-267.

He, B., Gross, M., and Roizman, B. (1997). The gamma(1)34.5 protein of herpes simplex virus 1 complexes with protein phosphatase 1alpha to dephosphorylate the alpha subunit of the eukaryotic translation initiation factor 2 and preclude the shutoff of protein synthesis by double-stranded RNA-activated protein kinase. *Proc Natl Acad Sci USA* 94, 843-848.

Heinicke, L.A., Wong, C.J., Lary, J., Nallagatla, S.R., Diegelman-Parente, A., Zheng, X., Cole, J.L., and Bevilacqua, P.C. (2009). RNA dimerization promotes PKR dimerization and activation. *J Mol Biol* 390, 319-338.

Henkel, T., Zabel, U., van Zee, K., Muller, J.M., Fanning, E., and Baeuerle, P.A. (1992). Intramolecular masking of the nuclear location signal and dimerization domain in the precursor for the p50 NF-kappa B subunit. *Cell* 68, 1121-1133.

Herber, D.L., Maloney, J.L., Roth, L.M., Freeman, M.J., Morgan, D., and Gordon, M.N. (2006). Diverse microglial responses after intrahippocampal administration of lipopolysaccharide. *Glia* 53, 382-391.

Hesseltger, J., Taub, D., Baskar, P., Greenberg, M., Hoxie, J., Kolson, D.L., and Horuk, R. (1998). Neuronal apoptosis induced by HIV-1 gp120 and the chemokine SDF-1 alpha is mediated by the chemokine receptor CXCR4. *Curr Biol* 8, 595-598.

Heyes, M.P., Brew, B.J., Martin, A., Price, R.W., Salazar, A.M., Sidtis, J.J., Yergey, J.A., Mouradian, M.M., Sadler, A.E., Keilp, J., *et al.* (1991). Quinolinic acid in cerebrospinal fluid and serum in HIV-1 infection: relationship to clinical and neurological status. *Ann Neurol* 29, 202-209.

Heyes, M.P., Ellis, R.J., Ryan, L., Childers, M.E., Grant, I., Wolfson, T., Archibald, S., and Jernigan, T.L. (2001). Elevated cerebrospinal fluid quinolinic acid levels are associated with region-specific cerebral volume loss in HIV infection. *Brain* 124, 1033-1042.

Ho, D.D., Rota, T.R., Schooley, R.T., Kaplan, J.C., Allan, J.D., Groopman, J.E., Resnick, L., Felsenstein, D., Andrews, C.A., and Hirsch, M.S. (1985). Isolation of HTLV-III from cerebrospinal fluid and neural tissues of patients with neurologic syndromes related to the acquired immunodeficiency syndrome. *N Engl J Med* 313, 1493-1497.

Hoberg, J.E., Popko, A.E., Ramsey, C.S., and Mayo, M.W. (2006). Ikb Kinase  $\alpha$ -Mediated Derepression of SMRT Potentiates Acetylation of RelA/p65 by p300. *Mol Cell Biol* 26, 457-471.

Hoffman, P.M., Dhib-Jalbut, S., Mikovits, J.A., Robbins, D.S., Wolf, A.L., Bergey, G.K., Lohrey, N.C., Weislow, O.S., and Ruscetti, F.W. (1992). Human T-cell leukemia virus type I infection of monocytes and microglial cells in primary human cultures. *Proc Natl Acad Sci USA* 89, 11784-11788.

Holcomb, T., Taylor, L., Trohkimainen, J., and Curthoys, N.P. (2000). Isolation, characterization and expression of a human brain mitochondrial glutaminase cDNA. *Brain Res Mol Brain Res* 76, 56-63.

Holtz, W.A., Turetzky, J.M., Jong, Y.J., and O'Malley, K.L. (2006). Oxidative stress-triggered unfolded protein response is upstream of intrinsic cell death evoked by parkinsonian mimetics. *J Neurochem* 99, 54-69.

Hosokawa, N., Wada, I., Hasegawa, K., Yorihuzi, T., Tremblay, L.O., Herscovics, A., and Nagata, K. (2001). A novel ER alpha-mannosidase-like protein accelerates ER-associated degradation. *Eur Mol Biol Org Rep* 2, 415-422.

Hoxtermann, S., Nuchel, C., and Altmeyer, P. (1998). Fumaric acid esters suppress peripheral CD4- and CD8-positive lymphocytes in psoriasis. *Dermatology* 196, 223-230.

Hunot, S., Dugas, N., Faucheux, B., Hartmann, A., Tardieu, M., Debre, P., Agid, Y., Dugas, B., and Hirsch, E.C. (1999). FcepsilonRII/CD23 is expressed in Parkinson's disease and induces, in vitro, production of nitric oxide and tumor necrosis factor-alpha in glial cells. *J Neurosci* 19, 3440-3447.

Hunt, S.L., and Jackson, R.J. (1999). Polypyrimidine-tract binding protein (PTB) is necessary, but not sufficient, for efficient internal initiation of translation of human rhinovirus-2 RNA. *Rna* 5, 344-359.

Hurwitz, A.A., Berman, J.W., and Lyman, W.D. (1994). The role of the blood-brain barrier in HIV infection of the central nervous system. *Adv Neuroimmunol* 4, 249-256.

Imai, T., Hieshima, K., Haskell, C., Baba, M., Nagira, M., Nishimura, M., Kakizaki, M., Takagi, S., Nomiyama, H., Schall, T.J., *et al.* (1997). Identification and molecular characterization of fractalkine receptor CX3CR1, which mediates both leukocyte migration and adhesion. *Cell* 91, 521-530.

Isler, J.A., Skalet, A.H., and Alwine, J.C. (2005). Human cytomegalovirus infection activates and regulates the unfolded protein response. *J Virol* 79, 6890-6899.

Itoh, K., Wakabayashi, N., Katoh, Y., Ishii, T., Igarashi, K., Engel, J.D., and Yamamoto, M. (1999). Keap1 represses nuclear activation of antioxidant responsive elements by Nrf2 through binding to the amino-terminal Neh2 domain. *Genes Dev* 13, 76-86.

Iwasaki, K., Mackenzie, E.L., Hailemariam, K., Sakamoto, K., and Tsuji, Y. (2006). Hemin-mediated regulation of an antioxidant-responsive element of the human ferritin H gene and role of Ref-1 during erythroid differentiation of K562 cells. *Mol Cell Biol* 26, 2845-2856.

Jacobson, S., Gupta, A., Mattson, D., Mingioli, E., and McFarlin, D.E. (1990a). Immunological studies in tropical spastic paraparesis. *Ann Neurol* 27, 149-156.

Jacobson, S., Shida, H., McFarlin, D.E., Fauci, A.S., and Koenig, S. (1990b). Circulating CD8+ cytotoxic T lymphocytes specific for HTLV-I pX in patients with HTLV-I associated neurological disease. *Nature* 348, 245-248.

Janic, B., Iskander, A.S., Rad, A.M., Soltanian-Zadeh, H., and Arbab, A.S. (2008). Effects of ferumoxides-protamine sulfate labeling on immunomodulatory characteristics of macrophage-like THP-1 cells. *PLoS ONE* 3, e2499.

Jaworowski, A., Kamwendo, D.D., Ellery, P., Sonza, S., Mwapasa, V., Tadesse, E., Molyneux, M.E., Rogerson, S.J., Meshnick, S.R., and Crowe, S.M. (2007). CD16+ monocyte subset preferentially harbors HIV-1 and is expanded in pregnant Malawian women with *Plasmodium falciparum* malaria and HIV-1 infection. *J Infectious Dis* 196, 38-42.

Jiang, H.Y., and Wek, R.C. (2005). GCN2 phosphorylation of eIF2alpha activates NF-kappaB in response to UV irradiation. *Biochem J* 385, 371-380.

- Jiang, Z.G., Piggee, C., Heyes, M.P., Murphy, C., Quearry, B., Bauer, M., Zheng, J., Gendelman, H.E., and Markey, S.P. (2001). Glutamate is a mediator of neurotoxicity in secretions of activated HIV-1-infected macrophages. *J Neuroimmunol* **117**, 97-107.
- Jordan, R., Wang, L., Graczyk, T.M., Block, T.M., and Romano, P.R. (2002). Replication of a cytopathic strain of bovine viral diarrhea virus activates PERK and induces endoplasmic reticulum stress-mediated apoptosis of MDBK cells. *J Virol* **76**, 9588-9599.
- Kanmogne, G.D., Kennedy, R.C., and Grammas, P. (2002). HIV-1 gp120 proteins and gp160 peptides are toxic to brain endothelial cells and neurons: possible pathway for HIV entry into the brain and HIV-associated dementia. *J Neuropathol Exp Neurol* **61**, 992-1000.
- Kasuya, T., Nakajima, H., and Kitamoto, K. (1999). Cloning and characterization of the bipA gene encoding ER chaperone BiP from *Aspergillus oryzae*. *J Biosci Bioeng* **88**, 472-478.
- Kaufman, R.J., Scheuner, D., Schroder, M., Shen, X., Lee, K., Liu, C.Y., and Arnold, S.M. (2002). The unfolded protein response in nutrient sensing and differentiation. *Nat Rev Mol Cell Biol* **3**, 411-421.
- Kaul, M. (2009). HIV-1 associated dementia: update on pathological mechanisms and therapeutic approaches. *Curr Opin Neurol* **22**, 315-320.
- Kaul, M., Garden, G.A., and Lipton, S.A. (2001). Pathways to neuronal injury and apoptosis in HIV-associated dementia. *Nature* **410**, 988-994.
- Kaul, M., and Lipton, S.A. (1999). Chemokines and activated macrophages in HIV gp120-induced neuronal apoptosis. *Proc Natl Acad Sci USA* **96**, 8212-8216.
- Kaul, M., and Lipton, S.A. (2006). Mechanisms of neuroimmunity and neurodegeneration associated with HIV-1 infection and AIDS. *J Neuroimmune Pharmacol* **1**, 138-151.
- Kaul, M., Zheng, J., Okamoto, S., Gendelman, H.E., and Lipton, S.A. (2005). HIV-1 infection and AIDS: consequences for the central nervous system. *Cell Death Different* **12 Suppl 1**, 878-892.
- Kelder, W., McArthur, J.C., Nance-Sproson, T., McClernon, D., and Griffin, D.E. (1998). Beta-chemokines MCP-1 and RANTES are selectively increased in cerebrospinal fluid of patients with human immunodeficiency virus-associated dementia. *Ann Neurol* **44**, 831-835.
- Ketzler, S., Weis, S., Haug, H., and Budka, H. (1990). Loss of neurons in the frontal cortex in AIDS brains. *Acta Neuropathol* **80**, 92-94.
- Khan, M.Z., Brandimarti, R., Patel, J.P., Huynh, N., Wang, J., Huang, Z., Fatatis, A., and Meucci, O. (2004). Apoptotic and antiapoptotic effects of CXCR4: is it a matter of intrinsic efficacy? Implications for HIV neuropathogenesis. *AIDS Res Hum Retroviruses* **20**, 1063-1071.
- Khanlou, N., Moore, D.J., Chana, G., Cherner, M., Lazzaretto, D., Dawes, S., Grant, I., Masliah, E., and Everall, I.P. (2009). Increased frequency of alpha-synuclein in the substantia nigra in human immunodeficiency virus infection. *J Neurovirol* **15**, 131-138.
- Kilareski, E., Shah, S., Nonnemacher, M., and Wigdahl, B. (2009). Regulation of HIV-1 transcription in cells of the monocyte-macrophage lineage. *Retrovirology* **6**, 118.

- Kobayashi S, H.Y., Kobayashi N, Yamamoto N (1990). Serum level of TNF[alpha] in HIV-infected individuals. *AIDS* 4, 169-170.
- Koenig, S., Gendelman, H.E., Orenstein, J.M., Dal Canto, M.C., Pezeshekpour, G.H., Yungbluth, M., Janotta, F., Aksamit, A., Martin, M.A., and Fauci, A.S. (1986). Detection of AIDS virus in macrophages in brain tissue from AIDS patients with encephalopathy. *Science* 233, 1089-1093.
- Kolb, S.A., Sporer, B., Lahrtz, F., Koedel, U., Pfister, H.W., and Fontana, A. (1999). Identification of a T cell chemotactic factor in the cerebrospinal fluid of HIV-1-infected individuals as interferon-gamma inducible protein 10. *J Neuroimmunol* 93, 172-181.
- Kolson, D.L., and Gonzalez-Scarano, F. (2000). HIV-1 and dementia. *J Clin Invest* 106, 11-13.
- Kopito, R.R. (1997). ER quality control: the cytoplasmic connection. *Cell* 88, 427-430.
- Koumenis, C., and Wouters, B.G. (2006). "Translating" tumor hypoxia: unfolded protein response (UPR)-dependent and UPR-independent pathways. *Mol Cancer Res* 4, 423-436.
- Koyanagi, Y., O'Brien, W., Zhao, J., Golde, D., Gasson, J., and Chen, I. (1988). Cytokines alter production of HIV-1 from primary mononuclear phagocytes. *Science* 241, 1673-1675.
- Kraft-Terry, S.D., Buch, S.J., Fox, H.S., and Gendelman, H.E. (2009). A coat of many colors: neuroimmune crosstalk in human immunodeficiency virus infection. *Neuron* 64, 133-145.
- Kramer-Hammerle, S., Rothenaigner, I., Wolff, H., Bell, J.E., and Brack-Werner, R. (2005). Cells of the central nervous system as targets and reservoirs of the human immunodeficiency virus. *Virus Res* 111, 194-213.
- Krishnamoorthy, T., Pavitt, G.D., Zhang, F., Dever, T.E., and Hinnebusch, A.G. (2001). Tight binding of the phosphorylated alpha subunit of initiation factor 2 (eIF2alpha) to the regulatory subunits of guanine nucleotide exchange factor eIF2B is required for inhibition of translation initiation. *Mol Cell Biol* 21, 5018-5030.
- Kruger, R., Hardt, C., Tschentscher, F., Jackel, S., Kuhn, W., Muller, T., Werner, J., Voitalla, D., Berg, D., Kuhn, N., *et al.* (2000). Genetic analysis of immunomodulating factors in sporadic Parkinson's disease. *J Neural Transm* 107, 553-562.
- Kutsch, O., Oh, J., Nath, A., and Benveniste, E.N. (2000). Induction of the chemokines interleukin-8 and IP-10 by human immunodeficiency virus type 1 tat in astrocytes. *J Virol* 74, 9214-9221.
- Lawrence, D.M., and Major, E.O. (2002). HIV-1 and the brain: connections between HIV-1-associated dementia, neuropathology and neuroimmunology. *Microbes Infect* 4, 301-308.
- Lazarini, F., Casanova, P., Tham, T.N., De Clercq, E., Arenzana-Seisdedos, F., Baleux, F., and Dubois-Dalcq, M. (2000). Differential signalling of the chemokine receptor CXCR4 by stromal cell-derived factor 1 and the HIV glycoprotein in rat neurons and astrocytes. *Eur J Neurosci* 12, 117-125.
- Lee, A.H., Iwakoshi, N.N., and Glimcher, L.H. (2003). XBP-1 regulates a subset of endoplasmic reticulum resident chaperone genes in the unfolded protein response. *Mol Cell Biol* 23, 7448-7459.

- Lee, K., Tirasophon, W., Shen, X., Michalak, M., Prywes, R., Okada, T., Yoshida, H., Mori, K., and Kaufman, R.J. (2002). IRE1-mediated unconventional mRNA splicing and S2P-mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein response. *Genes Dev* 16, 452-466.
- Lehmann, J.C., Listopad, J.J., Rentzsch, C.U., Igney, F.H., von Bonin, A., Hennekes, H.H., Asadullah, K., and Docke, W.D. (2007). Dimethylfumarate induces immunosuppression via glutathione depletion and subsequent induction of heme oxygenase 1. *J Invest Dermatol* 127, 835-845.
- Lehmann, M.H., Masanetz, S., Kramer, S., and Erfle, V. (2006). HIV-1 Nef upregulates CCL2/MCP-1 expression in astrocytes in a myristoylation- and calmodulin-dependent manner. *J Cell Sci* 119, 4520-4530.
- Leonard, C.K., Spellman, M.W., Riddle, L., Harris, R.J., Thomas, J.N., and Gregory, T.J. (1990). Assignment of intrachain disulfide bonds and characterization of potential glycosylation sites of the type 1 recombinant human immunodeficiency virus envelope glycoprotein (gp120) expressed in Chinese hamster ovary cells. *J Biol Chem* 265, 10373-10382.
- Letendre, S., Marquie-Beck, J., Capparelli, E., Best, B., Clifford, D., Collier, A.C., Gelman, B.B., McArthur, J.C., McCutchan, J.A., Morgello, S., *et al.* (2008). Validation of the CNS Penetration-Effectiveness rank for quantifying antiretroviral penetration into the central nervous system. *Arch Neurol* 65, 65-70.
- Letendre, S.L., Lanier, E.R., and McCutchan, J.A. (1999). Cerebrospinal fluid beta chemokine concentrations in neurocognitively impaired individuals infected with human immunodeficiency virus type 1. *J Infect Dis* 180, 310-319.
- Letendre, S.L., Marquie-Beck, J., Ellis, R.J., Woods, S.P., Best, B., Clifford, D.B., Collier, A.C., Gelman, B.B., Marra, C., McArthur, J.C., *et al.* (2007). The role of cohort studies in drug development: clinical evidence of antiviral activity of serotonin reuptake inhibitors and HMG-CoA reductase inhibitors in the central nervous system. *J Neuroimmune Pharmacol* 2, 120-127.
- Letendre, S.L., Woods, S.P., Ellis, R.J., Atkinson, J.H., Masliah, E., van den Brande, G., Durelle, J., Grant, I., and Everall, I. (2006). Lithium improves HIV-associated neurocognitive impairment. *AIDS* 20, 1885-1888.
- Levin, M.C., Lehky, T.J., Flierlage, A.N., Katz, D., Kingma, D.W., Jaffe, E.S., Heiss, J.D., Patronas, N., McFarland, H.F., and Jacobson, S. (1997). Immunologic analysis of a spinal cord-biopsy specimen from a patient with human T-cell lymphotropic virus type I-associated neurologic disease. *N Engl J Med* 336, 839-845.
- Li, M., and Ransohoff, R.M. (2008). Multiple roles of chemokine CXCL12 in the central nervous system: a migration from immunology to neurobiology. *Progress Neurobiol* 84, 116-131.
- Li, X.D., Lankinen, H., Putkuri, N., Vapalahti, O., and Vaheri, A. (2005). Tula hantavirus triggers pro-apoptotic signals of ER stress in Vero E6 cells. *Virology* 333, 180-189.
- Lieberman, E., Fong, Y.L., Selby, M.J., Choo, Q.L., Cousens, L., Houghton, M., and Yen, T.S. (1999). Activation of the grp78 and grp94 promoters by hepatitis C virus E2 envelope protein. *J Virol* 73, 3718-3722.

- Limatola, C., Lauro, C., Catalano, M., Ciotti, M.T., Bertollini, C., Di Angelantonio, S., Ragozzino, D., and Eusebi, F. (2005). Chemokine CX3CL1 protects rat hippocampal neurons against glutamate-mediated excitotoxicity. *J Neuroimmunol* 166, 19-28.
- Liner, K.J., 2nd, Hall, C.D., and Robertson, K.R. (2008). Effects of antiretroviral therapy on cognitive impairment. *Curr HIV/AIDS Rep* 5, 64-71.
- Linker, R.A., Lee, D.H., Ryan, S., van Dam, A.M., Conrad, R., Bista, P., Zeng, W., Hronowsky, X., Buko, A., Chollate, S., *et al.* (2011). Fumaric acid esters exert neuroprotective effects in neuroinflammation via activation of the Nrf2 antioxidant pathway. *Brain* 134, 678-692.
- Lipton, S.A. (2004). Failures and successes of NMDA receptor antagonists: molecular basis for the use of open-channel blockers like memantine in the treatment of acute and chronic neurologic insults. *NeuroRx* 1, 101-110.
- Litjens, N.H., Burggraaf, J., van Strijen, E., van Gulpen, C., Mattie, H., Schoemaker, R.C., van Dissel, J.T., Thio, H.B., and Nibbering, P.H. (2004). Pharmacokinetics of oral fumarates in healthy subjects. *Br J Clin Pharmacol* 58, 429-432.
- Liu, S., Suragani, R.N., Wang, F., Han, A., Zhao, W., Andrews, N.C., and Chen, J.J. (2007). The function of heme-regulated eIF2alpha kinase in murine iron homeostasis and macrophage maturation. *J Clin Invest* 117, 3296-3305.
- Loewe, R., Holnthoner, W., Groger, M., Pillinger, M., Gruber, F., Mechtcheriakova, D., Hofer, E., Wolff, K., and Petzelbauer, P. (2002). Dimethylfumarate inhibits TNF-induced nuclear entry of NF-kappa B/p65 in human endothelial cells. *J Immunol* 168, 4781-4787.
- Lovell, M.A., and Markesbery, W.R. (2007). Oxidative damage in mild cognitive impairment and early Alzheimer's disease. *J Neurosci Res* 85, 3036-3040.
- Lu, P.D., Harding, H.P., and Ron, D. (2004). Translation reinitiation at alternative open reading frames regulates gene expression in an integrated stress response. *J Cell Biol* 167, 27-33.
- Ma, Y., Brewer, J.W., Diehl, J.A., and Hendershot, L.M. (2002). Two distinct stress signaling pathways converge upon the CHOP promoter during the mammalian unfolded protein response. *J Mol Biol* 318, 1351-1365.
- Ma, Y., and Hendershot, L.M. (2003). Delineation of a negative feedback regulatory loop that controls protein translation during endoplasmic reticulum stress. *J Biol Chem* 278, 34864-34873.
- Maciejewski-Lenoir, D., Chen, S., Feng, L., Maki, R., and Bacon, K.B. (1999). Characterization of fractalkine in rat brain cells: migratory and activation signals for CX3CR-1-expressing microglia. *J Immunol* 163, 1628-1635.
- Magyar, K., and Szende, B. (2004). (-)-Deprenyl, a selective MAO-B inhibitor, with apoptotic and anti-apoptotic properties. *Neurotoxicology* 25, 233-242.
- Mahad, D.J., Howell, S.J., and Woodroffe, M.N. (2002a). Expression of chemokines in cerebrospinal fluid and serum of patients with chronic inflammatory demyelinating polyneuropathy. *J Neurol Neurosurg Psychiatry* 73, 320-323.

- Mahad, D.J., Howell, S.J., and Woodroffe, M.N. (2002b). Expression of chemokines in the CSF and correlation with clinical disease activity in patients with multiple sclerosis. *J Neurol Neurosurg Psychiatry* 72, 498-502.
- Maitra, R.K., McMillan, N.A., Desai, S., McSwiggen, J., Hovanessian, A.G., Sen, G., Williams, B.R., and Silverman, R.H. (1994). HIV-1 TAR RNA has an intrinsic ability to activate interferon-inducible enzymes. *Virology* 204, 823-827.
- Malik, M., Chen, Y.Y., Kienzle, M.F., Tomkowicz, B.E., Collman, R.G., and Ptasznik, A. (2008). Monocyte migration and LFA-1-mediated attachment to brain microvascular endothelia is regulated by SDF-1 alpha through Lyn kinase. *J Immunol* 181, 4632-4637.
- Malmestrom, C., Andersson, B.A., Haghighi, S., and Lycke, J. (2006). IL-6 and CCL2 levels in CSF are associated with the clinical course of MS: implications for their possible immunopathogenic roles. *J Neuroimmunol* 175, 176-182.
- Mamidi, A., DeSimone, J.A., and Pomerantz, R.J. (2002). Central nervous system infections in individuals with HIV-1 infection. *J Neurovirol* 8, 158-167.
- Maragos, W.F., Tillman, P., Jones, M., Bruce-Keller, A.J., Roth, S., Bell, J.E., and Nath, A. (2003). Neuronal injury in hippocampus with human immunodeficiency virus transactivating protein, Tat. *Neuroscience* 117, 43-53.
- Marra, C.M., Zhao, Y., Clifford, D.B., Letendre, S., Evans, S., Henry, K., Ellis, R.J., Rodriguez, B., Coombs, R.W., Schifitto, G., *et al.* (2009). Impact of combination antiretroviral therapy on cerebrospinal fluid HIV RNA and neurocognitive performance. *AIDS* 23, 1359-1366.
- Martin, M., Serradji, N., Dereuddre-Bosquet, N., Le Pavec, G., Fichet, G., Lamouri, A., Heymans, F., Godfroid, J.J., Clayette, P., and Dormont, D. (2000). PMS-601, a new platelet-activating factor receptor antagonist that inhibits human immunodeficiency virus replication and potentiates zidovudine activity in macrophages. *Antimicrob Agents Chemother* 44, 3150-3154.
- Martin-Garcia, J., Kolson, D.L., and Gonzalez-Scarano, F. (2002). Chemokine receptors in the brain: their role in HIV infection and pathogenesis. *AIDS* 16, 1709-1730.
- Maslah, E., Heaton, R.K., Marcotte, T.D., Ellis, R.J., Wiley, C.A., Mallory, M., Achim, C.L., McCutchan, J.A., Nelson, J.A., Atkinson, J.H., *et al.* (1997). Dendritic injury is a pathological substrate for human immunodeficiency virus-related cognitive disorders. HNRC Group. The HIV Neurobehavioral Research Center. *Ann Neurol* 42, 963-972.
- Matsumoto, R., Akama, K., Rakwal, R., and Iwahashi, H. (2005). The stress response against denatured proteins in the deletion of cytosolic chaperones SSA1/2 is different from heat-shock response in *Saccharomyces cerevisiae*. *BMC Genomics* 6, 141.
- McArthur, J.C., Brew, B.J., and Nath, A. (2005). Neurological complications of HIV infection. *Lancet Neurol* 4, 543-555.
- McArthur, J.C., Haughey, N., Gartner, S., Conant, K., Pardo, C., Nath, A., and Sacktor, N. (2003). Human immunodeficiency virus-associated dementia: an evolving disease. *Journal of Neurovirology* 9, 205-221.

- McArthur, J.C., McClernon, D.R., Cronin, M.F., Nance-Sproson, T.E., Saah, A.J., St Clair, M., and Lanier, E.R. (1997). Relationship between human immunodeficiency virus-associated dementia and viral load in cerebrospinal fluid and brain. *Ann Neurol* 42, 689-698.
- McArthur, J.C., McDermott, M.P., McClernon, D., St Hillaire, C., Conant, K., Marder, K., Schifitto, G., Selnes, O.A., Sacktor, N., Stern, Y., *et al.* (2004). Attenuated central nervous system infection in advanced HIV/AIDS with combination antiretroviral therapy. *Arch Neurol* 61, 1687-1696.
- McArthur, J.C., Nance-Sproson, T.E., Griffin, D.E., Hoover, D., Selnes, O.A., Miller, E.N., Margolick, J.B., Cohen, B.A., Farzadegan, H., and Saah, A. (1992). The diagnostic utility of elevation in cerebrospinal fluid beta 2-microglobulin in HIV-1 dementia. Multicenter AIDS Cohort Study. *Neurology* 42, 1707-1712.
- McArthur, J.C., Steiner, J., Sacktor, N., and Nath, A. (2010). Human immunodeficiency virus-associated neurocognitive disorders: Mind the gap. *Ann Neurol* 67, 699-714.
- McCombe, J.A., Auer, R.N., Maingat, F.G., Houston, S., Gill, M.J., and Power, C. (2009). Neurologic immune reconstitution inflammatory syndrome in HIV/AIDS: outcome and epidemiology. *Neurology* 72, 835-841.
- McCrossan, M., Marsden, M., Carnie, F.W., Minnis, S., Hansoti, B., Anthony, I.C., Brett, R.P., Bell, J.E., and Simmonds, P. (2006). An immune control model for viral replication in the CNS during presymptomatic HIV infection. *Brain* 129, 503-516.
- McCullough, K.D., Martindale, J.L., Klotz, L.O., Aw, T.Y., and Holbrook, N.J. (2001). Gadd153 sensitizes cells to endoplasmic reticulum stress by down-regulating Bcl2 and perturbing the cellular redox state. *Mol Cell Biol* 21, 1249-1259.
- McFarlin, D.E., and Blattner, W.A. (1991). Non-AIDS retroviral infections in humans. *Annu Rev Med* 42, 97-105.
- McGeer, E.G., and McGeer, P.L. (2010). Neuroinflammation in Alzheimer's disease and mild cognitive impairment: a field in its infancy. *J Alzheimers Dis* 19, 355-361.
- McGeer, P.L., Itagaki, S., Tago, H., and McGeer, E.G. (1987). Reactive microglia in patients with senile dementia of the Alzheimer type are positive for the histocompatibility glycoprotein HLA-DR. *Neurosci Lett* 79, 195-200.
- McManus, C.M., Liu, J.S., Hahn, M.T., Hua, L.L., Brosnan, C.F., Berman, J.W., and Lee, S.C. (2000). Differential induction of chemokines in human microglia by type I and II interferons. *Glia* 29, 273-280.
- Meisner, F., Scheller, C., Kneitz, S., Sopper, S., Neuen-Jacob, E., Riederer, P., ter Meulen, V., and Koutsilieri, E. (2008). Memantine upregulates BDNF and prevents dopamine deficits in SIV-infected macaques: a novel pharmacological action of memantine. *Neuropsychopharmacology* 33, 2228-2236.
- Mengozzi, M., De Filippi, C., Transidico, P., Biswas, P., Cota, M., Ghezzi, S., Vicenzi, E., Mantovani, A., Sozzani, S., and Poli, G. (1999). Human immunodeficiency virus replication induces monocyte chemotactic protein-1 in human macrophages and U937 promonocytic cells. *Blood* 93, 1851-1857.



- Meucci, O., Fatatis, A., Simen, A.A., Bushell, T.J., Gray, P.W., and Miller, R.J. (1998). Chemokines regulate hippocampal neuronal signaling and gp120 neurotoxicity. *Proc Natl Acad Sci USA* 95, 14500-14505.
- Meucci, O., Fatatis, A., Simen, A.A., and Miller, R.J. (2000). Expression of CX3CR1 chemokine receptors on neurons and their role in neuronal survival. *Proc Natl Acad Sci USA* 97, 8075-8080.
- Miljkovic, D., Drulovic, J., Trajkovic, V., Mesaros, S., Dujmovic, I., Maksimovic, D., Samardzic, T., Stojasavljevic, N., Levic, Z., and Mostarica Stojkovic, M. (2002). Nitric oxide metabolites and interleukin-6 in cerebrospinal fluid from multiple sclerosis patients. *Eur J Neurol* 9, 413-418.
- Mizuno, T., Kawanokuchi, J., Numata, K., and Suzumura, A. (2003). Production and neuroprotective functions of fractalkine in the central nervous system. *Brain Res* 979, 65-70.
- Mordechai, E., Kon, N., Henderson, E.E., and Suhadolnik, R.J. (1995). Activation of the interferon-inducible enzymes, 2',5'-oligoadenylate synthetase and PKR by human T-cell leukemia virus type I Rex-response element. *Virology* 206, 913-922.
- Mori, K. (2000). Tripartite management of unfolded proteins in the endoplasmic reticulum. *Cell* 101, 451-454.
- Mrak, R.E., and Griffin, W.S. (2005). Glia and their cytokines in progression of neurodegeneration. *Neurobiol Aging* 26, 349-354.
- Mulvey, M., Arias, C., and Mohr, I. (2007). Maintenance of endoplasmic reticulum (ER) homeostasis in herpes simplex virus type 1-infected cells through the association of a viral glycoprotein with PERK, a cellular ER stress sensor. *J Virol* 81, 3377-3390.
- Nabel, G., and Baltimore, D. (1987). An inducible transcription factor activates expression of human immunodeficiency virus in T cells. *Nature* 326, 711-713.
- Nagai, M., Brennan, M.B., Sakai, J.A., Mora, C.A., and Jacobson, S. (2001a). CD8(+) T cells are an in vivo reservoir for human T-cell lymphotropic virus type I. *Blood* 98, 1858-1861.
- Nagai, M., Yamano, Y., Brennan, M.B., Mora, C.A., and Jacobson, S. (2001b). Increased HTLV-I proviral load and preferential expansion of HTLV-I Tax-specific CD8+ T cells in cerebrospinal fluid from patients with HAM/TSP. *Ann Neurol* 50, 807-812.
- Nakagawa, T., Zhu, H., Morishima, N., Li, E., Xu, J., Yankner, B.A., and Yuan, J. (2000). Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* 403, 98-103.
- Nath, A., Haughey, N.J., Jones, M., Anderson, C., Bell, J.E., and Geiger, J.D. (2000). Synergistic neurotoxicity by human immunodeficiency virus proteins Tat and gp120: protection by memantine. *Ann Neurol* 47, 186-194.
- Nath, A., Schiess, N., Venkatesan, A., Rumbaugh, J., Sacktor, N., and McArthur, J. (2008). Evolution of HIV dementia with HIV infection. *Int Rev Psychiatry* 20, 25-31.
- Neumann, M., Fries, H.-W., Scheicher, C., Keikavoussi, P., Kolb-Mäurer, A., Bröcker, E.-B., Serfling, E., and Kämpgen, E. (2000). Differential expression of Rel/NF- $\kappa$ B and octamer factors is a hallmark of the generation and maturation of dendritic cells. *Blood* 95, 277-285.

Nguyen, D.G., Booth, A., Gould, S.J., and Hildreth, J.E. (2003). Evidence that HIV budding in primary macrophages occurs through the exosome release pathway. *J Biol Chem* 278, 52347-52354.

Nicklas, W.J., Zeevalk, G., and Hyndman, A. (1987). Interactions between neurons and glia in glutamate/glutamine compartmentation. *Biochem Soc Trans* 15, 208-210.

Nishitoh, H., Matsuzawa, A., Tobiume, K., Saegusa, K., Takeda, K., Inoue, K., Hori, S., Kakizuka, A., and Ichijo, H. (2002). ASK1 is essential for endoplasmic reticulum stress-induced neuronal cell death triggered by expanded polyglutamine repeats. *Genes Dev* 16, 1345-1355.

Nottet, H.S., Persidsky, Y., Sasseville, V.G., Nukuna, A.N., Bock, P., Zhai, Q.H., Sharer, L.R., McComb, R.D., Swindells, S., Soderland, C., *et al.* (1996). Mechanisms for the transendothelial migration of HIV-1-infected monocytes into brain. *J Immunol* 156, 1284-1295.

O'Donnell, L.A., Agrawal, A., Jordan-Sciutto, K.L., Dichter, M.A., Lynch, D.R., and Kolson, D.L. (2006). Human immunodeficiency virus (HIV)-induced neurotoxicity: roles for the NMDA receptor subtypes. *J Neurosci* 26, 981-990.

Oda, Y., Hosokawa, N., Wada, I., and Nagata, K. (2003). EDEM as an acceptor of terminally misfolded glycoproteins released from calnexin. *Science* 299, 1394-1397.

Ohtani, Y., Minami, M., Kawaguchi, N., Nishiyori, A., Yamamoto, J., Takami, S., and Satoh, M. (1998). Expression of stromal cell-derived factor-1 and CXCR4 chemokine receptor mRNAs in cultured rat glial and neuronal cells. *Neurosci Lett* 249, 163-166.

Ollenschlager, G., Jansen, S., Schindler, J., Rasokat, H., Schrappe-Bacher, M., and Roth, E. (1988). Plasma amino acid pattern of patients with HIV infection. *Clin Chem* 34, 1787-1789.

Osame, M., Janssen, R., Kubota, H., Nishitani, H., Igata, A., Nagataki, S., Mori, M., Goto, I., Shimabukuro, H., Khabbaz, R., *et al.* (1990). Nationwide survey of HTLV-I-associated myelopathy in Japan: association with blood transfusion. *Ann Neurol* 28, 50-56.

Oster, W., Lindemann, A., Horn, S., Mertelsmann, R., and Herrmann, F. (1987). Tumor necrosis factor (TNF)-alpha but not TNF-beta induces secretion of colony stimulating factor for macrophages (CSF-1) by human monocytes. *Blood* 70, 1700-1703.

Ownby, R., Kumar, A., Benny Fernandez, J., Moleon-Borodowsky, I., Gonzalez, L., Eisdorfer, S., Waldrop-Valverde, D., and Kumar, M. (2009). Tumor Necrosis Factor-alpha Levels in HIV-1 Seropositive Injecting Drug Users. *J Neuroimmune Pharmacol* 4, 350-358.

Oyadomari, S., Araki, E., and Mori, M. (2002a). Endoplasmic reticulum stress-mediated apoptosis in pancreatic beta-cells. *Apoptosis* 7, 335-345.

Oyadomari, S., Koizumi, A., Takeda, K., Gotoh, T., Akira, S., Araki, E., and Mori, M. (2002b). Targeted disruption of the Chop gene delays endoplasmic reticulum stress-mediated diabetes. *J Clin Invest* 109, 525-532.

Park, I.W., Wang, J.F., and Groopman, J.E. (2001). HIV-1 Tat promotes monocyte chemoattractant protein-1 secretion followed by transmigration of monocytes. *Blood* 97, 352-358.

Pelchen-Matthews, A., Kramer, B., and Marsh, M. (2003). Infectious HIV-1 assembles in late endosomes in primary macrophages. *J Cell Biol* 162, 443-455.

Peled, A., Grabovsky, V., Habler, L., Sandbank, J., Arenzana-Seisdedos, F., Petit, I., Ben-Hur, H., Lapidot, T., and Alon, R. (1999). The chemokine SDF-1 stimulates integrin-mediated arrest of CD34(+) cells on vascular endothelium under shear flow. *J Clin Invest* 104, 1199-1211.

Pelletier, J., and Sonenberg, N. (1988). Internal initiation of translation of eukaryotic mRNA directed by a sequence derived from poliovirus RNA. *Nature* 334, 320-325.

Peng, H., Erdmann, N., Whitney, N., Dou, H., Gorantla, S., Gendelman, H.E., Ghorpade, A., and Zheng, J. (2006). HIV-1-infected and/or immune activated macrophages regulate astrocyte SDF-1 production through IL-1beta. *Glia* 54, 619-629.

Pereira, C.F., Middel, J., Jansen, G., Verhoef, J., and Nottet, H.S. (2001). Enhanced expression of fractalkine in HIV-1 associated dementia. *J Neuroimmunol* 115, 168-175.

Perrella, O., Guerriero, M., Izzo, E., Soscia, M., and Carrieri, P.B. (1992). Interleukin-6 and granulocyte macrophage-CSF in the cerebrospinal fluid from HIV infected subjects with involvement of the central nervous system. *Arq Neuropsiquiatr* 50, 180-182.

Persidsky, Y., Heilman, D., Haorah, J., Zelivyanskaya, M., Persidsky, R., Weber, G.A., Shimokawa, H., Kaibuchi, K., and Ikezu, T. (2006). Rho-mediated regulation of tight junctions during monocyte migration across the blood-brain barrier in HIV-1 encephalitis (HIVE). *Blood* 107, 4770-4780.

Persidsky, Y., Stins, M., Way, D., Witte, M.H., Weinand, M., Kim, K.S., Bock, P., Gendelman, H.E., and Fiala, M. (1997). A model for monocyte migration through the blood-brain barrier during HIV-1 encephalitis. *J Immunol* 158, 3499-3510.

Persidsky, Y., Zheng, J., Miller, D., and Gendelman, H.E. (2000). Mononuclear phagocytes mediate blood-brain barrier compromise and neuronal injury during HIV-1-associated dementia. *J Leukoc Biol* 68, 413-422.

Petito, C.K., Cho, E.S., Lemann, W., Navia, B.A., and Price, R.W. (1986). Neuropathology of acquired immunodeficiency syndrome (AIDS): an autopsy review. *J Neuropathol Exp Neurol* 45, 635-646.

Petito, C.K., and Roberts, B. (1995). Evidence of apoptotic cell death in HIV encephalitis. *Am J Pathol* 146, 1121-1130.

Poli, G., Kinter, A., Justement, J.S., Kehrl, J.H., Bressler, P., Stanley, S., and Fauci, A.S. (1990). Tumor necrosis factor alpha functions in an autocrine manner in the induction of human immunodeficiency virus expression. *Proc Natl Acad Sci USA* 87, 782-785.

Porcheray, F., Leone, C., Samah, B., Rimaniol, A.C., Dereuddre-Bosquet, N., and Gras, G. (2006). Glutamate metabolism in HIV-infected macrophages: implications for the CNS. *Am J Physiol Cell Physiol* 291, C618-626.

Portis, J.L., Askovich, P., Austin, J., Gutierrez-Cotto, Y., and McAtee, F.J. (2009). The degree of folding instability of the envelope protein of a neurovirulent murine retrovirus correlates with the severity of the neurological disease. *J Virol* 83, 6079-6086.

Power, C., McArthur, J.C., Nath, A., Wehrly, K., Mayne, M., Nishio, J., Langelier, T., Johnson, R.T., and Chesebro, B. (1998). Neuronal death induced by brain-derived human

immunodeficiency virus type 1 envelope genes differs between demented and nondemented AIDS patients. *J Virol* 72, 9045-9053.

Pulliam, L., Gascon, R., Stubblebine, M., McGuire, D., and McGrath, M.S. (1997). Unique monocyte subset in patients with AIDS dementia. *Lancet* 349, 692-695.

Qiao, X., Cummins, D.J., and Paul, S.M. (2001). Neuroinflammation-induced acceleration of amyloid deposition in the APPV717F transgenic mouse. *Eur J Neurosci* 14, 474-482.

Ragin, A.B., Wu, Y., Storey, P., Cohen, B.A., Edelman, R.R., and Epstein, L.G. (2006). Monocyte chemoattractant protein-1 correlates with subcortical brain injury in HIV infection. *Neurology* 66, 1255-1257.

Ransohoff, R.M. (1999). Mechanisms of inflammation in MS tissue: adhesion molecules and chemokines. *J Neuroimmunol* 98, 57-68.

Ransohoff, R.M., and Zamvil, S.S. (2007). Neuroimmunotherapeutics comes of age. *Neurotherapeutics* 4, 569-570.

Rappaport, J., Joseph, J., Croul, S., Alexander, G., Del Valle, L., Amini, S., and Khalili, K. (1999). Molecular pathway involved in HIV-1-induced CNS pathology: role of viral regulatory protein, Tat. *J Leukoc Biol* 65, 458-465.

Re, D.B., and Przedborski, S. (2006). Fractalkine: moving from chemotaxis to neuroprotection. *Nat Neurosci* 9, 859-861.

Rieckmann, P., Albrecht, M., Kitze, B., Weber, T., Tumani, H., Broocks, A., Luer, W., Helwig, A., and Poser, S. (1995). Tumor necrosis factor-alpha messenger RNA expression in patients with relapsing-remitting multiple sclerosis is associated with disease activity. *Ann Neurol* 37, 82-88.

Robertson, H.D., Manche, L., and Mathews, M.B. (1996). Paradoxical interactions between human delta hepatitis agent RNA and the cellular protein kinase PKR. *J Virol* 70, 5611-5617.

Robertson, K.R., Smurzynski, M., Parsons, T.D., Wu, K., Bosch, R.J., Wu, J., McArthur, J.C., Collier, A.C., Evans, S.R., and Ellis, R.J. (2007). The prevalence and incidence of neurocognitive impairment in the HAART era. *AIDS* 21, 1915-1921.

Roederer, M., Staal, F.J., Raju, P.A., Ela, S.W., and Herzenberg, L.A. (1990). Cytokine-stimulated human immunodeficiency virus replication is inhibited by N-acetyl-L-cysteine. *Proc Natl Acad Sci USA* 87, 4884-4888.

Romisch, K. (2004). A cure for traffic jams: small molecule chaperones in the endoplasmic reticulum. *Traffic* 5, 815-820.

Ron, D., and Walter, P. (2007). Signal integration in the endoplasmic reticulum unfolded protein response. *Nat Rev Mol Cell Biol* 8, 519-529.

Rosati, G. (2001). The prevalence of multiple sclerosis in the world: an update. *Neurol Sci* 22, 117-139.

Rostasy, K., Egles, C., Chauhan, A., Kneissl, M., Bahrani, P., Yiannoutsos, C., Hunter, D.D., Nath, A., Hedreen, J.C., and Navia, B.A. (2003). SDF-1alpha is expressed in astrocytes and

neurons in the AIDS dementia complex: an in vivo and in vitro study. *J Neuropathol Exp Neurol* 62, 617-626.

Roulet, E. (1999). Opportunistic infections of the central nervous system during HIV-1 infection (emphasis on cytomegalovirus disease). *J Neurol* 246, 237-243.

Rowell, J.F., Stanhope, P.E., and Siliciano, R.F. (1995). Endocytosis of endogenously synthesized HIV-1 envelope protein. Mechanism and role in processing for association with class II MHC. *J Immunol* 155, 473-488.

Roy, S., Agy, M., Hovanessian, A.G., Sonenberg, N., and Katze, M.G. (1991). The integrity of the stem structure of human immunodeficiency virus type 1 Tat-responsive sequence of RNA is required for interaction with the interferon-induced 68,000-Mr protein kinase. *J Virol* 65, 632-640.

Sabri, F., Titanji, K., De Mito, A., and Chiodi, F. (2003). Astrocyte activation and apoptosis: their roles in the neuropathology of HIV infection. *Brain Pathol* 13, 84-94.

Saccani, A., Saccani, S., Orlando, S., Sironi, M., Bernasconi, S., Ghezzi, P., Mantovani, A., and Sica, A. (2000). Redox regulation of chemokine receptor expression. *Proc Natl Acad Sci USA* 97, 2761-2766.

Sacktor, N., McDermott, M.P., Marder, K., Schifitto, G., Selnes, O.A., McArthur, J.C., Stern, Y., Albert, S., Palumbo, D., Kieburtz, K., *et al.* (2002). HIV-associated cognitive impairment before and after the advent of combination therapy. *J Neurovirol* 8, 136-142.

Sacktor, N., Schifitto, G., McDermott, M.P., Marder, K., McArthur, J.C., and Kieburtz, K. (2000). Transdermal selegiline in HIV-associated cognitive impairment: pilot, placebo-controlled study. *Neurology* 54, 233-235.

Sadagopal, S., Lorey, S.L., Barnett, L., Basham, R., Lebo, L., Erdem, H., Haman, K., Avison, M., Waddell, K., Haas, D.W., *et al.* (2008). Enhancement of human immunodeficiency virus (HIV)-specific CD8+ T cells in cerebrospinal fluid compared to those in blood among antiretroviral therapy-naive HIV-positive subjects. *J Virol* 82, 10418-10428.

Sardar, A.M., Bell, J.E., and Reynolds, G.P. (1995). Increased concentrations of the neurotoxin 3-hydroxykynurenine in the frontal cortex of HIV-1-positive patients. *J Neurochem* 64, 932-935.

Sasseville, V.G., Newman, W., Brodie, S.J., Hesterberg, P., Pauley, D., and Ringler, D.J. (1994). Monocyte adhesion to endothelium in simian immunodeficiency virus-induced AIDS encephalitis is mediated by vascular cell adhesion molecule-1/alpha 4 beta 1 integrin interactions. *Am J Pathol* 144, 27-40.

Savarin-Vuillat, C., and Ransohoff, R.M. (2007). Chemokines and chemokine receptors in neurological disease: raise, retain, or reduce? *Neurotherapeutics* 4, 590-601.

Sawada, M., Imamura, K., and Nagatsu, T. (2006). Role of cytokines in inflammatory process in Parkinson's disease. *J Neural Transm Suppl*, 373-381.

Schaedler, S., Krause, J., Himmelsbach, K., Carvajal-Yepes, M., Lieder, F., Klingel, K., Nassal, M., Weiss, T.S., Werner, S., and Hildt, E. (2010). Hepatitis B virus induces expression of antioxidant response element-regulated genes by activation of Nrf2. *J Biol Chem* 285, 41074-41086.

Schenk, S., Saberi, M., and Olefsky, J.M. (2008). Insulin sensitivity: modulation by nutrients and inflammation. *J Clin Invest* 118, 2992-3002.

Schifitto, G., Navia, B.A., Yiannoutsos, C.T., Marra, C.M., Chang, L., Ernst, T., Jarvik, J.G., Miller, E.N., Singer, E.J., Ellis, R.J., *et al.* (2007a). Memantine and HIV-associated cognitive impairment: a neuropsychological and proton magnetic resonance spectroscopy study. *AIDS* 21, 1877-1886.

Schifitto, G., Peterson, D.R., Zhong, J., Ni, H., Cruttenden, K., Gaugh, M., Gendelman, H.E., Boska, M., and Gelbard, H. (2006). Valproic acid adjunctive therapy for HIV-associated cognitive impairment: a first report. *Neurology* 66, 919-921.

Schifitto, G., Yiannoutsos, C.T., Ernst, T., Navia, B.A., Nath, A., Sacktor, N., Anderson, C., Marra, C.M., and Clifford, D.B. (2009a). Selegiline and oxidative stress in HIV-associated cognitive impairment. *Neurology* 73, 1975-1981.

Schifitto, G., Zhang, J., Evans, S.R., Sacktor, N., Simpson, D., Millar, L.L., Hung, V.L., Miller, E.N., Smith, E., Ellis, R.J., *et al.* (2007b). A multicenter trial of selegiline transdermal system for HIV-associated cognitive impairment. *Neurology* 69, 1314-1321.

Schifitto, G., Zhong, J., Gill, D., Peterson, D.R., Gaugh, M.D., Zhu, T., Tivarus, M., Cruttenden, K., Maggirwar, S.B., Gendelman, H.E., *et al.* (2009b). Lithium therapy for human immunodeficiency virus type 1-associated neurocognitive impairment. *J Neurovirol* 15, 176-186.

Schilling, S., Goelz, S., Linker, R., Luehder, F., and Gold, R. (2006). Fumaric acid esters are effective in chronic experimental autoimmune encephalomyelitis and suppress macrophage infiltration. *Clin Exp Immunol* 145, 101-107.

Schreck R, R.P., Baeuerle PA (1991). Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. *Eur Mol Biol Org* 10, 2247-2258.

Schubert, U., Anton, L.C., Gibbs, J., Norbury, C.C., Yewdell, J.W., and Bennink, J.R. (2000). Rapid degradation of a large fraction of newly synthesized proteins by proteasomes. *Nature* 404, 770-774.

Schweinsburg, B.C., Taylor, M.J., Alhassoon, O.M., Gonzalez, R., Brown, G.G., Ellis, R.J., Letendre, S., Videen, J.S., McCutchan, J.A., Patterson, T.L., *et al.* (2005). Brain mitochondrial injury in human immunodeficiency virus-seropositive (HIV+) individuals taking nucleoside reverse transcriptase inhibitors. *J Neurovirol* 11, 356-364.

Scorziello, A., Florio, T., Bajetto, A., and Schettini, G. (1998). Intracellular signalling mediating HIV-1 gp120 neurotoxicity. *Cell Signal* 10, 75-84.

Seidel, P., Merfort, I., Hughes, J.M., Oliver, B.G., Tamm, M., and Roth, M. (2009). Dimethylfumarate inhibits NF- $\kappa$ B function at multiple levels to limit airway smooth muscle cell cytokine secretion. *Am J Physiol Lung Cell Mol Physiol* 297, L326-339.

Sevigny, J.J., Albert, S.M., McDermott, M.P., McArthur, J.C., Sacktor, N., Conant, K., Schifitto, G., Selnes, O.A., Stern, Y., McClernon, D.R., *et al.* (2004). Evaluation of HIV RNA and markers of immune activation as predictors of HIV-associated dementia. *Neurology* 63, 2084-2090.

Sevigny, J.J., Albert, S.M., McDermott, M.P., Schifitto, G., McArthur, J.C., Sacktor, N., Conant, K., Selnes, O.A., Stern, Y., McClernon, D.R., *et al.* (2007). An evaluation of neurocognitive status

and markers of immune activation as predictors of time to death in advanced HIV infection. *Arch Neurol* 64, 97-102.

Shen, J., Chen, X., Hendershot, L., and Prywes, R. (2002). ER stress regulation of ATF6 localization by dissociation of BiP/GRP78 binding and unmasking of Golgi localization signals. *Dev Cell* 3, 99-111.

Sheng, J.G., Griffin, W.S., Royston, M.C., and Mrak, R.E. (1998). Distribution of interleukin-1-immunoreactive microglia in cerebral cortical layers: implications for neuritic plaque formation in Alzheimer's disease. *Neuropathol Appl Neurobiol* 24, 278-283.

Shiramizu, B., Gartner, S., Williams, A., Shikuma, C., Ratto-Kim, S., Watters, M., Aguon, J., and Valcour, V. (2005). Circulating proviral HIV DNA and HIV-associated dementia. *AIDS* 19, 45-52.

Si, Q., Cosenza, M., Kim, M.O., Zhao, M.L., Brownlee, M., Goldstein, H., and Lee, S. (2004). A novel action of minocycline: inhibition of human immunodeficiency virus type 1 infection in microglia. *J Neurovirol* 10, 284-292.

Si, Q., Kim, M.O., Zhao, M.L., Landau, N.R., Goldstein, H., and Lee, S. (2002). Vpr- and Nef-dependent induction of RANTES/CCL5 in microglial cells. *Virology* 301, 342-353.

Soldan, S.S., Wu, G., Markowitz, C., and Kolson, D.L. (2008). Multiple Sclerosis and Other Demyelinating Diseases. In *Neuroimmune Pharmacology*, T. Ikezu, and H.E. Gendelman, eds. (Springer).

Song, L., Nath, A., Geiger, J.D., Moore, A., and Hochman, S. (2003). Human immunodeficiency virus type 1 Tat protein directly activates neuronal N-methyl-D-aspartate receptors at an allosteric zinc-sensitive site. *J Neurovirol* 9, 399-403.

Sood, R., Porter, A.C., Olsen, D.A., Cavener, D.R., and Wek, R.C. (2000). A mammalian homologue of GCN2 protein kinase important for translational control by phosphorylation of eukaryotic initiation factor-2alpha. *Genetics* 154, 787-801.

Sozzani, S., Introna, M., Bernasconi, S., Polentarutti, N., Cinque, P., Poli, G., Sica, A., and Mantovani, A. (1997). MCP-1 and CCR2 in HIV infection: regulation of agonist and receptor expression. *J Leukoc Biol* 62, 30-33.

Spitzenberger, T.J., Heilman, D., Diekmann, C., Batrakova, E.V., Kabanov, A.V., Gendelman, H.E., Elmquist, W.F., and Persidsky, Y. (2007). Novel delivery system enhances efficacy of antiretroviral therapy in animal model for HIV-1 encephalitis. *J Cereb Blood Flow Metab* 27, 1033-1042.

Sporer, B., Kastenbauer, S., Koedel, U., Arendt, G., and Pfister, H.W. (2003). Increased intrathecal release of soluble fractalkine in HIV-infected patients. *AIDS Res Hum Retroviruses* 19, 111-116.

Staal FJ, R.M., Raju PA, Anderson MT, Ela SW, Herzenberg LA, Herzenberg LA (1993). Antioxidants inhibit stimulation of HIV transcription. *AIDS Res Hum Retroviruses* 9, 299-306.

Stoof, T.J., Flier, J., Sampat, S., Nieboer, C., Tensen, C.P., and Boorsma, D.M. (2001). The antipsoriatic drug dimethylfumarate strongly suppresses chemokine production in human keratinocytes and peripheral blood mononuclear cells. *Br J Dermatol* 144, 1114-1120.

- Strachan, G.D., Kopp, A.S., Koike, M.A., Morgan, K.L., and Jordan-Sciutto, K.L. (2005). Chemokine- and neurotrophic factor-induced changes in E2F1 localization and phosphorylation of the retinoblastoma susceptibility gene product (pRb) occur by distinct mechanisms in murine cortical cultures. *Exp Neurol* 193, 455-468.
- Su, H.L., Liao, C.L., and Lin, Y.L. (2002). Japanese encephalitis virus infection initiates endoplasmic reticulum stress and an unfolded protein response. *J Virol* 76, 4162-4171.
- Sui, Y., Potula, R., Dhillon, N., Pinson, D., Li, S., Nath, A., Anderson, C., Turchan, J., Kolson, D., Narayan, O., *et al.* (2004). Neuronal apoptosis is mediated by CXCL10 overexpression in simian human immunodeficiency virus encephalitis. *Am J Pathol* 164, 1557-1566.
- Sui, Y., Stehno-Bittel, L., Li, S., Loganathan, R., Dhillon, N.K., Pinson, D., Nath, A., Kolson, D., Narayan, O., and Buch, S. (2006a). CXCL10-induced cell death in neurons: role of calcium dysregulation. *Eur J Neurosci* 23, 957-964.
- Sui, Z., Fan, S., Sniderhan, L., Reisinger, E., Litzburg, A., Schifitto, G., Gelbard, H.A., Dewhurst, S., and Maggirwar, S.B. (2006b). Inhibition of mixed lineage kinase 3 prevents HIV-1 Tat-mediated neurotoxicity and monocyte activation. *J Immunol* 177, 702-711.
- Sun, Z., Arendt, C.W., Ellmeier, W., Schaeffer, E.M., Sunshine, M.J., Gandhi, L., Annes, J., Petrzilka, D., Kupfer, A., Schwartzberg, P.L., *et al.* (2000). PKC-[thetas] is required for TCR-induced NF-[kappa]B activation in mature but not immature T lymphocytes. *Nature* 404, 402-407.
- Sundar, S.K., Cierpial, M.A., Kamaraju, L.S., Long, S., Hsieh, S., Lorenz, C., Aaron, M., Ritchie, J.C., and Weiss, J.M. (1991). Human immunodeficiency virus glycoprotein (gp120) infused into rat brain induces interleukin 1 to elevate pituitary-adrenal activity and decrease peripheral cellular immune responses. *Proc Natl Acad Sci USA* 88, 11246-11250.
- Swindells, S., Zheng, J., and Gendelman, H.E. (1999). HIV-associated dementia: new insights into disease pathogenesis and therapeutic interventions. *AIDS Patient Care STDS* 13, 153-163.
- Szeto, G.L., Brice, A.K., Yang, H.C., Barber, S.A., Siliciano, R.F., and Clements, J.E. (2010). Minocycline attenuates HIV infection and reactivation by suppressing cellular activation in human CD4+ T cells. *J Infect Dis* 201, 1132-1140.
- Szymocha, R., Akaoka, H., Brisson, C., Beurton-Marduel, P., Chalon, A., Bernard, A., Didier-Bazes, M., Belin, M.F., and Giraudon, P. (2000a). Astrocytic alterations induced by HTLV type 1-infected T lymphocytes: a role for Tax-1 and tumor necrosis factor alpha. *AIDS Res Hum Retroviruses* 16, 1723-1729.
- Szymocha, R., Brisson, C., Bernard, A., Akaoka, H., Belin, M.F., and Giraudon, P. (2000b). Long-term effects of HTLV-1 on brain astrocytes: sustained expression of Tax-1 associated with synthesis of inflammatory mediators. *J Neurovirol* 6, 350-357.
- Talalay, P. (1989). Mechanisms of induction of enzymes that protect against chemical carcinogenesis. *Adv Enzyme Regul* 28, 237-250.
- Tardieu, M., Hery, C., Peudenier, S., Boespflug, O., and Montagnier, L. (1992). Human immunodeficiency virus type 1-infected monocytic cells can destroy human neural cells after cell-to-cell adhesion. *Ann Neurol* 32, 11-17.



- Tardif, K.D., Mori, K., Kaufman, R.J., and Siddiqui, A. (2004). Hepatitis C virus suppresses the IRE1-XBP1 pathway of the unfolded protein response. *J Biol Chem* 279, 17158-17164.
- Tardif, K.D., Mori, K., and Siddiqui, A. (2002). Hepatitis C virus subgenomic replicons induce endoplasmic reticulum stress activating an intracellular signaling pathway. *J Virol* 76, 7453-7459.
- The Dana Consortium (1998). A randomized, double-blind, placebo-controlled trial of deprenyl and thioctic acid in human immunodeficiency virus-associated cognitive impairment. The Dana Consortium on the Therapy of HIV Dementia and Related Cognitive Disorders *Neurology* 50, 645-651.
- Thieblemont, N., Weiss, L., Sadeghi, H.M., Estcourt, C., and Haeffner-Cavaillon, N. (1995). CD14<sup>low</sup>CD16<sup>high</sup>: a cytokine-producing monocyte subset which expands during human immunodeficiency virus infection. *Eur J Immunol* 25, 3418-3424.
- Toggas, S.M., Masliah, E., and Mucke, L. (1996). Prevention of HIV-1 gp120-induced neuronal damage in the central nervous system of transgenic mice by the NMDA receptor antagonist memantine. *Brain Res* 706, 303-307.
- Tong, N., Perry, S.W., Zhang, Q., James, H.J., Guo, H., Brooks, A., Bal, H., Kinnear, S.A., Fine, S., Epstein, L.G., *et al.* (2000). Neuronal fractalkine expression in HIV-1 encephalitis: roles for macrophage recruitment and neuroprotection in the central nervous system. *J Immunol* 164, 1333-1339.
- Tong, N., Sanchez, J.F., Maggirwar, S.B., Ramirez, S.H., Guo, H., Dewhurst, S., and Gelbard, H.A. (2001). Activation of glycogen synthase kinase 3 beta (GSK-3beta) by platelet activating factor mediates migration and cell death in cerebellar granule neurons. *Eur J Neurosci* 13, 1913-1922.
- Ubogu, E.E., Cossoy, M.B., and Ransohoff, R.M. (2006). The expression and function of chemokines involved in CNS inflammation. *Trends Pharmacol Sci* 27, 48-55.
- Umareddy, I., Pluquet, O., Wang, Q.Y., Vasudevan, S.G., Chevet, E., and Gu, F. (2007). Dengue virus serotype infection specifies the activation of the unfolded protein response. *J Virol* 4, 91.
- Urano, F., Wang, X., Bertolotti, A., Zhang, Y., Chung, P., Harding, H.P., and Ron, D. (2000). Coupling of stress in the ER to activation of JNK protein kinases by transmembrane protein kinase IRE1. *Science* 287, 664-666.
- Uttara, B., Singh, A.V., Zamboni, P., and Mahajan, R.T. (2009). Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options. *Curr Neuropharmacol* 7, 65-74.
- van Marle, G., Henry, S., Todoruk, T., Sullivan, A., Silva, C., Rourke, S.B., Holden, J., McArthur, J.C., Gill, M.J., and Power, C. (2004). Human immunodeficiency virus type 1 Nef protein mediates neural cell death: a neurotoxic role for IP-10. *Virology* 329, 302-318.
- van Ommen, B., Koster, A., Verhagen, H., and van Bladeren, P.J. (1992). The glutathione conjugates of tert-butyl hydroquinone as potent redox cycling agents and possible reactive agents underlying the toxicity of butylated hydroxyanisole. *Biochem Biophys Res Commun* 189, 309-314.

- Vandermeeren, M., Janssens, S., Borgers, M., and Geysen, J. (1997). Dimethylfumarate is an inhibitor of cytokine-induced E-selectin, VCAM-1, and ICAM-1 expression in human endothelial cells. *Biochem Biophys Res Commun* 234, 19-23.
- Vandermeeren, M., Janssens, S., Wouters, H., Borghmans, I., Borgers, M., Beyaert, R., and Geysen, J. (2001). Dimethylfumarate is an inhibitor of cytokine-induced nuclear translocation of NF-kappa B1, but not RelA in normal human dermal fibroblast cells. *J Invest Dermatol* 116, 124-130.
- Vawter, M.P., Dillon-Carter, O., Tourtellotte, W.W., Carvey, P., and Freed, W.J. (1996). TGFbeta1 and TGFbeta2 concentrations are elevated in Parkinson's disease in ventricular cerebrospinal fluid. *Exp Neurol* 142, 313-322.
- Vergote, D., Butler, G.S., Ooms, M., Cox, J.H., Silva, C., Hollenberg, M.D., Jhamandas, J.H., Overall, C.M., and Power, C. (2006). Proteolytic processing of SDF-1alpha reveals a change in receptor specificity mediating HIV-associated neurodegeneration. *Proc Natl Acad Sci USA* 103, 19182-19187.
- von Giesen, H.J., Adams, O., Koller, H., and Arendt, G. (2005). Cerebrospinal fluid HIV viral load in different phases of HIV-associated brain disease. *J Neurol* 252, 801-807.
- Wang, H., Sun, J., and Goldstein, H. (2008). Human immunodeficiency virus type 1 infection increases the in vivo capacity of peripheral monocytes to cross the blood-brain barrier into the brain and the in vivo sensitivity of the blood-brain barrier to disruption by lipopolysaccharide. *J Virol* 82, 7591-7600.
- Wang, X.Z., Harding, H.P., Zhang, Y., Jolicoeur, E.M., Kuroda, M., and Ron, D. (1998). Cloning of mammalian Ire1 reveals diversity in the ER stress responses. *Eur Mol Biol Org J* 17, 5708-5717.
- Wang, Y., White, M.G., Akay, C., Chodroff, R.A., Robinson, J., Lindl, K.A., Dichter, M.A., Qian, Y., Mao, Z., Kolson, D.L., *et al.* (2007). Activation of cyclin-dependent kinase 5 by calpains contributes to human immunodeficiency virus-induced neurotoxicity. *J Neurochem* 103, 439-455.
- Watabe, K., Saida, T., and Kim, S.U. (1989). Human and simian glial cells infected by human T-lymphotropic virus type I in culture. *J Neuropathol Exp Neurol* 48, 610-619.
- Wei, M.C., Zong, W.X., Cheng, E.H., Lindsten, T., Panoutsakopoulou, V., Ross, A.J., Roth, K.A., MacGregor, G.R., Thompson, C.B., and Korsmeyer, S.J. (2001). Proapoptotic BAX and BAK: a requisite gateway to mitochondrial dysfunction and death. *Science* 292, 727-730.
- Wei, X., Decker, J.M., Wang, S., Hui, H., Kappes, J.C., Wu, X., Salazar-Gonzalez, J.F., Salazar, M.G., Kilby, J.M., Saag, M.S., *et al.* (2003). Antibody neutralization and escape by HIV-1. *Nature* 422, 307-312.
- Werner-Felmayer, G., Werner, E.R., Fuchs, D., Hausen, A., Reibnegger, G., and Wachter, H. (1989). Tumour necrosis factor-alpha and lipopolysaccharide enhance interferon-induced tryptophan degradation and pteridine synthesis in human cells. *Biol Chem Hoppe Seyler* 370, 1063-1069.
- Wesselingh, S.L., Power, C., Glass, J.D., Tyor, W.R., McArthur, J.C., Farber, J.M., Griffin, J.W., and Griffin, D.E. (1993). Intracerebral cytokine messenger RNA expression in acquired immunodeficiency syndrome dementia. *Ann Neurol* 33, 576-582.

West, M.J., Lowe, A.D., and Karn, J. (2001). Activation of Human Immunodeficiency Virus Transcription in T Cells Revisited: NF- $\kappa$ B p65 Stimulates Transcriptional Elongation. *J Virol* 75, 8524-8537.

White, M.G., Wang, Y., Akay, C., Lindl, K.A., Kolson, D.L., and Jordan-Sciutto, K.L. (2011). Parallel high throughput neuronal toxicity assays demonstrate uncoupling between loss of mitochondrial membrane potential and neuronal damage in a model of HIV-induced neurodegeneration. *Neurosci Res*.

WHO (2009). AIDS Epidemic update 2009 (UNAIDS).

WHO (2010). AIDS Epidemic update 2010 (UNAIDS).

Wiley, C.A., and Achim, C. (1994). Human immunodeficiency virus encephalitis is the pathological correlate of dementia in acquired immunodeficiency syndrome. *Ann Neurol* 36, 673-676.

Williams, R., Dhillon, N.K., Hegde, S.T., Yao, H., Peng, F., Callen, S., Chebloune, Y., Davis, R.L., and Buch, S.J. (2009a). Proinflammatory cytokines and HIV-1 synergistically enhance CXCL10 expression in human astrocytes. *Glia* 57, 734-743.

Williams, R., Yao, H., Dhillon, N.K., and Buch, S.J. (2009b). HIV-1 Tat co-operates with IFN-gamma and TNF-alpha to increase CXCL10 in human astrocytes. *PLoS One* 4, e5709.

Wilms, H., Sievers, J., Rickert, U., Rostami-Yazdi, M., Mrowietz, U., and Lucius, R. (2010). Dimethylfumarate inhibits microglial and astrocytic inflammation by suppressing the synthesis of nitric oxide, IL-1beta, TNF-alpha and IL-6 in an in-vitro model of brain inflammation. *J Neuroinflammation* 7, 30.

Wu, D.T., Woodman, S.E., Weiss, J.M., McManus, C.M., D'Aversa, T.G., Hesselgesser, J., Major, E.O., Nath, A., and Berman, J.W. (2000). Mechanisms of leukocyte trafficking into the CNS. *J Neurovirol* 6 Suppl 1, S82-85.

Wurdig, S., and Kugler, P. (1991). Histochemistry of glutamate metabolizing enzymes in the rat cerebellar cortex. *Neurosci Letters* 130, 165-168.

Wylie, K.M., Schrimpf, J.E., and Morrison, L.A. (2009). Increased eIF2alpha phosphorylation attenuates replication of herpes simplex virus 2 vhs mutants in mouse embryonic fibroblasts and correlates with reduced accumulation of the PKR antagonist ICP34.5. *J Virol* 83, 9151-9162.

Yadav, A., and Collman, R.G. (2009). CNS Inflammation and Macrophage/Microglial Biology Associated with HIV-1 Infection. *J Neuroimmune Pharmacol*.

Yamamoto, K., Yoshida, H., Kokame, K., Kaufman, R.J., and Mori, K. (2004). Differential contributions of ATF6 and XBP1 to the activation of endoplasmic reticulum stress-responsive cis-acting elements ERSE, UPR and ERSE-II. *J Biochem* 136, 343-350.

Yamano, Y., Nagai, M., Brennan, M., Mora, C.A., Soldan, S.S., Tomaru, U., Takenouchi, N., Izumo, S., Osame, M., and Jacobson, S. (2002). Correlation of human T-cell lymphotropic virus type 1 (HTLV-1) mRNA with proviral DNA load, virus-specific CD8(+) T cells, and disease severity in HTLV-1-associated myelopathy (HAM/TSP). *Blood* 99, 88-94.

- Yan, M., Shen, J., Person, M.D., Kuang, X., Lynn, W.S., Atlas, D., and Wong, P.K. (2008). Endoplasmic reticulum stress and unfolded protein response in Atm-deficient thymocytes and thymic lymphoma cells are attributable to oxidative stress. *Neoplasia* 10, 160-167.
- Yao, H., Peng, F., Dhillon, N., Callen, S., Bokhari, S., Stehno-Bittel, L., Ahmad, S.O., Wang, J.Q., and Buch, S. (2009). Involvement of TRPC channels in CCL2-mediated neuroprotection against tat toxicity. *J Neurosci* 29, 1657-1669.
- Ye, J., Rawson, R.B., Komuro, R., Chen, X., Dave, U.P., Prywes, R., Brown, M.S., and Goldstein, J.L. (2000). ER stress induces cleavage of membrane-bound ATF6 by the same proteases that process SREBPs. *Mol Cell* 6, 1355-1364.
- Yoshida, H., Matsui, T., Hosokawa, N., Kaufman, R.J., Nagata, K., and Mori, K. (2003). A time-dependent phase shift in the mammalian unfolded protein response. *Dev Cell* 4, 265-271.
- Yoshida, H., Matsui, T., Yamamoto, A., Okada, T., and Mori, K. (2001). XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell* 107, 881-891.
- Yoshida, H., Okada, T., Haze, K., Yanagi, H., Yura, T., Negishi, M., and Mori, K. (2000). ATF6 activated by proteolysis binds in the presence of NF-Y (CBF) directly to the cis-acting element responsible for the mammalian unfolded protein response. *Mol Cell Biol* 20, 6755-6767.
- Yu, C.Y., Hsu, Y.W., Liao, C.L., and Lin, Y.L. (2006). Flavivirus infection activates the XBP1 pathway of the unfolded protein response to cope with endoplasmic reticulum stress. *Journal of Virology* 80, 11868-11880.
- Zhan, Q., Lord, K.A., Alamo, I., Jr., Hollander, M.C., Carrier, F., Ron, D., Kohn, K.W., Hoffman, B., Liebermann, D.A., and Fornace, A.J., Jr. (1994). The gadd and MyD genes define a novel set of mammalian genes encoding acidic proteins that synergistically suppress cell growth. *Mol Cell Biol* 14, 2361-2371.
- Zhang, K., and Kaufman, R.J. (2008). From endoplasmic-reticulum stress to the inflammatory response. *Nature* 454, 455-462.
- Zhang, K., Rana, F., Silva, C., Ethier, J., Wehrly, K., Chesebro, B., and Power, C. (2003). Human immunodeficiency virus type 1 envelope-mediated neuronal death: uncoupling of viral replication and neurotoxicity. *J Virol* 77, 6899-6912.
- Zhang, L., He, T., Talal, A., Wang, G., Frankel, S.S., and Ho, D.D. (1998). In vivo distribution of the human immunodeficiency virus/simian immunodeficiency virus coreceptors: CXCR4, CCR3, and CCR5. *J Virol* 72, 5035-5045.
- Zhao, N., Fogg, J.M., Zechiedrich, L., and Zu, Y. (2011). Transfection of shRNA-encoding Minivector DNA of a few hundred base pairs to regulate gene expression in lymphoma cells. *Gene Ther* 18, 220-224.
- Zhao, Y., Navia, B.A., Marra, C.M., Singer, E.J., Chang, L., Berger, J., Ellis, R.J., Kolson, D.L., Simpson, D., Miller, E.N., *et al.* Memantine for AIDS dementia complex: open-label report of ACTG 301. *HIV Clin Trials* 11, 59-67.
- Zheng, J., Ghorpade, A., Niemann, D., Cotter, R.L., Thylin, M.R., Epstein, L., Swartz, J.M., Shepard, R.B., Liu, X., Nukuna, A., *et al.* (1999a). Lymphotropic virions affect chemokine

receptor-mediated neural signaling and apoptosis: implications for human immunodeficiency virus type 1-associated dementia. *J Virol* 73, 8256-8267.

Zheng, J., Thylin, M.R., Ghorpade, A., Xiong, H., Persidsky, Y., Cotter, R., Niemann, D., Che, M., Zeng, Y.C., Gelbard, H.A., *et al.* (1999b). Intracellular CXCR4 signaling, neuronal apoptosis and neuropathogenic mechanisms of HIV-1-associated dementia. *J Neuroimmunol* 98, 185-200.

Zhou, H., Lapointe, B.M., Clark, S.R., Zbytnuik, L., and Kubes, P. (2006). A requirement for microglial TLR4 in leukocyte recruitment into brain in response to lipopolysaccharide. *J Immunol* 177, 8103-8110.

Zhou, H., Pandak, W.M., Jr., Lyall, V., Natarajan, R., and Hylemon, P.B. (2005). HIV protease inhibitors activate the unfolded protein response in macrophages: implication for atherosclerosis and cardiovascular disease. *Mol Pharmacol* 68, 690-700.

Zhou, L.Z.H., Johnson, A.P., and Rando, T.A. (2001). NF[ $\kappa$ ]B and AP-1 mediate transcriptional responses to oxidative stress in skeletal muscle cells. *Free Rad Biol Med* 31, 1405-1416.

Zhu, R., Zhang, Y.B., Chen, Y.D., Dong, C.W., Zhang, F.T., Zhang, Q.Y., and Gui, J.F. (2006). Molecular cloning and stress-induced expression of *paralichthys olivaceus* heme-regulated initiation factor 2 $\alpha$  kinase. *Dev Comp Immunol* 30, 1047-1059.

Zink, M.C., Coleman, G.D., Mankowski, J.L., Adams, R.J., Tarwater, P.M., Fox, K., and Clements, J.E. (2001). Increased macrophage chemoattractant protein-1 in cerebrospinal fluid precedes and predicts simian immunodeficiency virus encephalitis. *J Infect Dis* 184, 1015-1021.

Zink, M.C., Spelman, J.P., Robinson, R.B., and Clements, J.E. (1998). SIV infection of macaques--modeling the progression to AIDS dementia. *J Neurovirol* 4, 249-259.

Zink, M.C., Uhrlaub, J., DeWitt, J., Voelker, T., Bullock, B., Mankowski, J., Tarwater, P., Clements, J., and Barber, S. (2005). Neuroprotective and anti-human immunodeficiency virus activity of minocycline. *J Am Med Assoc* 293, 2003-2011.

Zinszner, H., Kuroda, M., Wang, X., Batchvarova, N., Lightfoot, R.T., Remotti, H., Stevens, J.L., and Ron, D. (1998). CHOP is implicated in programmed cell death in response to impaired function of the endoplasmic reticulum. *Genes Dev* 12, 982-995.