## University of Pennsylvania Scholarly Commons

Department of Physics Papers

Department of Physics

11-13-2009

## Elasticity and Response in Nearly Isostatic Periodic Lattices

Anton Souslov *University of Pennsylvania*, asouslov@sas.upenn.edu

Andrea J. Liu *University of Pennsylvania*, ajliu@physics.upenn.edu

Tom C. Lubensky *University of Pennsylvania*, tom@physics.upenn.edu

Suggested Citation:

Souslov, A., A.J. Liu and T.C. Lubensky. (2009). "Elasticity and Response in Nearly Isostatic Periodic Lattices." Physical Review Letters. 103, 205503.

@ 2009 The American Physical Society http://dx.doi.org/10.1103/PhysRevLett.103.205503.

 $This paper is posted at Scholarly Commons. \\ http://repository.upenn.edu/physics_papers/42 \\ For more information, please contact repository@pobox.upenn.edu.$ 

## **Elasticity and Response in Nearly Isostatic Periodic Lattices**

Anton Souslov, Andrea J. Liu, and T. C. Lubensky

Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA (Received 12 June 2009; published 13 November 2009)

The square and kagome lattices with nearest-neighbor springs of spring constant k are isostatic with a number of zero-frequency modes that scale with their perimeter. We analytically study the approach to this isostatic limit as the spring constant k' for next-nearest-neighbor bonds vanishes. We identify a characteristic frequency  $\omega^* \sim \sqrt{k'}$  and length  $l^* \sim \sqrt{k/k'}$  for both lattices. The shear modulus  $C_{44} = k'$  of the square lattice vanishes with k', but that for the kagome lattice does not.

DOI: 10.1103/PhysRevLett.103.205503

An argument advanced by Maxwell [1] states that rigid assemblies of loose particles must have at least  $Z_{\rm iso}=2n_f$  contacts per particle, where  $n_f$  is the number of relevant degrees of freedom per particle. This result has been applied to many systems [2,3], including network glasses [4,5], rigidity percolation [6,7],  $\beta$ -cristobalite [8], granular media [9,10], protein folding [11], and elasticity in networks of semiflexible polymers [12].

Zero-temperature packings of particles can exhibit transitions with increasing particle density from an unjammed, disordered state with no interparticle contacts to a jammed, disordered state with an average of  $Z_c$  contacts per particle. These transitions are discontinuous because the coordination number  $Z_c$  must satisfy Maxwell's inequality. For the special case of frictionless spheres, the coordination number at the jamming transition,  $Z_c$ , appears to be exactly the minimal, or isostatic, value needed for mechanical stability,  $Z_{iso}$  [13–15]. The coincidence of the jamming transition with the threshold for mechanical stability gives rise to special properties at the transition: the coordination number Z jumps discontinuously but as the transition is approached from the high-density side, the shear modulus vanishes and length and time scales diverge as power laws [13–16]. In addition, the vibrational properties just above the transition are very different from those of typical solids [3,16-19].

The behavior of the isostatic jamming transition invites comparison to second-order structural transitions, in which the vanishing of an elastic constant in an isostatic lattice signals the instability towards a shape distortion. Jamming involves a transition between two disordered states, while critical structural transitions involve a crystalline state. This raises the question of which aspects of the jamming transition apply to all systems, periodic or disordered, near isostatic transitions and which do not. To answer these questions, it is important to study models for which results can be rigorously established via analytic calculations. In this Letter, we test the robustness of the connection between isostaticity, power-law elastic moduli and vibrational properties by an exact analytical exploration of the approach to the threshold of mechanical stability in fully periodic, nearly isostatic systems. Specifically, we study PACS numbers: 63.70.+h, 63.20.D-, 64.70.kj, 83.80.Fg

two lattices, the 2D square and kagome lattices, shown in Fig. 1, with nearest-neighbor (NN) harmonic springs of spring constant k and next-nearest-neighbor (NNN) harmonic springs with spring constant k'. The isostatic structural transition is then approached continuously as  $k' \rightarrow 0$  since both systems are isostatic there with  $Z_c = 4$ . To describe the phase that results when k' < 0, it is necessary to add a nonlinear term  $gx^4$  to the energy [20].

Our principal results are that in both the square and kagome lattices, there is a characteristic frequency  $\omega^* \sim \sqrt{k'}$  and a length  $l^* \sim \sqrt{k/k'}$  that follow directly form the form of the wave-number- and frequency-dependent response function but that can also be obtained from cutting arguments [3,16]. However, the shear modulus exponent is different in the three systems, showing that isostaticity does not confer universality on all power-law properties near the transition.

It is useful first to understand the two states that straddle the structural transition. Above the transition, where k' > 0, the square or kagome lattice is stable. Below the transition, where k' < 0 (and g > 0), the  $N_0$  zero modes at k' = 0 develop positive or negative amplitudes, and there are  $\sim a^{N_0}$ , where a > 1, distinct ground states. In this paper, we will restrict our attention to the case  $k' \ge 0$ , and we will use the harmonic approximation.

Now consider the square lattice shown in Fig. 1(a) with NNN nonlinear springs. This lattice has  $N = N_x N_y$  sites




FIG. 1 (color online). (a) Square and (b) kagome lattices with NN springs of spring constant k and NNN springs of spring constant k'. White circles in (a) and white triangles in (b) show a zero-energy distortion.

and  $N_{nn} = 2N_xN_y - N_x - N_y$  nearest-neighbor bonds. Thus when k' = 0, this system is isostatic with a number of zero-frequency modes,  $N_0 = 2N - N_{nn} = N_x + N_y$ , equal to half the perimeter of the system. When k' > 0, these modes become quasi-isostatic with nonzero frequencies that vanish as  $k' \rightarrow 0$ . The components of the dynamical matrix are easily calculated [21]

$$D_{xx}(\mathbf{q}) = D_{yy}(q_y, q_x)$$

$$= 4k\sin^2(q_x/2) + 4k'\sin^2(q_y/2) + 4k'\sin^2(q_x/2)$$

$$- 8k'\sin^2(q_x/2)\sin^2(q_y/2),$$

$$D_{xy}(\mathbf{q}) = D_{yx}(\mathbf{q}) = 2k' \sin(q_x) \sin(q_y), \tag{1}$$

where we set the lattice constant a equal to 1. In the continuum limit  $q \ll 1$ ,  $D_{ij}(\mathbf{q})$  obtains the form dictated by elasticity theory with elastic constants  $C_{11} = C_{xxxx} = C_{yyyy}$ ,  $C_{12} = C_{xxyy}$ , and  $C_{44} = C_{xyxy}$ : [21]

$$D_{xx}(\mathbf{q}) = C_{11}q_x^2 + C_{44}q_y^2; \qquad D_{yy}(\mathbf{q}) = C_{11}q_y^2 + C_{44}q_x^2;$$
  

$$D_{xy}(\mathbf{q}) = (C_{12} + C_{44})q_xq_y, \qquad (2)$$

and we can relate k and k' to the elastic constants:  $C_{11} = k + k'$ ,  $C_{44} = C_{12} = k'$ . Thus, the shear modulus vanishes as  $k' \to 0$ . When k' = 0,  $D_{ij}(\mathbf{q})$  breaks up into two independent one-dimensional compressional phonon systems: it is diagonal with  $D_{xx}(\mathbf{q}) = 4k\sin^2(q_x/2)$  independent of  $q_y$  and  $D_{yy}(\mathbf{q}) = 4k'\sin^2(q_y/2)$  independent of  $q_x$ . Thus at  $q_x = 0$ ,  $D_{xx}(\mathbf{q})$  vanishes for all points along the line  $-\pi < q_y < \pi$ . Note that at a standard structural phase transition, components of  $D_{ij}(\mathbf{q})$  vanish at one or possibly a discrete set of points in the Brillouin Zone (BZ) when an elastic modulus vanishes. In contrast, for the periodic isostatic system, components of  $D_{ij}(\mathbf{q})$  vanish along lines.

The one-dimensional nature of  $D_{ij}(\mathbf{q})$  gives rise to compressional phonons with frequencies  $\omega_{x,y}(\mathbf{q}) = 2\sqrt{k'}|\sin q_{x,y}/2|$  [Fig. 2(a)] and a one-dimensional density of states  $\rho(\omega)(2/\pi)/\sqrt{4k-\omega^2}$  with a nonzero value  $\rho_0 = (\pi\sqrt{k})^{-1}$  at  $\omega = 0$  as shown in Fig. 3(a).

When k' > 0, modes exhibit a  $\cos 4\theta$  modulation at low frequency and one-dimensional isostatic behavior at larger




FIG. 2 (color online). (a)  $\omega_x(\mathbf{q})$  when k'=0 showing the line of zero energy at  $q_x=0$  and one-dimensional dispersion as a function of  $q_x$  at fixed  $q_y$ . (b) Density plot of the low-energy phonon mode when k'=0.02k showing  $\cos 4\theta$  modulation at small  $\mathbf{q}$  and one-dimensional isostatic behavior at large  $\mathbf{q}$ .

**q** [Fig. 2(b)]. When  $0 < k' \ll k$ ,  $D_{ij}(\mathbf{q})$  is well approximated as a diagonal matrix with  $D_{xx}(\mathbf{q}) = kq_x^2 + 4k'\sin^2(q_y/2)$  with associated eigenfrequency  $\omega_x(\mathbf{q}) \sim \sqrt{D_{xx}(\mathbf{q})}$ . These expressions immediately define a characteristic frequency

$$\omega^* = 2\sqrt{k'} \tag{3}$$

as the frequency at the point  $M=(0,\pi)$  on the BZ edge. The first term in  $D_{xx}(\mathbf{q})$  represents the long-wavelength anomalous *isostatic (iso)* modes that are present when k'=0, whereas the second represents the effects of NNN coupling. When  $q_x=0$ , the only length scale in the problem is the unit lattice spacing, and no divergent length scale can be extracted from  $D_{xx}(0,q_y)$  as it can be in the case of the structural phase transition. When the first term is large compared to the second,  $D_{xx}(\mathbf{q})$  reduces to its form for the isostatic k'=0 limit, and we can extract a length by comparing these two terms. The shortest length we can extract comes from comparing  $kq_x^2$  to  $D_{xx}(\mathbf{q})$  at point M on the zone edge, i.e.,

$$l^* = (1/2)\sqrt{k/k'} \equiv 1/q^*. \tag{4}$$

If  $q_y < \pi$ , the isostatic limit is reached when  $q_x > q^*$ . A similar analysis applies to  $D_{yy}(\mathbf{q})$  when  $q_y > q^*$ . If a square of length l is cut from the bulk, the wave numbers of its excitations will be greater than  $\pi/l$ , and for  $ql^* > 1$ , all modes within the box will be effectively isostatic ones. This construction is equivalent to the cutting argument of Wyart *et al.* [3,22].

Equation (4) is identical to the length at which the frequency of the compressional mode  $\omega_x(q_x, 0) = \sqrt{k/l^*} \sim \sqrt{C_{11}}/l^*$  becomes equal to  $\omega^*$ . A meaningful length from the transverse mode  $\omega_x(0, q_y)$  cannot be extracted in a similar fashion. The full phonon spectrum [Fig. 3(a)] exhibits acoustic phonons identical to those of a standard square lattice at  $q \ll 1$  and a saddle point at the point M. Thus, the low-frequency density of states is Debye-like:  $\rho(\omega) = (\omega/(2\pi))/\sqrt{kk'}$  with a denominator that, because of the anisotropy of the square lattice, is proportional to the geometric mean of longitudinal and trans-




FIG. 3 (color online). Density of states as a function of  $\eta = \omega/\omega^*$  for (a) the square and (b) kagome lattices, showing their constant value when k' = 0 (dashed green line) and van Hove singularities, from the saddle at S and minima at M and  $\Gamma$ , whose frequencies all scale as  $\omega^* \propto \sqrt{k'}$  when k' > 0.

verse sound velocities rather than to a single velocity. In addition  $\rho(\omega)$  exhibits a logarithmic van Hove singularity at  $\omega^*$  and approaches the one-dimensional limit  $(1/\pi)/\sqrt{k}$  at  $\omega^* \ll \omega \ll 2\sqrt{k}$ . The frequency  $\omega^*$  [Eq. (3)] is recovered by equating the low-frequency Debye form at  $\omega^*$  to the high-frequency isostatic form of the density of states.

The kagome lattice can be viewed as an array of one-dimensional staggered linear rows, parallel to the x axis, of pairs of opposing triangles as shown in Fig. 1. Identical arrays with rows parallel to  $(\cos 2\pi/3, \pm \sin 2\pi/3)$  can be identified. As in the square lattice, each site has 2d = 4 nearest neighbors in the bulk. A counting procedure similar to that used for the square lattice yields  $N_0$  proportional to the  $\sqrt{N}$  for lattice of N sites.

The isostatic (iso) modes correspond to identical rotations about their top vertices of all of the "up" pointing triangles in any row in the horizontal (or symmetryequivalent) grid. These rotations require counterrotations of connected "down" pointing triangles as shown in Fig. 1(b). There are three sites per unit cell (which we take to be those in the up triangles) in the kagome lattice and six phonon branches [Fig. 4(a)]. Three of these are high-energy optical branches, two are acoustic and one is isostatic. The zero modes of the latter show up as three lines of zero frequency along  $q_x = 0$  [ $\Gamma = (0, 0)$  to M = $(0, G_0/2)$ , where  $G_0 = 4\pi/\sqrt{3}$ , in the BZ] and the two symmetry-related lines as shown in Fig. 5(a). Away from  $q_x = 0$ , the isostatic mode frequency is  $\omega_I(\mathbf{q}) = c_x q_x$ , where  $c_x = \sqrt{3k}/4$  for small  $q_x$ . This behavior is identical to that of the square lattice. The resulting density of states decreases linearly with  $\omega$  from a nonzero value  $\rho_0$  =  $3G_0/(4\pi^2c_x) = 8/(\pi\sqrt{k})$  at  $\omega = 0$ . The total lowfrequency density of states from the two acoustic modes and the isostatic mode is independent of  $\omega$  and equal to  $\rho_0$ at small  $\omega$  [Fig. 3(b)].

When springs of spring constant k' are added to the NNN bonds shown in Fig. 1(b), the quasi-isostatic mode along  $q_x = 0$  [Fig. 4(b)] has nonzero frequency of order  $\sqrt{k'}$  for all  $q_y > 0$  and gives rise to various lengths of order  $l^* \sim$ 




FIG. 4 (color online). (a) Phonon dispersion along symmetry directions. The dotted lines are for k'=0 and the solid lines are for k'=0.02. The isostatic and quasi-isostatic branches are in red. (b) shows isostatic and shear modes along  $\Gamma M$  and indicates characteristic frequencies and wave-numbers.

 $\sqrt{k/k'}$ . At  $q_x = 0$  and low values of  $q_y$ , this mode hybridizes with the transverse phonon mode to produce a gapped translation-rotation mode with frequency

$$\omega^* = \sqrt{6k'} \tag{5}$$

at  ${\bf q}=0$ . At small  ${\bf q}$ , there are isotropic longitudinal and transverse sound modes with respective velocities  $c_L=\sqrt{3k}/4$  and  $c_T=\sqrt{k}/4$ . The lowest frequency mode after hybridization at  $q_x=0$  is a transverse phonon near  $q_y=0$  and predominantly an isostatic rotation mode at  $q_y>q_H^*$ , where  $q_H^*=4\sqrt{3k'/k}\equiv 1/l_H^*$  can be termed a hybridization wave number. This mode reaches a maximum frequency  $\omega_S^*=\omega^*/\sqrt{2}$  at a saddle point at  $q_y=Q_S=4(3k'/2k)^{1/4}\sim (l^*)^{-1/2}$  and a local minimum with frequency  $\omega_M^*=\omega^*/\sqrt{3}$  at the zone-edge point M. At low frequency, the DOS is Debye-like:  $\rho(\omega)=(\omega/2\pi)\times (c_L^{-2}+c_T^{-2})=32\omega/(2\pi k)$ . The points M,  $\Gamma$  and  $S=(0,Q_S)$  give rise to van Hove singularities in the DOS. The minimum point M produces a jump  $\Delta\rho_M=8\sqrt{2}/\pi\sqrt{k}>\rho_0$  at  $\omega_M^*$ , the saddle S produces a logarithmic singularity at  $\omega_S^*$ , and the minimum at  $\Gamma$  a jump  $\Delta\rho_\Gamma=(16/\pi)\sqrt{3k'/2k}\ll\rho_0$ , which is just visible in Fig. 3(b).

Lengths that scale as  $\sqrt{k/k'}$  can be introduced in much the same way as in the square lattice. The square of the low-frequency isostatic-shear mode at  $q_x=0$  increases as  $c_x^2q_x^2$  for nonzero shear, and lengths  $l_S^*=(1/4)\sqrt{k/k'}$  and  $l_M^*=(\sqrt{6}/8)\sqrt{k/k'}$  follow from comparing  $c_xq_x$  to  $\omega_S^*$  and  $\omega_M^*$ , respectively. These two lengths are longer than the hybridization length,  $l_H^*$ . Thus, it is only at length scales less than  $l_H^*$  that isostatic modes are retrieved completely, and we should take

$$l^* = l_H^* = (1/4)\sqrt{k/(3k')}.$$
 (6)

The less divergent length  $\xi_S = Q_S^{-1} \sim \sqrt{l^*}$  determines the position of the saddle S and does not describe the same physics as the other lengths.

The long-wavelength, low-frequency properties of the kagome lattice are best understood by considering the effective low-energy dynamical matrix  $\tilde{D}_{ij}$  obtained by integrating out the high-energy modes. This matrix is con-




FIG. 5 (color online). (a) Plot of iso mode at k' = 0. (b) density plot of iso mode at k' = 0.02k. Inset: circular symmetry at origin.

TABLE I. Frequencies, lengths, and elastic moduli in square and kagome lattices and the marginally jammed (MJ) state of disordered sphere packings.  $\omega^*$ ,  $l^*$  and B scale the same way in all lattices if we take  $k' \sim (\Delta z)^2$ , but G does not.

Quantity	Square	Kagome	MJ
$l^*$	$\frac{(\frac{k}{k'})^{1/2}}{(k')^{1/2}}$	$\frac{(\frac{k}{k'})^{1/2}}{(k')^{1/2}}$	$(\Delta z)^{-1}$
$\omega^*$	$(k')^{1/2}$	$(k')^{1/2}$	$(\Delta z)^1$
G	k'	k	$(\Delta z)^1$ $(\Delta z)^0$
B	k	k	$(\Delta z)^0$

veniently represented in a basis consisting of the longitudinal (L) and transverse (T) phonons and the mixed rotation-translation gapped mode  $(\eta)$  at  $\mathbf{q}$ :  $\tilde{D}_{LT} = \tilde{D}_{TL} = 0$ , and components are

$$\begin{split} \tilde{D}_{LL} &= 3kq^2/16; & \tilde{D}_{TT} = kq^2/16; \\ \tilde{D}_{\eta\eta} &= 6k' + kq^2/16 & \tilde{D}_{L\eta} = \tilde{D}_{\eta L}^* = kq^2\cos(3\theta)/16 \\ \tilde{D}_{T\eta} &= \tilde{D}_{\eta T}^* = -i(\sqrt{3}/2)k'q + kq^2\sin(3\theta)/16, \end{split}$$
 (7

where  $\theta$  is the angle that  $\mathbf{q}$  makes with the x axis. When k'=0, the transverse phonon and rotation-translation mode mix at  $q_x=0$  ( $\theta=\pi/2$ ) to produce the zero isostatic rotation mode and a sound mode with  $\omega=\sqrt{k/8}q_y$ . In addition, there is the longitudinal mode with  $\omega=(\sqrt{3k}/4)q_y$ . When  $k'\neq 0$ , the off-diagonal terms can be ignored to lowest order in  $\mathbf{q}$ , leading to longitudinal and transverse sound modes with respective velocities  $c_L=\sqrt{B}$  and  $c_T=\sqrt{G}$ , arising from bulk and shear moduli B=3k/16 and G=k/16. Note that the shear modulus is proportional to k; this is in contrast to the square lattice where  $G \propto k'$ . To understand how this comes about, it is instructive to look at the TT component of the phonon susceptibility,

$$\chi_{TT}(q,\theta) = D_{TT}^{-1}(q,\theta) = \frac{16}{kq^2} \frac{6 + (ql^*)^2 (3 - \cos^2 3\theta)}{6 + 2(ql^*)^2 \cos^2 3\theta}.$$
 (8)

When  $ql^* \ll 1$ , this reduces to the required isotropic form  $1/(Gq^2)$ . When  $ql^* \gg 1$ ,  $\chi_{TT}$  is anisotropic with value  $1/(Gq^2)$  for  $\cos 3\theta = 1$  and 1/(6k') for  $\cos 3\theta = 0$ .

We can now compare the properties of nearly isostatic lattices with those of disordered sphere packings. Marginally-jammed packings with volume fraction  $\phi \equiv \phi_c + \Delta \phi$ , just above the volume fraction  $\phi_c$  at the jamming threshold, have an average number of contacts per sphere  $z \equiv z_c + \Delta z$ , where  $z_c = 2d$ , and  $\Delta z \sim (\Delta \phi)^{1/2}$  [13,15]. They are macroscopically isotropic with bulk and shear moduli that for harmonic interparticle potentials scale, respectively, as  $B \sim (\Delta z)^0$  and  $G \sim (\Delta z)^1$  [13,15]. The longitudinal and transverse sound velocities then scale as  $c_L \sim (\Delta z)^0$  and  $c_T \sim (\Delta z)^{1/2}$ . The density of states has a plateau above a frequency  $\omega^*$  [15,16]. A length scale  $l^* \sim 1/\Delta z$  can be extracted by equating  $c_L/l^*$  to  $\omega^*$  [16], or from the response to a point perturbation [23]. This scaling emerges from the cutting arguments of Refs. [3,17]. A

second length  $l_T^* \sim (\Delta z)^{-1/2}$  can be defined from  $c_T/l_T^* = \omega^*$  [16] and is important for energy transport [19].

The most robust features are the scalings of  $\omega^* \sim \sqrt{k'}$ and  $l^* \sim 1/\sqrt{k'}$  in both the square and kagome lattices. Effective medium calculations [24] for these lattices with NNN bonds added with probability  $P \sim \Delta z$  yield  $k' \sim$  $(\Delta z)^2$ ; this correspondence yields the observed scalings of  $\omega^*$  and  $l^*$  for marginally-jammed systems. These scalings are likely to be robust for all nearly-isostatic systems. However, the power-law scaling of the shear modulus is not universal (Table I). The shear moduli of marginallyjammed systems and the square lattice both vanish with  $\Delta z$ (with different powers), but G of the kagome lattice is proportional to k and does not vanish. Thus, different isostatic transitions can fall into different universality classes. This conclusion is consistent with recent numerical results for disordered systems near isostaticity, which show different scalings of bulk moduli [25].

We thank NSF-DMR-0520020 (A.S.), NSF-DMR-0804900 (T.C.L.), and DE-FG02-05ER46199 (A.J.L.).

- [1] J. C. Maxwell, Philos. Mag. 27, 294 (1865).
- [2] S. Alexander, Phys. Rep. 296, 65 (1998).
- [3] M. Wyart, Ann. Phys. (Paris) 30, 1 (2005).
- [4] J. C. Phillips, J. Non-Cryst. Solids 43, 37 (1981).
- [5] M. F. Thorpe, J. Non-Cryst. Solids 57, 355 (1983).
- [6] D. J. Jacobs and M. F. Thorpe, Phys. Rev. Lett. 75, 4051 (1995).
- [7] P. M. Duxbury et al., Phys. Rev. E 59, 2084 (1999).
- [8] I.P. Swainson and M.T. Dove, Phys. Rev. Lett. **71**, 193 (1993); **71**, 3610 (1993).
- [9] S. F. Edwards and D. V. Grinev, Phys. Rev. Lett. 82, 5397 (1999).
- [10] A. V. Tkachenko and T. A. Witten, Phys. Rev. E 60, 687 (1999).
- [11] A. J. Rader et al., Proc. Natl. Acad. Sci. U.S.A. 99, 3540 (2002).
- [12] C. Heussinger, B. Schaefer, and E. Frey, Phys. Rev. E 76, 031906 (2007).
- [13] D. J. Durian, Phys. Rev. Lett. 75, 4780 (1995).
- [14] C. S. O'Hern et al., Phys. Rev. Lett. 88, 075507 (2002).
- [15] C.S. O'Hern et al., Phys. Rev. E 68, 011306 (2003).
- [16] L. E. Silbert, A. J. Liu, and S. R. Nagel, Phys. Rev. Lett. 95, 098301 (2005).
- [17] M. Wyart, S. R. Nagel, and T. A. Witten, Europhys. Lett. 72, 486 (2005).
- [18] N. Xu et al., Phys. Rev. Lett. 98, 175502 (2007).
- [19] N. Xu et al., Phys. Rev. Lett. 102, 038001 (2009).
- [20] A. Souslov and T. C. Lubensky (unpublished).
- [21] N. W. Ashcroft and N. D. Mermin, *Solid State Physics* (Hold, Rinehart, and Winston, New York, 1976).
- [22] M. Wyart et al., Phys. Rev. E 72, 051306 (2005).
- [23] W. G. Ellenbroek et al., Phys. Rev. Lett. 97, 258001 (2006).
- [24] X. Mao and T. Lubensky (unpublished).
- [25] W. G. Ellenbroek, Z. Zeravcic, W. van Saarloos and M. van Hecke, arXiv:0907.0012.