



10-15-2009

# Flat 3-brane with Tension in Cascading Gravity

Claudia de Rham

*Perimeter Institute for Theoretical Physics, McMaster University*

Justin Khoury

*Perimeter Institute for Theoretical Physics, University of Pennsylvania, jkhoury@sas.upenn.edu*

Andrew Tolley

*Perimeter Institute for Theoretical Physics*

Follow this and additional works at: [http://repository.upenn.edu/physics\\_papers](http://repository.upenn.edu/physics_papers)

 Part of the [Physics Commons](#)

## Recommended Citation

Rham, C. d., Khoury, J., & Tolley, A. (2009). Flat 3-brane with Tension in Cascading Gravity. Retrieved from [http://repository.upenn.edu/physics\\_papers/50](http://repository.upenn.edu/physics_papers/50)

De Rham, C., Khoury, J. and Tolley, A. (2009). "Flat 3-brane with Tension in Cascading Gravity." *Physical Review Letters*. **103** 161601.

© 2009 The American Physical Society

<http://dx.doi.org/10.1103/PhysRevLett.103.161601>.

This paper is posted at ScholarlyCommons. [http://repository.upenn.edu/physics\\_papers/50](http://repository.upenn.edu/physics_papers/50)

For more information, please contact [repository@pobox.upenn.edu](mailto:repository@pobox.upenn.edu).

---

# Flat 3-brane with Tension in Cascading Gravity

## Abstract

In the cascading gravity brane-world scenario, our 3-brane lies within a succession of lower codimension branes, each with their own induced gravity term, embedded into each other in a higher dimensional space-time. In the  $(6 + 1)$ -dimensional version of this scenario, we show that a 3-brane with tension remains flat, at least for sufficiently small tension that the weak-field approximation is valid. The bulk solution is singular nowhere and remains in the perturbative regime everywhere.

## Disciplines

Physical Sciences and Mathematics | Physics

## Comments

De Rham, C., Khoury, J. and Tolley, A. (2009). "Flat 3-brane with Tension in Cascading Gravity." *Physical Review Letters*. **103** 161601.

© 2009 The American Physical Society

<http://dx.doi.org/10.1103/PhysRevLett.103.161601>.

## Flat 3-brane with Tension in Cascading Gravity

Claudia de Rham,<sup>1,2</sup> Justin Khoury,<sup>1,3</sup> and Andrew Tolley<sup>1</sup>

<sup>1</sup>*Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, Ontario, N2L 2Y5, Canada*

<sup>2</sup>*Department of Physics and Astronomy, McMaster University, Hamilton Ontario, L8S 4M1, Canada*

<sup>3</sup>*Center for Particle Cosmology, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6395, USA*

(Received 23 July 2009; published 15 October 2009)

In the cascading gravity brane-world scenario, our 3-brane lies within a succession of lower-codimension branes, each with their own induced gravity term, embedded into each other in a higher-dimensional space-time. In the  $(6 + 1)$ -dimensional version of this scenario, we show that a 3-brane with tension remains flat, at least for sufficiently small tension that the weak-field approximation is valid. The bulk solution is singular nowhere and remains in the perturbative regime everywhere.

DOI: 10.1103/PhysRevLett.103.161601

PACS numbers: 11.25.Wx, 95.36.+x, 98.80.Cq

An old idea to address the vexing problem of the cosmological constant is to confine the visible Universe on a 3-brane in a higher-dimensional space-time: vacuum energy on the brane curves the extra dimensions, but leaves the  $4d$  geometry flat [1]. While tantalizing, this proposal fails as soon as the extra dimensions are compactified; since  $4d$  general relativity is recovered below the compactification scale, the theory unavoidably succumbs to Weinberg's no-go theorem [2]. (An alternative strategy is to use compact extra dimensions to suppress radiative corrections to the cosmological constant [3].)

The situation is drastically different, and far more promising, if the extra dimensions have infinite volume [4]. In this case, gravity is approximately  $4d$  only at short distances, thanks to an Einstein-Hilbert term on the brane, but becomes *higher dimensional* in the infrared. In the Dvali-Gabadadze-Porrati (DGP) scenario [5] with one extra dimension, the gravitational force law on the brane scales as the usual  $1/r^2$  at short distances, but the asymptotes scales as  $1/r^3$  at large distances. Gravity therefore behaves as a high-pass filter [6]. This weakening of gravity suggests that vacuum energy, by virtue of being the longest-wavelength source, only *appears* small because it is *degravitated* [6,7].

The degravitation phenomenon is not realized in the original DGP model because the weakening of the force law is too shallow in the infrared [7]. This motivates exploring higher-codimension branes, i.e., a higher-dimensional bulk. Realizing these higher-codimension scenarios has proven difficult. To begin with, the simplest constructions are plagued by ghost instabilities [8,9]. Second, the  $4d$  propagator is divergent and must be regularized [10]. Furthermore, for a static bulk, the geometry for codimension  $N > 2$  has a naked singularity at finite distance from the brane, for arbitrarily small tension [4]. (Allowing the brane to inflate gives a Hubble rate on the brane *inversely* proportional to the brane tension for codimension  $N > 2$  [4].)

Recently, it was argued that these pathologies are resolved by embedding our 3-brane within a succession of

higher-dimensional branes, each with their own induced gravity term [11–13]. We refer to this framework as cascading gravity. The induced graviton kinetic term acts as a regulator for the 3-brane propagator [11,12]. In the case  $N = 2$  studied in [11], consisting of a 3-brane embedded in a 4-brane within a  $(5 + 1)$ -dimensional bulk, the ghost is cured by including a sufficiently large tension on the (flat) 3-brane [11,14]. Alternatively, the ghost is also cured when considering a higher-dimensional Einstein-Hilbert term localized on the brane [9,12].

Already with  $N = 2$ , the solution exhibits degravitation: a 3-brane with tension creates a deficit angle in the bulk while remaining flat [14]. We stress that this self-tuning mechanism crucially relies on the extra dimensions having infinite volume: if the dimensions were compact, the brane tension would have to be tuned against other branes and/or bulk fluxes [15].

Since the deficit angle must be less than  $2\pi$ , the tension allowed by the solutions considered in [11,14] is bounded by  $M_6^4$ , where the  $6d$  Planck mass  $M_6$  is itself constrained to be at most  $\sim \text{meV}$ . Given its geometrical nature, this bound is most likely an artifact of the codimension-2 case and is expected to be absent in the higher codimension.

Motivated by these considerations, in this Letter we explore cascading gravity with  $N = 3$ , consisting of a 3-brane living on a 4-brane, itself embedded in a 5-brane, together in a  $(6 + 1)$ -dimensional bulk, as sketched in Fig. 1. Including tension on the 3-brane, we derive a solution for which (i) the bulk metric is nonsingular everywhere (except, of course, for a delta-function in curvature at the 3-brane location) and asymptotically flat, and (ii) the induced 3-brane geometry is exactly flat.

Since the metric depends on 3 spatial coordinates, to proceed analytically we restrict ourselves to the weak-field approximation, corresponding to sufficiently small tension. For consistency, we check that our solution remains perturbative everywhere. We are currently working on numerically extending these solutions to the nonlinear regime of large tension.




FIG. 1 (color online). Sketch of the codimension-3 cascading gravity setup.

Unlike the case of a pure codimension-3 DGP brane in  $(6 + 1)$  dimensions, where the static bulk geometry has a naked singularity for arbitrarily small tension [4], here the bulk metric is completely smooth. This traces back to the cascading mechanism of regulating the propagator: the presence of parent branes removes the power-law divergence in the  $4d$  propagator.

As illustrated in Fig. 1, the 3 extra spatial dimensions are denoted by  $y$ ,  $z$ , and  $w$ , with the codimension-1 brane located at  $w = 0$ , the codimension-2 brane at  $z = w = 0$ , and the codimension-3 brane at  $y = z = w = 0$ . Indices in  $7d$  are denoted by  $A, B, \dots$ , indices in  $6d$  by  $a, b, \dots$ , indices in  $5d$  by  $\alpha, \beta, \dots$ , and finally indices in  $4d$  by  $\mu, \nu, \dots$ .

*I. Scalar Green's Functions.*—In solving for the metric perturbations, it is useful to first consider the scalar Green's functions, determined from the action

$$S = \frac{1}{2} \int d^7x \Psi [M_7^5 \square_7 + \delta(w) M_6^4 \square_6 + \delta^2(z, w) M_5^3 \square_5 + \delta^3(y, z, w) M_4^2 \square_4] \Psi, \quad (1)$$

where  $M_d$  denotes the “Planck” mass in  $d$  dimensions. The model has three cross-over scales:

$$m_5 = \frac{M_5^3}{M_4^2}, \quad m_6 = \frac{M_6^4}{M_5^3}, \quad \text{and} \quad m_7 = \frac{M_7^5}{M_6^4}, \quad (2)$$

marking, respectively, the transition scale from  $4d$  to  $5d$ , from  $5d$  to  $6d$ , and finally from  $6d$  to  $7d$ .

In the absence of the  $5d$  and  $4d$  kinetic terms, the propagator on the codimension-1 brane is of the DGP form [5]

$$G_6^{(0)}(z - z') = \frac{1}{M_6^4} \int \frac{d\omega}{2\pi} \frac{e^{i\omega(z-z')}}{\omega^2 + q^2 + m_7 \sqrt{q^2 + \omega^2}}, \quad (3)$$

where  $q^\alpha$  is the  $5d$  momentum, and  $\omega$  is the momentum associated with the  $z$  coordinate. The exact  $6d$  propagator is then obtained by treating the  $5d$  kinetic term as a

perturbation localized at  $z = 0$ :

$$\begin{aligned} G_6(z, z') &= G_6^{(0)}(z - z') - M_5^3 G_6^{(0)}(z) q^2 G_6^{(0)}(-z') \\ &\quad + M_5^6 G_6^{(0)}(z) q^4 G_6^{(0)}(0) G_6^{(0)}(-z') + \dots \\ &= G_6^{(0)}(z - z') - \frac{G_6^{(0)}(z) M_5^3 q^2 G_6^{(0)}(-z')}{1 + M_5^3 q^2 G_6^{(0)}(0)}. \end{aligned} \quad (4)$$

In particular the induced propagator on the codimension-2 brane is determined in terms of the integral of the higher-dimensional Green's function:

$$G_5^{(0)}(q^2) = G_6(0, 0) = \frac{1}{M_5^3} \frac{1}{q^2 + g(q^2)}; \quad (5)$$

$$g(q^2) \equiv \frac{1}{M_5^3 G_6^{(0)}(0)} = \frac{\pi m_6}{2} \frac{\sqrt{m_7^2 - q^2}}{\tanh^{-1}(\sqrt{\frac{m_7 - |q|}{m_7 + |q|}})}. \quad (6)$$

(For  $|q| > m_7$ , we assume analytic continuation from the hyperbolic tangent to its trigonometric counterpart.)

Remarkably, the codimension-1 kinetic term makes the  $5d$  propagator finite, thereby regulating the logarithmic divergence characteristic of pure codimension-2 branes. Indeed,  $G_5^{(0)} \rightarrow M_7^{-5} \log(m_7 q)$  as  $M_6 \rightarrow 0$ , and thus  $M_6$  plays the role of a physical cutoff. As another check, note that in the limit  $m_7 \rightarrow 0$  in which the bulk decouples, we recover the usual DGP result:  $G_5^{(0)} \sim 1/(q^2 + m_6 q)$ .

It is straightforward to repeat the same steps to derive the induced  $4d$  propagator on the codimension-3 brane.

*II. Cascading Gravity.*—We now proceed to the gravitational case. The  $7d$  Einstein equations are given by

$$\begin{aligned} M_7^5 G_{AB}^{(7)} &= -\delta(w) \{ \delta_A^\alpha \delta_B^\beta M_6^4 G_{ab}^{(6)} + \delta(z) \delta_A^\alpha \delta_B^\beta M_5^3 G_{\alpha\beta}^{(5)} \\ &\quad + \delta(z) \delta(y) \delta_A^\mu \delta_B^\nu [M_4^2 G_{\mu\nu}^{(4)} + \Lambda g_{\mu\nu}] \}. \end{aligned} \quad (7)$$

The effective source therefore consists of induced gravity terms on each of the branes, as well as tension  $\Lambda$  on the codimension-3 brane.

In the weak-field approximation, the  $7d$  line element can be written as  $ds^2 = (\eta_{AB} + h_{AB}) dx^A dx^B$ . As shown in Appendix A, there is enough symmetry and gauge freedom to simplify the metric to the form

$$\begin{aligned} ds^2 &= [1 + \Phi(y, z, w)] (dw^2 + dz^2 + dy^2) \\ &\quad - \frac{\Theta(w)}{2m_7} \partial_\alpha \Phi_0(y, z) dx^\alpha dw \\ &\quad + \left( 1 - \frac{\Phi(y, z, w)}{4} \right) \eta_{\mu\nu} dx^\mu dx^\nu, \end{aligned} \quad (8)$$

where  $\Phi_0(y, z) \equiv \Phi(y, z, w = 0)$  is the induced profile on the codimension-1 brane. Here  $\Theta(w)$  is the theta function:  $\Theta(w) = +1$  for  $w > 0$ , and  $-1$  for  $w < 0$ .

Substituting this ansatz into Einstein's equations (7), we find that  $\Phi$  satisfies

$$\left(\square_7 + \frac{\delta(w)}{m_7}\square_6 - \frac{3}{5}\frac{\delta^2(z, w)}{m_7 m_6}\square_5\right)\Phi = \frac{8}{5}\frac{\delta^3(y, z, w)}{M_7^5}\Lambda. \quad (9)$$

This equation is of the cascading form [12], as reviewed above. The asymptotically flat bulk solution is given by

$$\Phi(y, z, w) = e^{-|w|\sqrt{-\square_6}}\Phi_0(y, z), \quad (10)$$

where the induced profile  $\Phi_0(y, z)$  satisfies

$$\left(\square_6 - m_7\sqrt{-\square_6} - \frac{3}{5}\frac{\delta(z)}{m_6}\square_5\right)\Phi_0 = \frac{8}{5}\frac{\delta^2(y, z)}{M_6^4}\Lambda. \quad (11)$$

To solve (11), we Fourier transform to momentum space and use the 6d and 5d Green's functions given, respectively, by (3) and (5). The result is

$$\Phi_0(y, z) = \int \frac{dq_y d\omega}{(2\pi)^2} \frac{e^{i\omega z} e^{iq_y y} g(q_y) \phi(q_y)}{\omega^2 + q_y^2 + m_7\sqrt{\omega^2 + q_y^2}}, \quad (12)$$

where the Fourier transform of the codimension-2 profile,  $\phi(q_y) = \int dy e^{-iq_y y} \Phi_0(z=0, y)$ , satisfies

$$\left(\frac{3}{5}q_y^2 - g(q_y^2)\right)\phi(q_y) = \frac{8}{5M_5^3}\Lambda. \quad (13)$$

The solution to (13) can be expressed as the sum of a principal part  $\mathcal{P}$  and two homogeneous modes:

$$\phi(q_y) = \frac{8\Lambda}{5M_5^3} \mathcal{P}\left[\frac{1}{\frac{3}{5}q_y^2 - g(q_y^2)}\right] + \sum_{\sigma=\pm} C_\sigma \delta(q_y - \sigma q_0),$$

where  $\frac{3}{5}q_0^2 = g(q_0^2)$ . Requiring the field  $\Phi_0$  to be real imposes  $C_+ = C_- \equiv C$ , while requiring  $\Phi_0$  to fall as  $y \rightarrow 0$  sets  $C = 0$ . Using the resulting expression for  $\phi(q_y)$  into (12) and then into (10), we obtain the final expression for the scalar potential  $\Phi(y, z, w) = \frac{8\Lambda}{5M_6^4} \hat{\Phi}(y, z, w)$ :

$$\hat{\Phi} = \int \frac{d\omega dq_y}{(2\pi)^2} \frac{e^{-|w|\sqrt{\omega^2 + q_y^2}} e^{i\omega z} e^{iq_y y}}{\omega^2 + q_y^2 + m_7\sqrt{\omega^2 + q_y^2}} \mathcal{P}\left[\frac{g(q_y)}{\frac{3}{5}q_y^2 - g(q_y)}\right]. \quad (14)$$

This is our main result. Thanks to the cascading mechanism, which has regularized all potential divergences, this solution is finite everywhere. Figure 2 shows that  $\hat{\Phi}(y, z, w)$  is smooth everywhere and decreases with  $w$ .

As it stands, however, our framework has a ghost [8,9], as indicated by the poles at  $q_y = \pm q_0$ . There are two ways to resolve this issue. One can introduce sufficiently large tension on both the codimension-2 and -3 branes [11]: to remove the ghost, the codimension-2 tension should be  $\geq M_5^3 m_7^2$ , whereas the corresponding bound on the codimension-3 tension is yet to be determined.

Alternatively, one can regularize codimension-2 and -3 branes and include the 6d Einstein-Hilbert term localized on these objects [9,12]. Following this route, we demon-




FIG. 2 (color online). Plot of the solution for the metric potential  $\hat{\Phi}(y, z, w)$  for  $w = 0$  and  $w = 2m_7^{-1}$  in the case where  $m_6 = m_7$ .

strate in Appendix B that the poles do disappear, and that the profile for  $\hat{\Phi}(y, z, w)$  is qualitatively unchanged.

*III. Discussion.*—In this Letter, we have shown that a 3-brane with tension remains flat in the (6+1)-dimensional cascading gravity framework. In the weak-field approximation, we have obtained a bulk solution which is nowhere singular and remains perturbative everywhere.

These properties crucially depend on the existence of parent branes with finite Planck masses. Indeed, our solution goes outside the perturbative regime and acquires divergences in the limit  $M_5, M_6 \rightarrow 0$ , consistent with [4].

We are currently extending our solutions to the nonlinear regime through numerical analysis. For now, we view the present results as a tantalizing first step towards realizing the idea of Rubakov and Shaposhnikov.

We thank G. Dvali and G. Gabadadze for helpful discussions. This work was supported in part by NSERC and Ontario's MRI.

*Appendix A.*—We show that the weak-field metric can be brought to the form (10) by symmetry and gauge freedom. In the de Donder gauge,  $\partial_A h_B^A = \frac{1}{2} \partial_B h_C^C$ , (7) reduces to

$$-\frac{M_7^5}{2} \square_7 \left( h_{AB} - \frac{1}{2} \eta_{AB} h_C^C \right) = \delta(w) (T_{ab}^{(6)} - M_6^4 G_{ab}^{(6)}),$$

where the effective stress energy on the codimension-1 brane,  $T_{ab}^{(6)}$ , includes contributions from the 5d and 6d induced gravity terms. Since there is no stress energy along the  $(a, w)$  and  $(w, w)$  directions, the corresponding equations are consistently satisfied by setting  $h_{aw} = 0$  and  $h_{ww} = h^c_c$ , where  $h^c_c$  is the 6d trace. It follows that the induced gauge choice in 6d is given by  $\partial_a h_b^a = \partial_b h^c_c$ ; hence, the  $(a, b)$  components of (A1) reduce to

$$-\frac{M_7^5}{2} \square_7 (h_{ab} - \eta_{ab} h^c_c) = \delta(w) \frac{M_6^4}{2} (\square_6 h_{ab} - \partial_a \partial_b h^c_c) + \delta(w) T_{ab}^{(6)}. \quad (A1)$$

To proceed further, it is convenient to decompose  $h_{ab}$  into its trace and transverse-traceless (TT) parts:

$$h_{ab} = h_{ab}^{6dTT} + \frac{\partial_a \partial_b}{\square_6} h^c{}_c. \quad (\text{A2})$$

From (A1), the 6dTT components satisfy

$$-\frac{M_7^5}{2} \left( \square_7 + \frac{\delta(w)}{m_7} \square_6 \right) h_{ab}^{6dTT} = \delta(w) \left( T_{ab}^{(6)} - \frac{1}{5} \eta_{ab} T^{(6)} + \frac{1}{5} \frac{\partial_a \partial_b}{\square_6} T^{(6)} \right). \quad (\text{A3})$$

The symmetries of the problem allow a simple expression for the 5d components of the 6dTT part:

$$h_{\alpha\beta}^{6dTT} = -\frac{1}{4} \Phi \eta_{\alpha\beta} - \left( \frac{\square_5}{\square_6} - \frac{5}{4} \right) \frac{\partial_\alpha \partial_\beta}{\square_5} \Phi. \quad (\text{A4})$$

This follows from setting  $h_{\alpha\beta}^{5dTT} = 0$ , which is consistent with the equations of motion for the case of interest. Substituting into (A3), and using  $T_{\alpha\beta}^{(5)} = -\delta^\mu{}_\alpha \delta^\nu{}_\beta \Lambda \eta_{\mu\nu} \delta(y)$ , the resulting equation of motion for  $\Phi$  agrees with (9).

We can now be explicit about the form of the various metric components. Combining (A2) and (A4), we get

$$h_{\alpha\beta} = -\frac{1}{4} \Phi \eta_{\alpha\beta} - \left( \frac{\square_5}{\square_6} - \frac{5}{4} \right) \frac{\partial_\alpha \partial_\beta}{\square_5} \Phi + \frac{\partial_\alpha \partial_\beta}{\square_6} h^c{}_c. \quad (\text{A5})$$

And since everything is independent of  $x^\mu$ , we get  $h_{y\mu} = 0$  and  $h_{\mu\nu} = -\frac{1}{4} \Phi \eta_{\mu\nu}$ . Similarly, from (A2) we obtain

$$h_{yz} = \frac{\partial_y \partial_z}{\square_6} (h^c{}_c - \Phi); \quad h_{zz} = \frac{\partial_z^2}{\square_6} (h^c{}_c - \Phi) + \Phi; \\ h_{yy} = \frac{\partial_y^2}{\square_6} (h^c{}_c - \Phi) + \Phi. \quad (\text{A6})$$

This is equivalent to (8) after a small diffeomorphism.

*Appendix B.*—One way to cure the ghost of higher-dimension DGP models [8,9] is to consider a higher-dimensional Einstein-Hilbert term localized on the regularized brane [9,12]. Following this prescription, we will show that the solution remains finite everywhere.

When adding a 6d Einstein-Hilbert term on the regularized 4-brane, on the top of the usual 5d Einstein-Hilbert term of the form “ $\square_5 h_{\alpha\beta}$ ” we must consider excitations of transverse modes along the extra dimensions as well as the higher-dimensional mode  $h_{zz}$ . In the thin-brane limit, however, the excitations along the extra dimension become very massive, so that any term containing  $z$  derivatives can be neglected. Meanwhile,  $h_{zz}$  survives in the limit; see [12] for details.

In the 7d de Donder gauge, the Einstein equations are the same as in (A1). Setting  $h_{aw} = 0$  and  $h_{ww} = h^c{}_c$ , we have

$$-\frac{M_7^5}{2} \left( \square_7 + \frac{\delta(w)}{m_7} \square_6 \right) h_{ab} = \delta(w) \left( T_{ab}^{(6)} - \frac{1}{5} T^{(6)} \eta_{ab} \right) \quad (\text{B1})$$

with  $T_{z\alpha}^{(6)} = 0$ ,  $T_{zz}^{(6)} = M_5^3 \delta(z) R_5/2$ , and

$$T_{\alpha\beta}^{(6)} = -M_5^3 \delta(z) \left[ G_{\alpha\beta}^{(5)} + \frac{1}{2} (\square_5 h_{zz} \eta_{\alpha\beta} - \partial_\alpha \partial_\beta h_{zz}) \right] - \delta(z) \delta(y) \Lambda \eta_{\mu\nu} \delta^\mu{}_\alpha \delta^\nu{}_\beta. \quad (\text{B2})$$

Using this in the 6d part of the Einstein equations, we get  $h_{zz} = -\psi$ ,  $\square_5 h_{yy} = -4\square_5 \psi + \partial_y^2 h^\alpha{}_\alpha$ ,  $h_{\mu y} = 0$ , and  $\square_5 h_{\mu\nu} = \square_5 \psi \eta_{\mu\nu} + \partial_\mu \partial_\nu h^\alpha{}_\alpha$ , with

$$\left[ \square_7 + \frac{\delta(w)}{m_7} \square_6 + \frac{\delta^{(2)}(w, z)}{m_7 m_6} \square_5 \right] \psi = \frac{2}{5} \frac{\delta^{(3)}(w, z, y)}{M_7^5} \Lambda. \quad (\text{B3})$$

We notice that the kinetic term for  $\psi$  is now everywhere positive, signaling that the ghost has been cured. Equation (B3) is similar to (9) for  $\Phi$ , except for a redefinition of  $m_6$  and  $M_7$ . The profile for  $\psi(y, z, w) = -\frac{2\Lambda}{5M_6^5} \hat{\Psi}(y, z, w)$ ,

$$\hat{\Psi} = \int \frac{d\omega dq_y}{(2\pi)^2} \frac{e^{-|w|\sqrt{\omega^2 + q_y^2}} e^{i\omega z} e^{iq_y y}}{\omega^2 + q_y^2 + m_7 \sqrt{\omega^2 + q_y^2}} \frac{g(q_y)}{q_y^2 + g(q_y)}, \quad (\text{B4})$$

is similar to that of  $\hat{\Phi}$ , and, in particular, is free of divergences. The static solution for a codimension-3 brane with tension remains therefore well defined, at least in the weak-field approximation, in a ghost-free setup.

- 
- [1] V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. B **125**, 139 (1983).
  - [2] S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989).
  - [3] Y. Aghababae, C. P. Burgess, S. L. Parameswaran, and F. Quevedo, Nucl. Phys. **B680**, 389 (2004).
  - [4] G. Dvali, G. Gabadadze, and M. Shifman, Phys. Rev. D **67**, 044020 (2003).
  - [5] G. R. Dvali, G. Gabadadze, and M. Porrati, Phys. Lett. B **485**, 208 (2000).
  - [6] N. Arkani-Hamed, S. Dimopoulos, G. Dvali, and G. Gabadadze, arXiv:hep-th/0209227.
  - [7] G. R. Dvali, S. Hofmann, and J. Khoury, Phys. Rev. D **76**, 084006 (2007).
  - [8] S. L. Dubovsky and V. A. Rubakov, Phys. Rev. D **67**, 104014 (2003).
  - [9] G. Gabadadze and M. Shifman, Phys. Rev. D **69**, 124032 (2004).
  - [10] R. P. Geroch and J. H. Traschen, Phys. Rev. D **36**, 1017 (1987); C. de Rham, J. High Energy Phys. 01 (2008) 060.
  - [11] C. de Rham *et al.*, Phys. Rev. Lett. **100**, 251603 (2008).
  - [12] C. de Rham, S. Hofmann, J. Khoury, and A. J. Tolley, J. Cosmol. Astropart. Phys. 02 (2008) 011.
  - [13] C. de Rham, Can. J. Phys. **87**, 201 (2009).
  - [14] C. de Rham, J. Khoury, and A. J. Tolley (to be published).
  - [15] G. W. Gibbons, R. Kallosh, and A. D. Linde, J. High Energy Phys. 01 (2001) 022.