

# LOWER BOUNDS FOR GENERALIZED REGULATORS

James D. Sundstrom

A DISSERTATION

in

Mathematics

Presented to the Faculties of the University of Pennsylvania

in

Partial Fulfillment of the Requirements for the

Degree of Doctor of Philosophy

2016

Supervisor of Dissertation

---

Ted Chinburg, Professor of Mathematics

Graduate Group Chairperson

---

David Harbater, Professor of Mathematics

Dissertation Committee:

Ted Chinburg, Professor of Mathematics

Ching-Li Chai, Professor of Mathematics

Robert Strain, Professor of Mathematics

# ABSTRACT

## LOWER BOUNDS FOR GENERALIZED REGULATORS

James D. Sundstrom

Ted Chinburg

In 1999, Friedman and Skoruppa demonstrated a method to derive lower bounds for the relative regulator of an extension  $L/K$  of number fields. The relative regulator is defined using the subgroup  $E_{L/K}$  of relative units of  $L/K$ . It appears in the theta series  $\Theta_{E_{L/K}}$  associated to  $E_{L/K}$ , so an inequality relating  $\Theta_{E_{L/K}}$  and  $\Theta'_{E_{L/K}}$  provides an inequality for  $\text{Reg}(L/K)$ . This same technique can be applied to other subgroups  $E$  of the units of a number field  $L$ . In this thesis, we consider the case  $E = E_{L/K_1} \cap E_{L/K_2}$ , where  $K_1$  and  $K_2$  are real quadratic fields; the corresponding regulator grows exponentially in  $[L : \mathbb{Q}]$ .

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The Generalized Regulator</b>	<b>6</b>
<b>3</b>	<b>Theta series</b>	<b>8</b>
<b>4</b>	<b>Single-variable calculus</b>	<b>14</b>
<b>5</b>	<b>Estimation of <math>f</math> and <math>f'/f</math></b>	<b>21</b>
<b>6</b>	<b>Properties of <math>\Psi</math></b>	<b>40</b>
<b>7</b>	<b>Properties of the Saddle Point</b>	<b>45</b>
<b>8</b>	<b>Conclusion</b>	<b>55</b>
	<b>Appendices</b>	<b>57</b>
<b>A</b>	<b>Proof of Lemma 5.8</b>	<b>58</b>
A.1	Proof for $\sigma \geq 0.65$ . . . . .	59
A.2	Proof for $m \geq 140$ . . . . .	65

A.3	Proof for remaining cases . . . . .	69
-----	-------------------------------------	----

# Chapter 1

## Introduction

This thesis demonstrates how a technique of Friedman and Skoruppa [8] can be generalized. Before proceeding to the generalization, we first review their paper. Friedman and Skoruppa proved lower bounds for the relative regulator  $\text{Reg}(L/K)$  associated to an extension  $L/K$  of number fields. The relative regulator was defined by Bergé and Martinet [3], [4], [5] as follows.

Given a number field  $K$ , let  $\mathcal{O}_K$  denote the algebraic integers of  $K$ , with unit group  $\mathcal{O}_K^*$  and roots of unity  $\mu_K \subseteq \mathcal{O}_K^*$ . Let  $\mathcal{A}_K$  be the set of archimedean places of  $K$ ; for each  $v \in \mathcal{A}_K$ , let

$$e_v = \begin{cases} 1 & \text{if } v \text{ is real} \\ 2 & \text{if } v \text{ is complex.} \end{cases}$$

Let  $r_1(K)$  and  $r_2(K)$  be, respectively, the number of real and complex places of  $K$ .

Given an extension of number fields  $L/K$ , let  $E_{L/K}$  denote the group of relative units of  $L/K$ , i.e.,

$$E_{L/K} = \{\epsilon \in \mathcal{O}_L^* \mid N_{L/K}(\epsilon) \in \mu_K\}.$$

Note that  $E_{L/K}$  has rank  $r = r_{L/K} := |\mathcal{A}_L| - |\mathcal{A}_K|$ . Let  $\epsilon_1, \dots, \epsilon_r$  be fundamental relative units (free generators for  $E_{L/K}$  modulo torsion). For each  $w \in \mathcal{A}_K$ , fix some  $\tilde{w} \in \mathcal{A}_L$  lying above  $w$ . Let  $\mathcal{A}'_L$  denote the remaining places of  $L$  after each  $\tilde{w}$  is removed from  $\mathcal{A}_L$ . Then the relative regulator of  $L/K$  is defined by

$$\text{Reg}(L/K) = \left| \det(e_v \log |\epsilon_j|_v)_{\substack{v \in \mathcal{A}'_L \\ 1 \leq j \leq r}} \right|.$$

Costa and Friedman [6] proved that

$$\text{Reg}(L/K) = \frac{1}{[\mathcal{O}_K^* : \mu_K N_{L/K}(\mathcal{O}_L^*)]} \frac{\text{Reg}(L)}{\text{Reg}(K)} \leq \frac{\text{Reg}(L)}{\text{Reg}(K)}.$$

Hence a lower bound for  $\text{Reg}(L/K)$  is also a lower bound for  $\text{Reg}(L)/\text{Reg}(K)$ . Furthermore,  $\text{Reg}(L/\mathbb{Q}) = \text{Reg}(L)$ ; of course, this was already clear from the definition of  $\text{Reg}(L/\mathbb{Q})$ . Thus a lower bound for relative regulators includes a lower bound for the classical regulator as a special case.

To any subgroup  $E$  of  $\mathcal{O}_L^*$ , we can associate a theta series  $\Theta_E$ . Let  $E_{\text{tor}} = E \cap \mu_L$  denote the torsion subgroup of  $E$ . Let  $E_{\mathbb{R}} = E \otimes \mathbb{R}$ , and fix a Haar measure  $\mu$  on  $E_{\mathbb{R}}$ , so that  $\mu(E_{\mathbb{R}}/E)$  is the volume of any fundamental domain for the action of  $E$  on  $E_{\mathbb{R}}$ .

There is an embedding of  $E_{\mathbb{R}}$  into  $\mathbb{R}_+^{\mathcal{A}_L}$  given by

$$x = \sum_j \epsilon_j \otimes \xi_j \mapsto (x_v)_{v \in \mathcal{A}_L}, \quad x_v = \prod_j |\epsilon_j|_v^{\xi_j}. \quad (1.1)$$

For  $a \in L$  and  $x \in E_{\mathbb{R}}$ , set

$$\|ax\|^2 = \sum_{v \in \mathcal{A}_L} e_v |a|_v^2 x_v^2.$$

For any fractional ideal  $\mathfrak{a}$  of  $L$  and any  $t > 0$ , define

$$\Theta_E(t; \mathfrak{a}) = \frac{\mu(E_{\mathbb{R}}/E)}{|E_{\text{tor}}|} + \sum_{\substack{a \in \mathfrak{a}/E \\ a \neq 0}} \int_{E_{\mathbb{R}}} \exp(-c_{\mathfrak{a}} t \|ax\|^2) d\mu(x), \quad (1.2)$$

where the sum is taken over a complete set of representatives for the non-zero  $E$ -orbits in  $\mathfrak{a}$ ,

$$c_{\mathfrak{a}} = \pi \left( \sqrt{|\text{disc}(L)|} N_{L/\mathbb{Q}}(\mathfrak{a}) \right)^{-2/[L:\mathbb{Q}]}.$$

Friedman and Skoruppa give a proof that  $\Theta_E$  is well-defined: it is independent of the choice of representatives  $a$ , and the sum is absolutely convergent. They also observe that  $t^{[L:\mathbb{Q}]/2} \Theta_E(t; \mathfrak{a})$  is an increasing function (Prop. 2.1). Differentiating, it follows that

$$\Theta_E(t; \mathfrak{a}) + \frac{2}{[L:\mathbb{Q}]} t \Theta'_E(t; \mathfrak{a}) \geq 0. \quad (1.3)$$

Since the definition of  $\Theta_E$  involves  $\mu(E_{\mathbb{R}}/E)$  as a constant term, this inequality can be understood as a lower bound for  $\mu(E_{\mathbb{R}}/E)$ . In particular, if we take  $E = E_{L/K}$ , then it is fairly natural to normalize  $\mu$  by  $\mu(E_{\mathbb{R}}/E) = \text{Reg}(L/K)$ . Thus, by estimating the integrals in the definition of  $\Theta_E$ , we will obtain the desired lower bound for  $\text{Reg}(L/K)$ .

As a first step to understanding these integrals, Friedman and Skoruppa use the Mellin transform to prove the following (Prop. 3.1).

**Proposition 1.1.** *With notation as above (including  $E = E_{L/K}$ ),*

$$\int_{E_{\mathbb{R}}} \exp(-t \|ax\|^2) d\mu(x) = A \prod_{w \in \mathcal{A}_K} f_w(a_w + \log t),$$

where  $A = 2^{-r_{L/K}} \pi^{-r_2(L)/2}$ ,

$$f_w(y) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{-se_w[L:K]y} \Gamma(s)^{p_w+q_w} \Gamma(s + \frac{1}{2})^{q_w} ds,$$

$$a_w = \frac{2}{[L:K]} \log |N_{L/K}(a)|_w.$$

Here  $c$  is any positive number, and  $p_w$  and  $q_w$  are respectively the number of real and complex places of  $L$  extending  $w \in \mathcal{A}_K$ .

This proposition puts the integrals into a more tractable form; instead of estimating the original integrals, it suffices to understand  $f_w$ . More precisely, setting  $y = \log t$ , inequality (1.3) becomes

$$\frac{\text{Reg}(L/K)}{\#\mu_L} \geq A \sum_{\substack{a \in \mathfrak{a}/E_{L/K} \\ a \neq 0}} \left\{ -1 - \frac{2}{[L:\mathbb{Q}]} \sum_{w \in \mathcal{A}_K} \frac{f'_w}{f_w}(a_w + y) \right\} \prod_{w \in \mathcal{A}_K} f_w(a_w + y), \quad (1.4)$$

so we need to estimate  $f_w$  and  $f'_w/f_w$ . This is accomplished by the saddle-point method.

The saddle point method is summarized below, in a somewhat simplified form. See de Bruijn [7] for more details. The method applies to a contour integral of the form

$$\int_C e^{f(z)} dz.$$

We shift the contour  $C$  so that it passes through the saddle point  $\sigma$ , the point where  $f'(\sigma) = 0$ . The idea then is that we can replace  $f$  with its degree-2 Taylor approximation at  $\sigma$ , so that the integral is approximated by an integral of the form

$$\int_{-\infty}^{\infty} e^{a-bt^2} dt,$$

which we know how to evaluate. Furthermore, if we want to understand

$$\int_C e^{nf(z)} dz \quad \text{as } n \rightarrow \infty,$$

then this approximation gets better as  $n$  gets larger, so that we get an excellent description of the asymptotic behavior of the integral.

Once we have estimates for  $f_w$  and  $f'_w/f_w$ , we can plug them into inequality (1.4) and we are essentially done. Since inequality (1.4) holds for any  $y \in \mathbb{R}$ , it remains only to choose a  $y$  which gives a good bound. However, our estimates for  $f_w$  and  $f'_w/f_w$  depend on  $p_w$  and  $q_w$ . We would prefer to have lower bounds for  $\text{Reg}(L/K)$  that do not require such detailed



information about the places of  $L$ . Hence we make some effort to transform the bounds in terms of the  $p_w$  and  $q_w$  into bounds depending only on  $[L : K]$  and  $r_1(L)$ .

In short, Friedman and Skoruppa's method consists of four main steps:

1. Use the Mellin transform to replace the  $\Theta_E$  integrals with complex integrals.
2. Use the saddle-point method to estimate the complex integrals.
3. Replace these estimates with estimates that do not depend on the  $p_w$  and  $q_w$ .
4. Plug these estimates into inequality (1.3) to get lower bounds for the regulator.

In this thesis, we apply these methods to a generalized regulator for a number field  $L$  containing two real quadratic fields  $K_1$  and  $K_2$ . Specifically, we consider the regulator associated to  $E = E_{L/K_1} \cap E_{L/K_2}$ . Chapter 2 defines this regulator. Chapter 3 computes the necessary inverse Mellin transform; we find that we need to study a triple integral. Chapters 4-5 carry out step 2. Chapter 4 summarizes some results in single-variable calculus which will be needed; many of these results are quite similar (or identical) to results from Friedman and Skoruppa's original paper. Then Chapter 5 applies these results to study the relevant triple integrals. Step 3 is done in Chapters 6 and 7. Once again, Chapter 6 provides some simple results, which are applied in Chapter 7. Finally, Chapter 8 completes the argument, proving that the generalized regulator  $\text{Reg}_{K_1, K_2}(L)$  grows exponentially in  $[L : K]$ .

There is a mistake in Friedman and Skoruppa's proof of their Lemma 5.6. Fortunately, Lemma 5.6 is used only to prove Lemma 5.8. Appendix A gives a correct proof of Lemma 5.8, so their main results are all correct.

## Chapter 2

# The Generalized Regulator

Let  $K_1$  and  $K_2$  be distinct real quadratic fields, and  $L$  a number field containing  $K := K_1 K_2$ . Let  $m = [L : K] = [L : \mathbb{Q}]/4$ . Let  $\mathcal{A}_{K_1} = \{w_1, w_2\}$  and  $\mathcal{A}_{K_2} = \{w_3, w_4\}$  be the sets of archimedean places of  $K_1$  and  $K_2$ . Let  $\mathcal{A}_K = \{w_{13}, w_{14}, w_{23}, w_{24}\}$  denote the set of archimedean places of  $K$ , labeled so that  $w_{ij}$  extends  $w_i \in \mathcal{A}_{K_1}$  and  $w_j \in \mathcal{A}_{K_2}$ . Note that for any  $i \in \{1, 2\}$  and  $j \in \{3, 4\}$ ,

$$\sum_{\substack{v \in \mathcal{A}_L \\ v|w_i}} e_v = \sum_{\substack{v \in \mathcal{A}_L \\ v|w_j}} e_v = 2m, \quad \sum_{\substack{v \in \mathcal{A}_L \\ v|w_{ij}}} e_v = m. \quad (2.1)$$

For any  $w \in \mathcal{A}_{K_1 K_2}$ , let  $p_w$  and  $q_w$  denote respectively the number of real and complex places of  $L$  extending  $w$ , so that  $p_w + 2q_w = m$ . Let  $E_i$  denote the relative units of  $L/K_i$ , and define  $E = E_1 \cap E_2$ . Let  $E_{\text{tor}}$  denote the torsion subgroup of  $E$ .

We define a generalized regulator  $\text{Reg}_{K_1, K_2}(L)$  as follows. Let  $\epsilon_1, \dots, \epsilon_r$  ( $r = |\mathcal{A}_L| - 3$ ) be free generators of  $E/E_{\text{tor}}$ . Let  $\tilde{\mathcal{A}}_K$  be a set containing any three places of  $K$ , and select one place of  $L$  above each place in  $\tilde{\mathcal{A}}_K$ . Let  $\mathcal{A}'_L$  denote  $\mathcal{A}_L$  with these three places removed.

Define

$$\text{Reg}_{K_1, K_2}(L) = \left| \det(e_v \log|\epsilon_j|_v)_{\substack{v \in \mathcal{A}'_L \\ 1 \leq j \leq r}} \right|.$$

**Lemma 2.1.**  $\text{Reg}_{K_1, K_2}(L)$  is well-defined, i.e., it is independent of the choice of the  $\epsilon_j$  and of  $\mathcal{A}'_L$ .

*Proof.* Define  $\lambda: \mathcal{O}_L^* \rightarrow \mathbb{R}^{\mathcal{A}_L}$  by

$$\lambda(\epsilon) = (e_v \log|\epsilon|_v)_{v \in \mathcal{A}_L}.$$

Define  $\mathbf{x}_1 \in \mathbb{R}^{\mathcal{A}_L}$  by

$$(\mathbf{x}_1)_v = \begin{cases} e_v & \text{if } v \mid w_1, \\ 0 & \text{otherwise.} \end{cases}$$

Define similarly  $\mathbf{x}_2$  with respect to  $w_2$  and  $\mathbf{x}_3$  with respect to  $w_3$ . Let  $M$  denote the matrix with columns  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \lambda(\epsilon_1), \dots, \lambda(\epsilon_r)$ . Note that  $|\det(M)|$  is the covolume of the lattice generated by  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ , and  $\lambda(\mathcal{O}_L^*)$ , so it is independent of our choice of the  $\epsilon_j$ . Row operations show that

$$|\det(M)| = |\det(M')| \text{Reg}_{K_1, K_2}(L),$$

where  $M'$  is the  $3 \times 3$  matrix defined by

$$M' = \left( \sum_{\substack{v \mid w_i \text{ and } v \mid w_j}} e_v \right)_{\substack{1 \leq i \leq 3 \\ 1 \leq j \leq 3}}.$$

(The precise row operations to be used depend on  $\tilde{\mathcal{A}}_K$ , but the result is the same.) Since  $|\det(M')| = 4m^3$ , this proves that  $\text{Reg}_{K_1, K_2}(L) = \frac{1}{4m^3} |\det(M)|$ , so  $\text{Reg}_{K_1, K_2}(L)$  is well-defined.  $\square$

## Chapter 3

# Theta series

Let  $E \subset \mathcal{O}_L^*$  be as in the previous chapter. Let  $G = \mathbb{R}_+^{\mathcal{A}_L}$  and let  $H = \mathbb{R}_+^{\mathcal{A}_{K_1} \cup \{w_3\}}$ . Let  $\mu_G$  denote the natural Haar measure on  $G$ , namely

$$d\mu_G(g) = \prod_{v \in \mathcal{A}_L} \frac{dg_v}{g_v}.$$

Define  $\mu_H$  similarly. Let  $\mu$  be the Haar measure on  $E_{\mathbb{R}}$ , normalized so that

$$\mu(E_{\mathbb{R}}/E) = \text{Reg}_{K_1, K_2}(L).$$

Define  $\delta: G \rightarrow H$  by  $\delta(g) = (h_w)_{w \in \mathcal{A}_{K_1} \cup \{w_3\}}$ , where

$$h_w = \prod_{v|w} g_v^{e_v}.$$

Using the embedding  $E_{\mathbb{R}} \rightarrow G$  from (1.1), we get an exact sequence  $1 \rightarrow E_{\mathbb{R}} \rightarrow G \rightarrow H \rightarrow 1$ .

Let  $\sigma: H \rightarrow G$  be a section of  $\delta$ . (We will choose a particular section  $\sigma$  below.) Define an isomorphism  $\phi: E_{\mathbb{R}} \times H \rightarrow G$  by  $\phi(x, h) = x\sigma(h)$ .

**Lemma 3.1.**  $2^{r_2(L)}\mu_G \circ \phi = \mu \times \mu_H$ .

*Proof.* Since  $\mu_G \circ \phi$  is a Haar measure on  $E_{\mathbb{R}} \times H$ , we know that  $c\mu_G \circ \phi = \mu \times \mu_H$  for some constant  $c$ . Consider  $E_{\mathbb{R}}$ ,  $G$ ,  $H$ , and  $\mathbb{R}_+$  as real vector spaces. Choose any  $v_{13}, v_{23}, v_{14} \in \mathcal{A}_L$  such that  $v_{ij}$  extends  $w_i$  and  $w_j$ . Define  $g_{13}, g_{23}, g_{14} \in G$  by

$$(g_{ij})_v = \begin{cases} \exp(1/e_v) & \text{if } v = v_{ij}, \\ 1 & \text{otherwise.} \end{cases}$$

Then  $\delta(g_{13}), \delta(g_{23}), \delta(g_{14})$  is a basis for  $H$ , so we can define the section  $\sigma: H \rightarrow G$  by  $\delta(g_{ij}) \mapsto g_{ij}$ .

As before, let  $\epsilon_1, \dots, \epsilon_r$  be a  $\mathbb{Z}$ -basis for  $E/E_{\text{tor}}$ . Then the  $x_j = \epsilon_j \otimes 1$  form an  $\mathbb{R}$ -basis of  $E_{\mathbb{R}}$ . Extend this to a basis for  $G$  by adjoining the three vectors  $g_{ij}$ . It follows that

$$c\mu_G([x_1, \dots, x_r, g_{13}, g_{23}, g_{14}]) = \mu([x_1, \dots, x_r]) \cdot \mu_H([\delta(g_{13}), \delta(g_{23}), \delta(g_{14})]),$$

where  $[\dots]$  denotes convex hull. We have  $\mu([x_1, \dots, x_r]) = \text{Reg}_{K_1, K_2}(L)$  by the normalization of  $\mu$ . The convex hull of the  $\delta(g_{ij})$  is the “unit cube,” so it has volume 1. It is easily seen that  $\mu_G([x_1, \dots, x_r, g_{13}, g_{23}, g_{14}]) = 2^{-r_2(L)} \text{Reg}_{K_1, K_2}(L)$ . Hence  $c = 2^{r_2(L)}$ .  $\square$

For a fractional ideal  $\mathfrak{a}$  of  $L$  and  $t > 0$ , recall the theta series  $\Theta_E(t; \mathfrak{a})$  defined in (1.2).

We will use the Mellin transform to study this function. First, we define some notation.

For any  $z \in \mathbb{C}$  with  $\text{Re}(z) > 0$  and any  $\kappa \in [0, 1]$ , set

$$\alpha_{\kappa}(z) = \kappa \log \Gamma(z) + (1 - \kappa) \log \Gamma(z + \tfrac{1}{2}).$$

For  $i \in \{1, 2\}$  and  $j \in \{3, 4\}$ , let

$$k_{ij} = \frac{1}{m}(p_{w_{ij}} + q_{w_{ij}}). \tag{3.1}$$

Given  $s = (s_1, s_2, s_3) \in \mathbb{C}^3$ , define

$$s_{13} = s_1 + s_3, \quad s_{23} = s_2 + s_3, \quad s_{14} = s_1, \quad s_{24} = s_2.$$

Let  $\mathcal{R}$  denote the region

$$\mathcal{R} = \{s \in \mathbb{C}^3 \mid \text{all } \text{Re}(s_{ij}) > 0\}.$$

For a given  $\vec{\kappa} = (\kappa_{13}, \kappa_{14}, \kappa_{23}, \kappa_{24})$ , we define a function  $\alpha_{\vec{\kappa}}: \mathcal{R} \rightarrow \mathbb{C}$  by

$$\alpha_{\vec{\kappa}}(s) = \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \alpha_{\kappa_{ij}}(s_{ij}). \quad (3.2)$$

Let  $\alpha$  denote  $\alpha_{\vec{k}}$ , where  $\vec{k} = (k_{13}, k_{14}, k_{23}, k_{24})$  as in (3.1).

For  $g \in G$ , set

$$\|g\|^2 = \sum_{v \in \mathcal{A}_L} e_v g_v^2$$

and

$$\Psi(g) = \int_{E_{\mathbb{R}}} \exp(-\|gx\|^2) d\mu(x).$$

We want to evaluate  $\Psi(g)$ ; since  $\Psi(g)$  depends only on  $g$  modulo  $E_{\mathbb{R}}$ , it suffices to consider

$\psi = \Psi \circ \sigma$ . Now we compute the Mellin transform of  $\psi$ :

$$\begin{aligned} (M\psi)(s) &= (M\psi)(s_1, s_2, s_3) = \int_H \Psi(\sigma(h)) h^s d\mu_H(h) \\ &= \int_H \left( \int_{E_{\mathbb{R}}} \exp(-\|x\sigma(h)\|^2) d\mu(x) \right) h^s d\mu_H(h) \\ &= \int_{E_{\mathbb{R}} \times H} \exp(-\|\phi(x, h)\|^2) \delta(\phi(x, h))^s (d\mu \times d\mu_H)(x, h) \\ &= 2^{r_2(L)} \int_G \exp(-\|g\|^2) \delta(g)^s d\mu_G(g). \end{aligned}$$

Next observe that

$$\delta(g)^s = \left( \prod_{v|w_1} g_v^{e_v} \right)^{s_1} \left( \prod_{v|w_2} g_v^{e_v} \right)^{s_2} \left( \prod_{v|w_3} g_v^{e_v} \right)^{s_3} = \prod_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \prod_{v|w_{ij}} g_v^{e_v s_{ij}}.$$

It follows that

$$\begin{aligned}
(M\psi)(s) &= 2^{r_2(L)} \int_G \exp(-\|g\|^2) \prod_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \prod_{v|w_{ij}} g_v^{e_v s_{ij}} d\mu_G(g) \\
&= 2^{r_2(L)} \prod_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \prod_{v|w_{ij}} \int_G e^{-e_v g_v^2} g_v^{e_v s_{ij}} \frac{dg_v}{g_v} \\
&= 2^{r_2(L) - |\mathcal{A}_L|} \prod_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \Gamma\left(\frac{s_{ij}}{2}\right)^{p_{w_{ij}}} (2^{-s_{ij}} \Gamma(s_{ij}))^{q_{w_{ij}}} \\
&= 2^{-|\mathcal{A}_L|} \pi^{-r_2(L)/2} \prod_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \Gamma\left(\frac{s_{ij}}{2}\right)^{p_{w_{ij}} + q_{w_{ij}}} \Gamma\left(\frac{s_{ij} + 1}{2}\right)^{q_{w_{ij}}} \\
&= 2^{-|\mathcal{A}_L|} \pi^{-r_2(L)/2} \exp(m\alpha(\tfrac{1}{2}s)),
\end{aligned}$$

where we have used the identity  $2^{-s}\Gamma(s) = (2\sqrt{\pi})^{-1}\Gamma(\frac{s}{2})\Gamma(\frac{s+1}{2})$ .

Setting  $h = \delta(g)$  and taking an inverse Mellin transform,

$$\begin{aligned}
\Psi(g) &= \psi(h) \\
&= \frac{2^{-|\mathcal{A}_L|} \pi^{-r_2(L)/2}}{(2\pi i)^3} \int_{c-i\infty}^{c+i\infty} \int_{c-i\infty}^{c+i\infty} \int_{c-i\infty}^{c+i\infty} h_{w_1}^{-s_1} h_{w_2}^{-s_2} h_{w_3}^{-s_3} \exp(m\alpha(\tfrac{1}{2}s)) ds_1 ds_2 ds_3 \\
&= \frac{2^{3-|\mathcal{A}_L|} \pi^{-r_2(L)/2}}{(2\pi i)^3} \int_{c-i\infty}^{c+i\infty} \int_{c-i\infty}^{c+i\infty} \int_{c-i\infty}^{c+i\infty} h_{w_1}^{-2s_1} h_{w_2}^{-2s_2} h_{w_3}^{-2s_3} \exp(m\alpha(s)) ds_1 ds_2 ds_3.
\end{aligned}$$

Given  $a \in L^*$  and  $t > 0$ , define  $g \in G$  by  $g_v = \sqrt{t}|a|_v$ . Then for any  $x \in E_{\mathbb{R}}$ ,  $t\|ax\|^2 = \|gx\|^2$ ,

so

$$\Psi(g) = \int_{E_{\mathbb{R}}} \exp(-t\|ax\|^2) d\mu(x)$$

is the integral that appears in  $\Theta_E$ . For  $w \in \mathcal{A}_{K_i}$ ,

$$h_w = \prod_{v|w} g_v^{e_v} = \prod_{v|w} t^{e_v/2} |a|_v^{e_v} = t^m |\mathbf{N}_{L/K_i}(a)|_w.$$

Let  $A = 2^{3-|\mathcal{A}_L|} \pi^{-r_2(L)/2}$  and let  $a_w = \frac{1}{m} \log |\mathbf{N}_{L/K_i}(a)|_w$ . For any  $y \in \mathbb{R}$ , define

$$g_{y,a}(s) = -2(a_{w_1} + y)s_1 - 2(a_{w_2} + y)s_2 - 2(a_{w_3} + y)s_3 + \alpha(s)$$

and

$$f(y, a) = \frac{1}{(2\pi i)^3} \int_{c-i\infty}^{c+i\infty} \int_{c-i\infty}^{c+i\infty} \int_{c-i\infty}^{c+i\infty} \exp(mg_{y,a}(s)) ds_1 ds_2 ds_3.$$

The preceding work shows that  $\Psi(g) = Af(\log t, a)$ . Hence

$$\Theta_E(t; \mathfrak{a}) = \frac{\mu(E_{\mathbb{R}}/E)}{|E_{\text{tor}}|} + A \sum_{\substack{a \in \mathfrak{a}/E \\ a \neq 0}} f(\log(c_{\mathfrak{a}}t), a).$$

Inequality (1.3) says that  $\Theta_E(t; \mathfrak{a}) + \frac{1}{2m}t\Theta'_E(t; \mathfrak{a}) \geq 0$ . Plugging in the above formula, and choosing  $t$  so that  $y = \log(c_{\mathfrak{a}}t)$ , this proves that for any  $y \in \mathbb{R}$ ,

$$\frac{\mu(E_{\mathbb{R}}/E)}{|E_{\text{tor}}|} \geq A \sum_{\substack{a \in \mathfrak{a}/E \\ a \neq 0}} \left( -1 - \frac{1}{2m} \frac{f'}{f}(y, a) \right) f(y, a). \quad (3.3)$$

Next we want to choose  $y$  such that  $-\frac{f'}{f}(y, a) \geq 2m$  for all  $a$ . Then we can drop terms for  $a \neq 1$  to conclude that

$$\frac{\mu(E_{\mathbb{R}}/E)}{|E_{\text{tor}}|} \geq A \left( -1 - \frac{1}{2m} \frac{f'}{f}(y, 1) \right) f(y, 1). \quad (3.4)$$

This is done by the saddle-point method. In order to apply the saddle-point method, we first need to know that there is a saddle point. That is, we would like to find a point  $(s_1, s_2, s_3)$  where

$$\frac{\partial g_{y,a}}{\partial s_1} = \frac{\partial g_{y,a}}{\partial s_2} = \frac{\partial g_{y,a}}{\partial s_3} = 0.$$

This means we need to solve

$$a_{w_1} + y = \frac{1}{2}\alpha'_{k_{13}}(s_1 + s_3) + \frac{1}{2}\alpha'_{k_{14}}(s_1),$$

$$a_{w_2} + y = \frac{1}{2}\alpha'_{k_{23}}(s_2 + s_3) + \frac{1}{2}\alpha'_{k_{24}}(s_2),$$

$$a_{w_3} + y = \frac{1}{2}\alpha'_{k_{13}}(s_1 + s_3) + \frac{1}{2}\alpha'_{k_{23}}(s_2 + s_3).$$

Note that for any  $k \in (0, 1]$ ,  $\alpha'_k : (0, \infty) \rightarrow \mathbb{R}$  is strictly increasing and surjective.



**Lemma 3.2.** *This system has a unique solution in  $\mathcal{R} \cap \mathbb{R}^3$ .*

*Proof.* The given system of equations is equivalent to

$$\frac{1}{2}\alpha'_{k_{24}}(s_2) = a_{w_1} + a_{w_2} - a_{w_3} + y - \frac{1}{2}\alpha'_{k_{14}}(s_1), \quad (3.5)$$

$$\frac{1}{2}\alpha'_{k_{23}}(s_2 + s_3) = a_{w_3} - a_{w_1} + \frac{1}{2}\alpha'_{k_{14}}(s_1), \quad (3.6)$$

$$\frac{1}{2}\alpha'_{k_{13}}(s_1 + s_3) = a_{w_1} + y - \frac{1}{2}\alpha'_{k_{14}}(s_1). \quad (3.7)$$

Note that equation (3.5) determines  $s_2 > 0$  as a strictly decreasing function of  $s_1 > 0$ , and equation (3.6) determines  $s_2 + s_3 > 0$  as a strictly increasing function of  $s_1 > 0$ . Therefore these two equations determine  $s_3$  as a strictly increasing function of  $s_1$ . Under this correspondence,  $s_3 \rightarrow \infty$  as  $s_1 \rightarrow \infty$ . On the other hand, equation (3.7) determines  $s_3$  as a strictly decreasing function of  $s_1$ . Under this correspondence,  $s_3 \rightarrow \infty$  as  $s_1 \rightarrow 0$ . Now we have two functions  $s_1 \mapsto s_3$ , and the solutions of the system correspond to choices of  $s_1$  at which these functions are equal. It follows from what we have said that there is exactly one solution. □

## Chapter 4

# Single-variable calculus

Before proceeding to the triple-integral estimates we need, we record some single-variable lemmas which will be useful. Throughout this chapter, we assume  $\frac{1}{2} \leq \kappa \leq 1$ ,  $m > 0$ , and  $\sigma > 0$  (sometimes adding an additional assumption on  $m$  where helpful). Recall the formula [1], 6.4.10: for  $n \geq 1$  and  $\operatorname{Re}(s) > 0$ ,

$$\frac{\Psi^{(n)}(s)}{n!} = (-1)^{n+1} \sum_{k=0}^{\infty} \frac{1}{(s+k)^{n+1}}, \quad \Psi = \frac{\Gamma'}{\Gamma}. \quad (4.1)$$

**Lemma 4.1.** *If  $m\kappa \geq 4$ , then*

$$\frac{\sqrt{\sigma^2 \alpha''_{\kappa}(\sigma)}}{1.25^{m\kappa[\sigma]/2}} < \sqrt{2} \quad \text{and} \quad \frac{\sigma^2 \alpha''_{\kappa}(\sigma)}{1.25^{m\kappa[\sigma]/2}} < 2.$$

*If  $m\kappa \geq 30$ , then*

$$\frac{\sqrt{\sigma^2 \alpha''_{\kappa}(\sigma)}}{(\frac{73}{72})^{m\kappa[\sigma]/2}} < \sqrt{2}.$$

*Proof.* The first inequality is given in the proof of Friedman and Skoruppa's Lemma 5.5; as the other inequalities are proven in the same way, the proof is repeated here. Note that

$$\sigma^2 \alpha''_{\kappa}(\sigma) \leq \sigma^2 \Psi'(\sigma) < 1 + \sigma,$$

where the last inequality follows from estimating the sum (4.1) by an integral. Thus we need to show that  $\sqrt{1+\sigma}/1.25^{m\kappa[\sigma]/2} < \sqrt{2}$ . We see that  $\sqrt{1+\sigma}/1.25^{m\kappa[\sigma]/2}$  is maximized as  $\sigma \rightarrow 1^-$ , because  $1.25^2 > \sqrt{3/2}$ .  $\square$

**Lemma 4.2.** *Let  $\epsilon$  be 0 or 1. Suppose  $u > 0$  and  $m\kappa > 2$ . Then*

$$\int_{u\sigma}^{\infty} |t^{\epsilon} e^{m\alpha(\sigma+it)}| dt \leq \frac{e^{m\alpha(\sigma)} \sigma^{1+\epsilon} u^{\epsilon-1} (1+u^2)}{(m\kappa-2)(1+\frac{u^2}{4})^{m\kappa[\sigma]/2} (1+u^2)^{m\kappa/2}},$$

where  $[\sigma]$  denotes the greatest integer less than or equal to  $\sigma$ .

*Proof.* This is Friedman and Skoruppa's Lemma 5.3.  $\square$

**Lemma 4.3.** *Let  $D$  be given with  $0 < D \leq m^{1/3}\sqrt{\kappa}$  and assume that  $m\kappa > 2$ . Define  $\delta = D/(m^{1/3}\sqrt{\alpha''_{\kappa}(\sigma)})$ . Then*

$$\begin{aligned} \int_{|t|>\delta} |e^{m\alpha_{\kappa}(\sigma+it)}| dt &\leq \frac{\sqrt{2\pi}e^{m\alpha_{\kappa}(\sigma)}}{\sqrt{m\alpha''_{\kappa}(\sigma)}} \left( \frac{2^{3/2}\sigma\sqrt{m\alpha''_{\kappa}(\sigma)}}{\sqrt{\pi}(m\kappa-2)1.25^{m\kappa[\sigma]/2}2^{m\kappa/2}} + \frac{2^{3/2}\exp(-m^{1/3}D^2/4)}{\sqrt{\pi}m^{1/6}D} \right), \\ \int_{|t|>\delta} e^{-mt^2\alpha''_{\kappa}(\sigma)/2} dt &\leq \frac{2e^{-m^{1/3}D^2/2}}{m^{2/3}D\sqrt{\alpha''_{\kappa}(\sigma)}}. \end{aligned}$$

where  $[\sigma]$  denotes the greatest integer less than or equal to  $\sigma$ .

*Proof.* These inequalities can be found in Friedman and Skoruppa (see their proof of Lemma 5.4).  $\square$

**Lemma 4.4.** *Suppose  $m\kappa > 2$ . Then for any  $0 < D < m^{1/3}\sqrt{\kappa}$ ,*

$$\begin{aligned} \int_{-\infty}^{\infty} |e^{m\alpha_{\kappa}(\sigma+it)}| dt &< \frac{\sqrt{2\pi}e^{m\alpha_{\kappa}(\sigma)}}{\sqrt{m\alpha''_{\kappa}(\sigma)}} \left( 1 + \frac{2^{3/2}\sigma\sqrt{m\alpha''_{\kappa}(\sigma)}}{\sqrt{\pi}(m\kappa-2)1.25^{m\kappa[\sigma]/2}2^{m\kappa/2}} + \right. \\ &\quad \left. \frac{2^{3/2}\exp(-m^{1/3}D^2/4)}{\sqrt{\pi}m^{1/6}D} + \frac{e^{D^4/(4m^{1/3}\kappa)} - 1}{D^4/(4m^{1/3}\kappa)} \frac{3}{4m\kappa} \right), \end{aligned}$$

where  $[\sigma]$  again denotes the greatest integer less than or equal to  $\sigma$ . If  $m \geq 1000$ , then

$$\int_{-\infty}^{\infty} |e^{m\alpha_{\kappa}(\sigma+it)}| dt < \frac{\sqrt{2\pi}e^{m\alpha_{\kappa}(\sigma)}}{\sqrt{m\alpha''_{\kappa}(\sigma)}} \cdot 1.00205.$$

*Proof.* The first claim essentially comes from Lemma 5.4 of Friedman and Skoruppa, which estimates

$$\int_{-\infty}^{\infty} e^{m(\alpha_{\kappa}(\sigma+it)-iyt)} dt$$

for  $y = \alpha'_{\kappa}(\sigma)$ . However, that lemma has two extra terms which are not needed here. Friedman and Skoruppa bound the integral over  $|t| \geq \delta := D/(m^{1/3}\sqrt{\alpha''_{\kappa}(\sigma)})$  by replacing  $e^{m(\alpha_{\kappa}(\sigma+it)-iyt)}$  with  $|e^{m\alpha_{\kappa}(\sigma+it)}|$ , so that part of the argument works in this case without any change. That is,

$$\int_{|t| \geq \delta} |e^{m\alpha_{\kappa}(\sigma+it)}| dt \leq \frac{\sqrt{2\pi}e^{m\alpha_{\kappa}(\sigma)}}{\sqrt{m\alpha''_{\kappa}(\sigma)}} \left( \frac{2^{3/2}\sigma\sqrt{m\alpha''_{\kappa}(\sigma)}}{\sqrt{\pi}(m\kappa-2)1.25^{m\kappa[\sigma]/2}2^{m\kappa/2}} + \frac{2^{3/2}\exp(-m^{1/3}D^2/4)}{\sqrt{\pi}m^{1/6}D} \right).$$

Next, we have

$$\begin{aligned} \int_{-\delta}^{\delta} |e^{m\alpha_{\kappa}(\sigma+it)-m\alpha_{\kappa}(\sigma)}| dt &= \int_{-\delta}^{\delta} e^{-\frac{1}{2}m\alpha''_{\kappa}(\sigma)t^2} dt + \int_{-\delta}^{\delta} (|e^{m\alpha_{\kappa}(\sigma+it)-m\alpha_{\kappa}(\sigma)}| - e^{-\frac{1}{2}m\alpha''_{\kappa}(\sigma)t^2}) dt \\ &< \frac{\sqrt{2\pi}}{\sqrt{m\alpha''_{\kappa}(\sigma)}} + \int_{-\delta}^{\delta} (|e^{m\alpha_{\kappa}(\sigma+it)-m\alpha_{\kappa}(\sigma)}| - e^{-\frac{1}{2}m\alpha''_{\kappa}(\sigma)t^2}) dt. \end{aligned}$$

Note that

$$|e^{m\alpha_{\kappa}(\sigma+it)-m\alpha_{\kappa}(\sigma)}| - e^{-\frac{1}{2}m\alpha''_{\kappa}(\sigma)t^2} = (e^{m\operatorname{Re}(\rho(t))} - 1)e^{-\frac{1}{2}m\alpha''_{\kappa}(\sigma)t^2},$$

where  $\rho(t) = \alpha_{\kappa}(\sigma+it) - iyt + \frac{1}{2}\alpha''_{\kappa}(\sigma)t^2$ . Therefore we can bound the last integral in the same way that Friedman and Skoruppa bounded the integral of  $(e^{m\rho(t)} - 1)e^{-\frac{1}{2}m\alpha''_{\kappa}(\sigma)t^2}$ , except that we do not get a term coming from  $\operatorname{Im}(\rho)$ . We conclude that

$$\int_{-\delta}^{\delta} (|e^{m\alpha_{\kappa}(\sigma+it)-m\alpha_{\kappa}(\sigma)}| - e^{-\frac{1}{2}m\alpha''_{\kappa}(\sigma)t^2}) dt < \frac{\sqrt{2\pi}}{\sqrt{m\alpha''_{\kappa}(\sigma)}} \frac{e^{D^4/(4m^{1/3}\kappa)} - 1}{D^4/(4m^{1/3}\kappa)} \frac{3}{4m\kappa}.$$

Now suppose  $m \geq 1000$ . Then

$$\frac{2^{3/2}\sigma\sqrt{m\alpha''_{\kappa}(\sigma)}}{\sqrt{\pi}(m\kappa-2)1.25^{m\kappa[\sigma]/2}2^{m\kappa/2}} = \frac{4}{m\kappa^{3/2}\sqrt{\pi}} \frac{\sqrt{m\kappa/2}}{2^{m\kappa/2}} \frac{m\kappa}{m\kappa-2} \frac{\sqrt{\sigma^2\alpha''_{\kappa}(\sigma)}}{1.25^{m\kappa[\sigma]/2}} < \frac{10^{-76}}{m}; \quad (4.2)$$

bounds for the first three terms are obvious, and the last term is addressed by Lemma 4.1.

Thus

$$\int_{-\infty}^{\infty} |e^{m\alpha_{\kappa}(\sigma+it)}| dt < \frac{\sqrt{2\pi}e^{m\alpha_{\kappa}(\sigma)}}{\sqrt{m\alpha_{\kappa}''(\sigma)}} \left( 1 + \frac{10^{-76}}{m} + \frac{2^{3/2} \exp(-m^{1/3}D^2/4)}{\sqrt{\pi}m^{1/6}D} + \frac{e^{D^4/(2m^{1/3})} - 1}{D^4/(2m^{1/3})} \frac{3}{2m} \right).$$

Set  $D = 1.76$ . The quantity in parentheses is decreasing in  $m$  for  $m > 0$ , so the claimed bound follows by plugging in  $m = 1000$ .  $\square$

**Lemma 4.5.** *Suppose  $m \geq 1000$ . Let  $0 < D \leq m^{1/3}\sqrt{\kappa}$  be given, and again set  $\delta = D/(m^{1/3}\sqrt{\alpha_{\kappa}''(\sigma)})$ . Then*

$$\int_{|t|>\delta} |e^{m\alpha_{\kappa}(\sigma+it)}| dt < \frac{\sqrt{2\pi}e^{m\alpha_{\kappa}(\sigma)}}{\sqrt{m\alpha_{\kappa}''(\sigma)}} \frac{1}{m} \cdot \left( 10^{-76} + \frac{2^{3/2}m^{5/6} \exp(-m^{1/3}D^2/4)}{\sqrt{\pi}D} \right).$$

*Proof.* This is immediate from Lemma 4.3 and inequality (4.2).  $\square$

**Lemma 4.6.** *Let  $C > 0$  be given. Then*

$$\int_{-\infty}^{\infty} e^{-Ct^2} dt = \sqrt{\pi}C^{-1/2}, \quad \int_0^{\infty} e^{-Ct^2} t dt = \frac{1}{2}C^{-1}, \quad \int_0^{\infty} e^{-Ct^2} t^3 dt = \frac{1}{2}C^{-2}.$$

*Proof.* Make the substitution  $u = Ct^2$ . For any  $n \geq 0$ , we have

$$\int_0^{\infty} e^{-Ct^2} t^n dt = \int_0^{\infty} e^{-u} (u/C)^{(n+1)/2} \frac{du}{2u} = \frac{1}{2} \Gamma\left(\frac{n+1}{2}\right) C^{-(n+1)/2}. \quad \square$$

**Lemma 4.7.** *Let any  $\sigma > 0$  and  $0 \leq \kappa \leq 1$  be given. Define  $\rho = \rho_{\kappa,\sigma}: \mathbb{R} \rightarrow \mathbb{C}$  by*

$$\rho(t) = \alpha_{\kappa}(\sigma + it) - \alpha_{\kappa}(\sigma) - i\alpha_{\kappa}'(\sigma)t + \frac{1}{2}\alpha_{\kappa}''(\sigma)t^2;$$

*i.e.,  $\rho$  is the error in the degree-2 Taylor approximation to  $\alpha_{\kappa}(\sigma + it)$ . Then for any  $t \in \mathbb{R}$ ,*

$$|\operatorname{Im}(\rho(t))| \leq -\frac{\alpha_{\kappa}^{(3)}(\sigma)}{3!}|t|^3, \quad |\operatorname{Re}(\rho(t))| \leq \frac{\alpha_{\kappa}^{(4)}(\sigma)}{4!}t^4. \quad (4.3)$$

*If  $|t| \leq \sigma$ , then*

$$0 \leq \operatorname{Re}(\rho(t)) \leq \frac{\alpha_{\kappa}''(\sigma)}{4}t^2; \quad (4.4)$$

if  $|t| \leq \frac{\sigma}{3\sqrt{2}}$ , then

$$0 \leq \operatorname{Re}(\rho(t)) \leq \frac{\alpha''_{\kappa}(\sigma)}{72} t^2.$$

Furthermore,  $\operatorname{Im}(\alpha_{\kappa}(\sigma + it))$  is odd and  $\operatorname{Re}(\alpha_{\kappa}(\sigma + it))$  is even as a function of  $t$ . Thus  $\operatorname{Im}(\rho(t))$  is an odd function and  $\operatorname{Re}(\rho(t))$  is an even function.

*Proof.* The odd/even statement is proven by Friedman and Skoruppa, as well as the fact that  $\operatorname{Re}(\rho(t)) \geq 0$  for  $|t| \leq \sigma$ . See the proof of their Lemma 5.1.

From (4.1), we know that  $|\alpha_{\kappa}^{(3)}(\sigma + it)|$  and  $|\alpha_{\kappa}^{(4)}(\sigma + it)|$  (considered as functions of  $t$ ) are both maximized at  $t = 0$ , with  $\alpha_{\kappa}^{(3)}(\sigma) < 0$  and  $\alpha_{\kappa}^{(4)}(\sigma) > 0$ . Now apply the Taylor remainder theorem to  $\operatorname{Im}(\alpha_{\kappa}(\sigma + it))$ : since

$$\frac{d^3}{dt^3} \operatorname{Im}(\alpha_{\kappa}(\sigma + it)) = -i \operatorname{Im}(\alpha_{\kappa}^{(3)}(\sigma + it)),$$

we see that for any  $t \in \mathbb{R}$ , there exists  $\theta_t$  between 0 and  $t$  such that

$$|\operatorname{Im}(\alpha_{\kappa}(\sigma + it))| = \left| \frac{-i \operatorname{Im}(\alpha_{\kappa}^{(3)}(\sigma + i\theta_t))}{3!} t^3 \right| \leq -\frac{\alpha_{\kappa}^{(3)}(\sigma)}{3!} |t|^3.$$

This proves (4.3) for the imaginary part; the proof for the real part is identical.

For any  $\sigma > 0$ , (4.1) shows that

$$\frac{\sigma^2 \Psi^{(3)}(\sigma)}{4!} = \frac{1}{4} \sum_{k=0}^{\infty} \frac{\sigma^2}{(\sigma + k)^4} < \frac{1}{4} \sum_{k=0}^{\infty} \frac{1}{(\sigma + k)^2} = \frac{\Psi'(\sigma)}{4}.$$

It follows that  $\sigma^2 \alpha_{\kappa}^{(4)}(\sigma)/4! < \alpha''_{\kappa}(\sigma)/4$ . Thus for  $|t| \leq \sigma$ ,

$$|\operatorname{Re}(\rho(t))| \leq \frac{\alpha_{\kappa}^{(4)}(\sigma)}{4!} t^4 \leq \frac{\sigma^2 \alpha_{\kappa}^{(4)}(\sigma)}{4!} t^2 \leq \frac{\alpha''_{\kappa}(\sigma)}{4} t^2.$$

The same argument works for  $|t| \leq \frac{\sigma}{3\sqrt{2}}$ . □

**Lemma 4.8.** *Let  $R > 0$  be given. Then for any  $0 \leq u \leq R$  and any  $v \in \mathbb{R}$ , we have*

$$|\operatorname{Re}(e^{u+iv} - 1)| \leq u \frac{e^R - 1}{R} + \frac{v^2}{2}.$$

*Proof.* This is inequality (5.11) from Friedman and Skoruppa.  $\square$

**Lemma 4.9.** For  $m \geq 1000$ ,

$$\int_{|t| > \frac{\sigma}{3\sqrt{2}}} |te^{m\alpha_\kappa(\sigma+it)}| dt < \frac{0.0002557\sigma e^{m\alpha_\kappa(\sigma)}}{m^{3/2}\sqrt{\alpha''_\kappa(\sigma)}}.$$

*Proof.* By Lemma 4.2,

$$\begin{aligned} \int_{\frac{\sigma}{3\sqrt{2}}}^{\infty} |te^{m\alpha_\kappa(\sigma+it)}| dt &\leq 2 \cdot \frac{\frac{19}{18}\sigma^2 e^{m\alpha_\kappa(\sigma)}}{(m\kappa - 2)\left(\frac{73}{72}\right)^{m\kappa[\sigma]/2}\left(\frac{19}{18}\right)^{m\kappa/2}} \\ &= \frac{19}{9(\kappa - \frac{2}{m})} \cdot \frac{\sqrt{m}}{\left(\frac{19}{18}\right)^{m\kappa/2}} \cdot \frac{\sqrt{\sigma^2\alpha''_\kappa(\sigma)}}{\left(\frac{73}{72}\right)^{m\kappa[\sigma]/2}} \cdot \frac{\sigma e^{m\alpha_\kappa(\sigma)}}{m^{3/2}\sqrt{\alpha''_\kappa(\sigma)}}. \end{aligned}$$

We have  $\sqrt{m}/(19/18)^{m\kappa/2} \leq \sqrt{m}/(19/18)^{m/4} \leq \sqrt{1000}/(19/18)^{1000/4} < 0.0000853$ . Combining this with Lemma 4.1, we conclude that

$$\begin{aligned} \int_{\frac{\sigma}{3\sqrt{2}}}^{\infty} |te^{m\alpha_\kappa(\sigma+it)}| dt &< \frac{19}{9(\frac{1}{2} - \frac{2}{1000})} \cdot 0.0000853 \cdot \sqrt{2} \cdot \frac{\sigma e^{m\alpha_\kappa(\sigma)}}{m^{3/2}\sqrt{\alpha''_\kappa(\sigma)}} \\ &< \frac{0.0002557\sigma e^{m\alpha_\kappa(\sigma)}}{m^{3/2}\sqrt{\alpha''_\kappa(\sigma)}}. \end{aligned} \quad \square$$

**Lemma 4.10.** For  $m \geq 1000$ ,

$$\int_{|t| > \frac{\sigma}{3\sqrt{2}}} |e^{m\alpha_\kappa(\sigma+it)}| dt < \frac{0.00003429e^{m\alpha_\kappa(\sigma)}}{m\sqrt{\alpha''_\kappa(\sigma)}}.$$

*Proof.* The argument is the same as the proof of Lemma 4.9; by Lemma 4.2,

$$\begin{aligned} e^{-m\alpha_\kappa(\sigma)} \int_{\frac{\sigma}{3\sqrt{2}}}^{\infty} |e^{m\alpha_\kappa(\sigma+it)}| dt &\leq 2 \cdot \frac{\frac{19\sqrt{2}}{6}\sigma}{(m\kappa - 2)\left(\frac{73}{72}\right)^{m\kappa[\sigma]/2}\left(\frac{19}{18}\right)^{m\kappa/2}} \\ &= \frac{19\sqrt{2}}{3(\kappa - \frac{2}{m})} \cdot \frac{1}{\left(\frac{19}{18}\right)^{m\kappa/2}} \cdot \frac{\sqrt{\sigma^2\alpha''_\kappa(\sigma)}}{\left(\frac{73}{72}\right)^{m\kappa[\sigma]/2}} \cdot \frac{1}{m\sqrt{\alpha''_\kappa(\sigma)}} \\ &< \frac{19\sqrt{2}}{3(\frac{1}{2} - \frac{2}{1000})} \cdot \frac{1}{\left(\frac{19}{18}\right)^{1000/4}} \cdot \sqrt{2} \cdot \frac{1}{m\sqrt{\alpha''_\kappa(\sigma)}} \\ &< \frac{0.00003429}{m\sqrt{\alpha''_\kappa(\sigma)}}. \end{aligned} \quad \square$$

**Lemma 4.11.** *For any  $m > 0$ ,*

$$\int_{-\frac{\sigma}{3\sqrt{2}}}^{\frac{\sigma}{3\sqrt{2}}} |te^{m\alpha_\kappa(\sigma+it)}| dt < \frac{\frac{72}{35}e^{m\alpha_\kappa(\sigma)}}{m\alpha''_\kappa(\sigma)}.$$

*Proof.* Lemma 4.7 shows that, for  $|t| \leq \sigma/(3\sqrt{2})$ ,

$$|e^{m\alpha_\kappa(\sigma+it)}| \leq e^{m\alpha_\kappa(\sigma) - m\alpha''_\kappa(\sigma)t^2/2 + m\alpha''_\kappa(\sigma)t^2/72} = e^{m\alpha_\kappa(\sigma) - (35/72)m\alpha''_\kappa(\sigma)t^2}.$$

Hence

$$\int_{-\frac{\sigma}{3\sqrt{2}}}^{\frac{\sigma}{3\sqrt{2}}} |te^{m\alpha_\kappa(\sigma+it)}| dt < 2 \int_0^\infty te^{m\alpha_\kappa(\sigma) - (35/72)m\alpha''_\kappa(\sigma)t^2} dt = \frac{\frac{72}{35}e^{m\alpha_\kappa(\sigma)}}{m\alpha''_\kappa(\sigma)}. \quad \square$$

**Lemma 4.12.** *For any  $\sigma > 0$ ,  $\alpha'_{1/2}(\sigma/\sqrt{2}) > \alpha'_1(\sigma) = \Psi'(\sigma)$ .*

*Proof.* The duplication formula for  $\Psi$  ([1], 6.3.8) says that

$$\alpha'_{1/2}(\sigma) = \frac{1}{2}\Psi(\sigma) + \frac{1}{2}\Psi(\sigma + \frac{1}{2}) = \Psi(2\sigma) - \log 2.$$

Differentiating,  $\alpha''_{1/2}(\sigma) = 2\Psi'(2\sigma)$ . Thus we want to prove that  $2\Psi'(\sqrt{2}\sigma) > \Psi'(\sigma)$ . Estimating the sum (4.1) by integrals, we find that

$$\frac{1}{\sigma^2} + \frac{1}{(\sigma+1)} < \Psi'(\sigma) < \frac{1}{\sigma^2} + \frac{1}{(\sigma+1)^2} + \frac{1}{\sigma+1}.$$

Hence

$$\begin{aligned} 2\Psi'(\sqrt{2}\sigma) - \Psi'(\sigma) &> 2 \left( \frac{1}{2\sigma^2} + \frac{1}{(\sqrt{2}\sigma+1)} \right) - \left( \frac{1}{\sigma^2} + \frac{1}{(\sigma+1)^2} + \frac{1}{\sigma+1} \right) \\ &= \frac{(2-\sqrt{2})\sigma^2 + (3-2\sqrt{2})\sigma}{(\sigma+1)^2(\sqrt{2}\sigma+1)} \\ &> 0. \end{aligned} \quad \square$$



## Chapter 5

# Estimation of $f$ and $f'/f$

Let  $m > 0$  be given. (We will primarily be interested in  $m \geq 1000$ , but we will note which lemmas hold for all  $m > 0$  and which require  $m \geq 1000$ .) Let  $\frac{1}{2} \leq k_{ij} \leq 1$  be given for  $i \in \{1, 2\}$  and  $j \in \{3, 4\}$ , and let  $\alpha = \alpha_{\vec{k}}$  as in (3.2). Also let  $\vec{y} = (y_1, y_2, y_3) \in \mathbb{R}^3$  be given. For  $s = (s_1, s_2, s_3) \in \mathcal{R}$ , define

$$g(s) = -\vec{y} \cdot s + \alpha(s)$$

and

$$G(s) = \exp(mg(s)).$$

For a given  $a \in L^*$  and  $y \in \mathbb{R}$ , if we take  $\vec{y} = (2(a_{w_1} + y), 2(a_{w_2} + y), 2(a_{w_3} + y))$ , then  $g = g_{y,a}$ . Let  $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$  be the unique point in  $\mathcal{R} \cap \mathbb{R}^3$  at which

$$\frac{\partial g}{\partial s_1} = \frac{\partial g}{\partial s_2} = \frac{\partial g}{\partial s_3} = 0;$$

$\vec{\sigma}$  exists by Lemma 3.2. Recall that we are interested in

$$f = \frac{1}{(2\pi i)^3} \int_{\sigma_1 - i\infty}^{\sigma_1 + i\infty} \int_{\sigma_2 - i\infty}^{\sigma_2 + i\infty} \int_{\sigma_3 - i\infty}^{\sigma_3 + i\infty} G(s) ds_3 ds_2 ds_1 = \frac{1}{(2\pi)^3} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1.$$

To simplify notation, define  $A_{ij} = \alpha''_{k_{ij}}(\sigma_{ij})$  and  $\Pi_{ij} = A_{13}A_{23}A_{14}A_{24}/A_{ij}$ . Let  $A_{\min} = \min(A_{ij})$ , and let  $\Pi_{\max} = \max(\Pi_{ij}) = A_{13}A_{23}A_{14}A_{24}/A_{\min}$ . (Without loss of generality, we will assume that  $A_{23} = A_{\min}$  and  $A_{24} \leq A_{13}$  wherever this helps.<sup>1</sup>) Define  $P = \Pi_{13} + \Pi_{23} + \Pi_{14} + \Pi_{24}$ ; i.e.,  $P = P_3(A_{13}, A_{14}, A_{23}, A_{24})$ , where  $P_3$  is the degree-3 elementary symmetric polynomial.

Define

$$\begin{aligned} H(\vec{t}) &:= \exp\left(m\left(g(\vec{\sigma}) - \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} A_{ij} t_{ij}^2/2\right)\right) \\ &= G(\vec{\sigma}) \exp\left(-m \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} A_{ij} t_{ij}^2/2\right). \end{aligned}$$

The idea is that  $H(\vec{t})$  is a good approximation to  $G(\vec{\sigma} + i\vec{t})$ ; we obtain  $H(\vec{t})$  from  $G(s) = \exp(mg(s))$  by replacing each  $\alpha_{k_{ij}}(s_{ij})$  in  $g(s) = -\vec{y} \cdot s + \sum \alpha_{k_{ij}}(s_{ij})$  with its degree-two Taylor approximation (as a function of  $t_{ij}$ ). The fact that  $\vec{\sigma}$  is a critical point ensures that the linear terms cancel. The main term in our estimate for  $f$  comes from integrating  $H$ :

**Lemma 5.1.** *We have*

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) dt_3 dt_2 dt_1 = \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}}.$$

*Proof.* It is well-known (see, e.g., [7], page 71) that if  $A = (a_{ij})$  is an  $n \times n$  positive-definite symmetric matrix, then

$$\int_{T \in \mathbb{R}^n} \exp(-\frac{1}{2} \sum a_{ij} T_i T_j) dT = (2\pi)^{n/2} \det(A)^{-1/2}.$$

---

<sup>1</sup>Note that the only relation among  $t_{13}$ ,  $t_{14}$ ,  $t_{23}$ , and  $t_{24}$  is  $t_{13} + t_{24} = t_{14} + t_{23}$ . Since this is symmetric, we are free to choose any  $A_{ij}$  as  $A_{\min}$ . After choosing  $A_{23} = A_{\min}$ , we are still free to swap  $A_{13}$  and  $A_{24}$  if necessary.

In this case,

$$\det(A) = m^3 \det \begin{pmatrix} A_{13} + A_{14} & 0 & A_{13} \\ 0 & A_{23} + A_{24} & A_{23} \\ A_{13} & A_{23} & A_{13} + A_{23} \end{pmatrix} = m^3 P. \quad \square$$

As a simple consequence of this lemma, we can evaluate some other integrals which will be useful later:

**Lemma 5.2.** *Let any  $i \in \{1, 2\}$  and  $j \in \{3, 4\}$  be given. Then*

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) t_{ij}^4 dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} \frac{3}{m^2 A_{ij}^2}$$

and

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) t_{ij}^6 dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} \frac{15}{m^3 A_{ij}^3}.$$

*Proof.* First observe that

$$\begin{aligned} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) t_{ij}^2 dt_3 dt_2 dt_1 &= -\frac{2}{m} \frac{\partial}{\partial A_{ij}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) dt_3 dt_2 dt_1 \\ &= -\frac{2}{m} \frac{\partial}{\partial A_{ij}} \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} \\ &= \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{5/2} P^{3/2}} \frac{\partial P}{\partial A_{ij}}. \end{aligned}$$

Note that  $\partial P / \partial A_{ij}$  does not depend on  $A_{ij}$ ; for example,  $\partial P / \partial A_{13} = A_{14} A_{23} + A_{14} A_{24} + A_{23} A_{24}$ . Hence

$$\begin{aligned} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) t_{ij}^4 dt_3 dt_2 dt_1 &= -\frac{2}{m} \frac{\partial}{\partial A_{ij}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) t_{ij}^2 dt_3 dt_2 dt_1 \\ &= -\frac{2}{m} \frac{\partial}{\partial A_{ij}} \left( \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{5/2} P^{3/2}} \frac{\partial P}{\partial A_{ij}} \right) \\ &= 3 \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{7/2} P^{5/2}} \left( \frac{\partial P}{\partial A_{ij}} \right)^2, \end{aligned}$$

so the first claim will follow once we check that

$$\frac{\partial P / \partial A_{ij}}{P} < \frac{1}{A_{ij}}. \quad (5.1)$$

This is obvious; for example,

$$A_{13} \frac{\partial P}{\partial A_{13}} = A_{13} A_{14} A_{23} + A_{13} A_{14} A_{24} + A_{13} A_{23} A_{24} < P.$$

The second claim is proven identically:

$$\begin{aligned} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) t_{ij}^6 dt_3 dt_2 dt_1 &= -\frac{2}{m} \frac{\partial}{\partial A_{ij}} \left( 3 \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{7/2} P^{5/2}} \left( \frac{\partial P}{\partial A_{ij}} \right)^2 \right) \\ &= 15 \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{9/2} P^{7/2}} \left( \frac{\partial P}{\partial A_{ij}} \right)^3 \\ &< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} \frac{15}{m^3 A_{ij}^3}. \quad \square \end{aligned}$$

As in Lemma 4.5, choose  $D$  such that  $0 < D \leq 1000^{1/3}/\sqrt{2}$ . (This ensures  $D \leq m^{1/3} \sqrt{k_{ij}}$  for all  $i, j$  when  $m \geq 1000$ .) Define

$$\delta_{ij} = \frac{D}{m^{1/3} \sqrt{A_{ij}}}.$$

Let  $\Delta \subseteq \mathbb{R}^3$  denote the set

$$\{(t_1, t_2, t_3) \in \mathbb{R}^3 \mid |t_{14}| \leq \delta_{14}, |t_{24}| \leq \delta_{24}, |t_{13}| \leq \delta_{13}\}.$$

Recall that we want to prove that the integrals of  $G$  and  $H$  have the same asymptotic behavior. We will do so by showing that  $H$  is a good approximation to  $G$  inside of  $\Delta$ , and that the contributions to the integrals outside of  $\Delta$  are (asymptotically) negligible.

**Lemma 5.3.** *Suppose  $m \geq 1000$ . Then*

$$\iiint_{\mathbb{R}^3 \setminus \Delta} |G(\vec{\sigma} + i\vec{t})| dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\Pi_{23}}} \frac{1}{m} \cdot 3.013 \left( 10^{-76} + \frac{2^{3/2} m^{5/6} \exp(-m^{1/3} D^2/4)}{\sqrt{\pi} D} \right).$$

*Proof.* Recall that

$$|\Gamma(\sigma + it)| \leq \Gamma(\sigma) \quad \text{for any } \sigma > 0 \text{ and } t \in \mathbb{R}. \quad (5.2)$$

Hence for any  $s \in \mathcal{R}$  with  $\operatorname{Re}(s_{23}) = \sigma_{23}$ , we have

$$|G(s)| \leq e^{m\alpha_{k_{23}}(\sigma_{23})} |\exp(m(-\vec{y} \cdot s + \alpha_{k_{13}}(s_{13}) + \alpha_{k_{14}}(s_{14}) + \alpha_{k_{24}}(s_{24})))|. \quad (5.3)$$

We can bound the triple integral of the right-hand side by splitting it into three single integrals. (We will use this strategy several more times in this chapter.) In order to do so, we will need to change variables from  $(t_1, t_2, t_3)$  to  $(t_{14}, t_{24}, t_{13})$ , so that the right-hand side of (5.3) becomes a product of three single-variable functions. The change-of-variable matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$$

has determinant 1, so the substitution does not introduce a Jacobian factor. Using Lemmas 4.4 and 4.5 to bound the resulting single integrals, we find that

$$\begin{aligned} \int_{|t_{14}| > \delta_{14}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |G(\vec{\sigma} + i\vec{t})| dt_3 dt_2 dt_1 \\ < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\Pi_{23}}} \frac{1}{m} \cdot 1.00205^2 \left( 10^{-76} + \frac{2^{3/2} m^{5/6} \exp(-m^{1/3} D^2/4)}{\sqrt{\pi} D} \right). \end{aligned}$$

We get the same bound for  $\int_{-\infty}^{\infty} \int_{|t_{24}| > \delta_{24}} \int_{-\infty}^{\infty}$  and for  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{|t_{13}| > \delta_{13}}$ . Note that  $3 \cdot 1.00205^2 < 3.013$ . □

**Lemma 5.4.** *We have*

$$\iiint_{\mathbb{R}^3 \setminus \Delta} H(\vec{t}) dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\Pi_{23}}} \frac{1}{m} \cdot 3 \sqrt{\frac{2}{\pi}} \frac{m^{5/6} e^{-m^{1/3} D^2/2}}{D}.$$

*Proof.* Arguing as in the proof of Lemma 5.3, Lemmas 4.3 and 4.6 show that

$$\begin{aligned} \int_{|t_{14}| > \delta_{14}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) dt_3 dt_2 dt_1 &\leq G(\vec{\sigma}) \left( \frac{2e^{-m^{1/3}D^2/2}}{m^{2/3}D\sqrt{A_{14}}} \right) \sqrt{\frac{2\pi}{mA_{24}}} \sqrt{\frac{2\pi}{mA_{13}}} \\ &= \frac{(2\pi)^{3/2}G(\vec{\sigma})}{m^{3/2}\sqrt{\Pi_{23}}} \frac{1}{m} \cdot \sqrt{\frac{2}{\pi}} \frac{m^{5/6}e^{-m^{1/3}D^2/2}}{D}. \end{aligned}$$

We get the same bound for  $\int_{-\infty}^{\infty} \int_{|t_{24}| > \delta_{24}} \int_{-\infty}^{\infty}$  and for  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{|t_{13}| > \delta_{13}}$ . The result follows.  $\square$

Define  $\rho_{13}(t)$  as in Lemma 4.7, i.e.,

$$\rho_{13}(t) = \alpha_{k_{13}}(\sigma_{13} + it) - \alpha_{k_{13}}(\sigma_{13}) - i\alpha'_{k_{13}}(\sigma_{13})t + \frac{1}{2}\alpha''_{k_{13}}(\sigma_{13})t^2;$$

define  $\rho_{23}$ ,  $\rho_{14}$ , and  $\rho_{24}$  similarly. Then define

$$\rho(\vec{t}) = \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \rho_{ij}(t_{ij}),$$

so that

$$G(\vec{\sigma} + i\vec{t}) - H(\vec{t}) = H(\vec{t})(e^{m\rho(\vec{t})} - 1).$$

**Lemma 5.5.** *For any  $\vec{t} \in \mathbb{R}^3$ ,*

$$|\operatorname{Im}(\rho(\vec{t}))| \leq \frac{\sqrt{2}}{3} \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} A_{ij}^{3/2} |t_{ij}|^3, \quad (5.4)$$

$$|\operatorname{Re}(\rho(\vec{t}))| \leq \frac{1}{2} \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} A_{ij}^2 t_{ij}^4. \quad (5.5)$$

Furthermore, if  $\vec{t} \in \Delta$  and  $A_{23} = A_{\min}$ , then  $|\operatorname{Re}(\rho(\vec{t}))| \leq 42D^4m^{-4/3}$ .

*Proof.* By Lemma 4.7,

$$|\operatorname{Im}(\rho(\vec{t}))| \leq -\frac{1}{6} \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \alpha_{k_{ij}}^{(3)}(\sigma_{ij}) |t_{ij}|^3,$$

$$|\operatorname{Re}(\rho(\vec{t}))| \leq \frac{1}{24} \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \alpha_{k_{ij}}^{(4)}(\sigma_{ij}) t_{ij}^4.$$

Friedman and Skoruppa proved (Lemma 5.2) that for any integer  $n \geq 2$ , any  $0 < \kappa \leq 1$ , and any  $\sigma > 0$ ,

$$\frac{|\alpha_{\kappa}^{(n)}(\sigma)|}{(\alpha_{\kappa}''(\sigma))^{n/2}} \leq \frac{(n-1)!}{\kappa^{\frac{n}{2}-1}},$$

thus the previous inequalities and  $k_{ij} \geq \frac{1}{2}$  imply inequalities (5.4) and (5.5).

When  $\vec{t} \in \Delta$ ,

$$A_{13}^2 t_{13}^4 \leq A_{13}^2 \delta_{13}^4 = A_{13}^2 \left( \frac{D}{m^{1/3} \sqrt{A_{13}}} \right)^4 = \frac{D^4}{m^{4/3}},$$

and similarly for  $A_{14}^2 t_{14}^4$  and  $A_{24}^2 t_{24}^4$ . By Jensen's inequality, we know that for any  $n \geq 1$  and any  $x, y, z \in \mathbb{R}$ ,

$$|x + y + z|^n \leq 3^{n-1}(|x|^n + |y|^n + |z|^n).$$

In particular,

$$t_{23}^4 = (t_{13} - t_{14} + t_{24})^4 \leq 27(t_{13}^4 + t_{14}^4 + t_{24}^4).$$

For  $A_{23} = A_{\min}$  and  $\vec{t} \in \Delta$ , it follows that

$$A_{23}^2 t_{23}^4 \leq 27[A_{13}^2 t_{13}^4 + A_{14}^2 t_{14}^4 + A_{24}^2 t_{24}^4] \leq 81 D^4 m^{-4/3}.$$

Then inequality (5.5) says that  $|\operatorname{Re}(\rho(\vec{t}))| \leq 42 D^4 m^{-4/3}$ . □

It follows from Lemma 4.7 that  $\operatorname{Re}(\rho)$  is an even function and  $\operatorname{Im}(\rho)$  is an odd function, in the sense that

$$\operatorname{Re}(\rho(t_1, t_2, t_3)) = \operatorname{Re}(\rho(-t_1, -t_2, -t_3)).$$

Thus  $H(\vec{t}) \operatorname{Im}(e^{m\rho(\vec{t})} - 1) = H(\vec{t})e^{m \operatorname{Re}(\rho(\vec{t}))} \sin(m \operatorname{Im}(\rho(\vec{t})))$  is odd, so

$$\begin{aligned} \iiint_{\Delta} (G(\vec{\sigma} + i\vec{t}) - H(\vec{t})) dt_3 dt_2 dt_1 &= \iiint_{\Delta} H(\vec{t})(e^{m\rho(\vec{t})} - 1) dt_3 dt_2 dt_1 \\ &= \iiint_{\Delta} H(\vec{t}) \operatorname{Re}(e^{m\rho(\vec{t})} - 1) dt_3 dt_2 dt_1. \end{aligned}$$

Now we use this fact to bound the integral.

**Lemma 5.6.** *Assume  $A_{23} = A_{\min}$ . Then*

$$\left| \iiint_{\Delta} (G(\vec{\sigma} + i\vec{t}) - H(\vec{t})) dt_3 dt_2 dt_1 \right| < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} \frac{1}{m} \left( 6 \frac{e^R - 1}{R} + \frac{80}{3} \right),$$

where  $R = 42D^4 m^{-1/3}$ .

*Proof.* By Lemmas 5.5 and 4.8,

$$\begin{aligned} \left| \iiint_{\Delta} (G(\vec{\sigma} + i\vec{t}) - H(\vec{t})) dt_3 dt_2 dt_1 \right| &= \iiint_{\Delta} H(\vec{t}) \operatorname{Re}(e^{m\rho(\vec{t})} - 1) dt_3 dt_2 dt_1 \\ &\leq \iiint_{\Delta} H(\vec{t}) \left( u(\vec{t}) \frac{e^R - 1}{R} + \frac{v(\vec{t})^2}{2} \right) dt_3 dt_2 dt_1, \end{aligned}$$

where

$$\begin{aligned} u(\vec{t}) &= \frac{1}{2} m (A_{13}^2 t_{13}^4 + A_{23}^2 t_{23}^4 + A_{14}^2 t_{14}^4 + A_{24}^2 t_{24}^4), \\ v(\vec{t}) &= \frac{\sqrt{2}}{3} m (A_{13}^{3/2} |t_{13}|^3 + A_{23}^{3/2} |t_{23}|^3 + A_{14}^{3/2} |t_{14}|^3 + A_{24}^{3/2} |t_{24}|^3). \end{aligned}$$

It follows from Lemma 5.2 that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) u(\vec{t}) \frac{e^R - 1}{R} dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} \cdot \frac{6}{m} \frac{e^R - 1}{R}.$$

By Jensen's inequality,

$$\frac{v(\vec{t})^2}{2} \leq \frac{4}{9} m^2 (A_{13}^3 t_{13}^6 + A_{14}^3 t_{14}^6 + A_{23}^3 t_{23}^6 + A_{24}^3 t_{24}^6).$$

Hence Lemma 5.2 shows that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} H(\vec{t}) \frac{v(\vec{t})^2}{2} dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} \cdot \frac{80/3}{m}.$$

□



**Lemma 5.7.** *Suppose  $m \geq 1000$ . Then*

$$f = \frac{1}{(2\pi)^3} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 = \frac{G(\vec{\sigma})}{(2\pi)^{3/2} m^{3/2} \sqrt{P}} (1 + \varphi),$$

where

$$|\varphi| < \frac{378.1}{m}.$$

*Proof.* Without loss of generality, assume  $A_{23} = A_{\min}$ . We split up the integral as

$$\iiint G = \iiint H + \iiint_{\mathbb{R}^3 \setminus \Delta} G - \iiint_{\mathbb{R}^3 \setminus \Delta} H + \iiint_{\Delta} (G - H).$$

Then Lemma 5.1 gives the main term in the estimate, and Lemmas 5.3, 5.4, and 5.6 provide the error terms. We get

$$m|\varphi| < \sqrt{\frac{P}{\Pi_{23}}} \left( 3.013 \left( 10^{-76} + \frac{2^{3/2} m^{5/6} \exp(-m^{1/3} D^2/4)}{\sqrt{\pi} D} \right) + 3 \sqrt{\frac{2}{\pi}} \frac{m^{5/6} e^{-m^{1/3} D^2/2}}{D} \right) + 6 \frac{e^{42D^4 m^{-1/3}} - 1}{42D^4 m^{-1/3}} + \frac{80}{3}.$$

Since  $P/\Pi_{23} = P/\Pi_{\max} \leq 4$ , we have

$$m|\varphi| < 2 \left( 3.013 \left( 10^{-76} + \frac{2^{3/2} m^{5/6} \exp(-m^{1/3} D^2/4)}{\sqrt{\pi} D} \right) + 3 \sqrt{\frac{2}{\pi}} \frac{m^{5/6} e^{-m^{1/3} D^2/2}}{D} \right) + 6 \frac{e^{42D^4 m^{-1/3}} - 1}{42D^4 m^{-1/3}} + \frac{80}{3}, \quad (5.6)$$

Set  $D = 1.01$ ; note that  $D < 1000^{1/3}/\sqrt{2}$ , so this choice is valid. A simple derivative check shows that the right-hand side of (5.6) is decreasing for  $m \geq 1000$ . Plugging in  $m = 1000$  yields  $|\varphi| < 378.1/m$ .  $\square$

Next we need to estimate  $f'/f$ , where  $f' = \partial f / \partial y$  is given by

$$\begin{aligned} f' &= \frac{1}{(2\pi)^3} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} -2m(s_1 + s_2 + s_3) G(s) dt_3 dt_2 dt_1 \\ &= -2m(\sigma_1 + \sigma_2 + \sigma_3) f - \frac{2im}{(2\pi)^3} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (t_1 + t_2 + t_3) G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1. \end{aligned}$$

We want to show that

$$-\frac{1}{2m} \frac{f'}{f} \rightarrow (\sigma_1 + \sigma_2 + \sigma_3) \quad \text{as } m \rightarrow \infty,$$

so we need to find an upper bound for the error term

$$-\frac{1}{2m} \frac{f'}{f} - (\sigma_1 + \sigma_2 + \sigma_3) = \frac{i}{(2\pi)^3 f} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (t_1 + t_2 + t_3) G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1. \quad (5.7)$$

Before attempting to bound the integral, we will need a few more lemmas.

**Lemma 5.8.** *Let  $i_0, i_1 \in \{1, 2\}$  and  $j_0, j_1 \in \{3, 4\}$  be given such that  $A_{i_0 j_0} \leq A_{i_1 j_1}$ . Then*

$$\sigma_{i_0 j_0} > \sigma_{i_1 j_1} / \sqrt{2}.$$

*Proof.* Recall that  $\alpha''_{\kappa}(\sigma)$  is increasing in  $\kappa$  and decreasing in  $\sigma$ . We have

$$\alpha''_{1/2}(\sigma_{i_0 j_0}) \leq A_{i_0 j_0} \leq A_{i_1 j_1} \leq \alpha''_1(\sigma_{i_1 j_1}) < \alpha''_{1/2}(\sigma_{i_1 j_1} / \sqrt{2}),$$

where we have used Lemma 4.12 for the last inequality. It follows that  $\sigma_{i_0 j_0} > \sigma_{i_1 j_1} / \sqrt{2}$ , as claimed.  $\square$

**Corollary 5.9.** *For any  $i \in \{1, 2\}$  and  $j \in \{3, 4\}$ ,*

$$\frac{\sigma_{ij}}{\sigma_1 + \sigma_2 + \sigma_3} < 1.$$

*If  $A_{23} \leq A_{14}$ , then*

$$\frac{\sigma_{14}}{\sigma_1 + \sigma_2 + \sigma_3} < 2 - \sqrt{2}.$$

*Similarly, if  $A_{24} \leq A_{13}$ , then*

$$\frac{\sigma_{13}}{\sigma_1 + \sigma_2 + \sigma_3} < 2 - \sqrt{2}.$$

*Proof.* The first claim follows from  $\sigma_1 + \sigma_2 + \sigma_3 = \sigma_{13} + \sigma_{24} = \sigma_{14} + \sigma_{23}$ . The last two claims are proven identically; we prove the first:

$$\frac{\sigma_{14}}{\sigma_1 + \sigma_2 + \sigma_3} = \frac{\sigma_{14}}{\sigma_{23} + \sigma_{14}} = \frac{1}{\frac{\sigma_{23}}{\sigma_{14}} + 1} < \frac{1}{\frac{1}{\sqrt{2}} + 1} = 2 - \sqrt{2}. \quad \square$$

**Corollary 5.10.** *Assume  $A_{23} \leq A_{14}$  and  $A_{24} \leq A_{13}$ . Then for any  $i \in \{1, 2\}$  and  $j \in \{3, 4\}$ ,*

$$\frac{1}{\sqrt{A_{ij}}} < \begin{cases} (2\sqrt{2} - 2)(\sigma_1 + \sigma_2 + \sigma_3) & \text{if } i = 1, \\ \sqrt{2}(\sigma_1 + \sigma_2 + \sigma_3) & \text{if } i = 2. \end{cases}$$

*Proof.* It follows from (4.1) that for any  $\sigma > 0$  and any  $\frac{1}{2} \leq \kappa \leq 1$ ,  $\sigma^2 \alpha''_{\kappa}(\sigma) \geq \kappa \sigma^2 \Psi'(\sigma) > \kappa$ .

Thus

$$\frac{1}{\sqrt{A_{ij}}} \frac{1}{\sigma_1 + \sigma_2 + \sigma_3} = \frac{1}{\sqrt{\sigma_{ij}^2 \alpha''_{k_{ij}}(\sigma_{ij})}} \frac{\sigma_{ij}}{\sigma_1 + \sigma_2 + \sigma_3} < \frac{1}{\sqrt{k_{ij}}} \frac{\sigma_{ij}}{\sigma_1 + \sigma_2 + \sigma_3} \leq \sqrt{2} \frac{\sigma_{ij}}{\sigma_1 + \sigma_2 + \sigma_3}.$$

Now use the previous lemma.  $\square$

Define

$$\Sigma = \{\vec{t} \in \mathbb{R}^3 \mid |t_{13}| \leq \frac{\sigma_{13}}{3\sqrt{2}}, |t_{14}| \leq \frac{\sigma_{14}}{3\sqrt{2}}, |t_{24}| \leq \frac{\sigma_{24}}{3\sqrt{2}}\}.$$

We will split the integral (5.7) into an integral over  $\Sigma$  and an integral over  $\mathbb{R}^3 \setminus \Sigma$ .

**Lemma 5.11.** *Assume that  $A_{23} = A_{min}$  and  $A_{24} \leq A_{13}$ . Then, for  $m \geq 1000$ ,*

$$\left| \iiint_{\mathbb{R}^3 \setminus \Sigma} (t_1 + t_2 + t_3) G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \right| < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{0.0008}{m}.$$

*Proof.* Note that  $t_1 + t_2 + t_3 = t_{13} + t_{24}$ , so we consider separately the integrals

$$\iiint_{\mathbb{R}^3 \setminus \Sigma} t_{13} G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \quad \text{and} \quad \iiint_{\mathbb{R}^3 \setminus \Sigma} t_{24} G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1,$$

starting with the  $t_{13}$  integral. As in the proof of Lemma 5.3, we can bound this integral by splitting it into a product of three single integrals. By Lemmas 4.9 and 4.4, the integral over the region  $|t_{13}| > \frac{\sigma_{13}}{3\sqrt{2}}$  is bounded above by

$$e^{m(\alpha_{k_{23}}(\sigma_{23}) - \vec{y} \cdot \vec{\sigma})} \frac{0.0002557\sigma_{13}e^{m\alpha_{k_{13}}(\sigma_{13})}}{m^{3/2}\sqrt{A_{13}}} \frac{1.00205\sqrt{2\pi}e^{m\alpha_{k_{24}}(\sigma_{24})}}{\sqrt{mA_{24}}} \frac{1.00205\sqrt{2\pi}e^{m\alpha_{k_{14}}(\sigma_{14})}}{\sqrt{mA_{14}}},$$

which gives an upper bound of

$$\frac{0.002\sigma_{13}G(\vec{\sigma})}{m^{5/2}\sqrt{A_{13}A_{24}A_{14}}}.$$

Similarly, Lemmas 4.11, 4.10, and 4.4 show that the integral over the region  $|t_{13}| \leq \frac{\sigma_{13}}{3\sqrt{2}}$ ,  $|t_{24}| > \frac{\sigma_{24}}{3\sqrt{2}}$  is bounded above by

$$e^{m(\alpha_{k_{23}}(\sigma_{23}) - \vec{y} \cdot \vec{\sigma})} \frac{\frac{72}{35}e^{m\alpha_{k_{13}}(\sigma_{13})}}{mA_{13}} \frac{0.00003429e^{m\alpha_{k_{24}}(\sigma_{24})}}{m\sqrt{A_{24}}} \frac{1.00205\sqrt{2\pi}e^{m\alpha_{k_{14}}(\sigma_{14})}}{\sqrt{mA_{14}}},$$

which gives an upper bound of

$$\frac{0.0002G(\vec{\sigma})}{m^{5/2}\sqrt{A_{13}^2A_{24}A_{14}}}.$$

Therefore

$$\left| \iiint_{\mathbb{R}^3 \setminus \Sigma} t_{13} G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \right| < \frac{G(\vec{\sigma})}{m^{5/2}\sqrt{A_{13}A_{24}A_{14}}} \left( 0.002\sigma_{13} + \frac{0.0004}{\sqrt{A_{13}}} \right). \quad (5.8)$$

Corollary 5.9 says that

$$0.002\sigma_{13} < 0.002(2 - \sqrt{2})(\sigma_1 + \sigma_2 + \sigma_3) < 0.002(\sigma_1 + \sigma_2 + \sigma_3).$$

Corollary 5.10 shows that

$$\frac{0.0004}{\sqrt{A_{13}}} < 0.0004(2\sqrt{2} - 2)(\sigma_1 + \sigma_2 + \sigma_3) < 0.0004(\sigma_1 + \sigma_2 + \sigma_3).$$

Since  $A_{23} = A_{\min}$ , we have  $A_{13}A_{24}A_{14} = \Pi_{\max} \geq P/4$ . Thus

$$\begin{aligned} \left| \iiint_{\mathbb{R}^3 \setminus \Sigma} t_{13} G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \right| &< (0.002 + 0.0004) \frac{G(\vec{\sigma})(\sigma_1 + \sigma_2 + \sigma_3)}{m^{5/2} \sqrt{A_{13}A_{24}A_{14}}} \\ &\leq \frac{G(\vec{\sigma})(\sigma_1 + \sigma_2 + \sigma_3)}{m^{3/2} \sqrt{P}} \cdot \frac{0.0048}{m} \\ &< \frac{(2\pi)^{3/2} G(\vec{\sigma})(\sigma_1 + \sigma_2 + \sigma_3)}{m^{3/2} \sqrt{P}} \cdot \frac{0.0004}{m}. \end{aligned}$$

Now we use the same method to bound the  $t_{24}$  integral. The argument used to prove (5.8) works equally well in this case; that is,

$$\left| \iiint_{\mathbb{R}^3 \setminus \Sigma} t_{24} G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \right| < \frac{G(\vec{\sigma})}{m^{5/2} \sqrt{A_{13}A_{24}A_{14}}} \left( 0.002\sigma_{24} + \frac{0.0004}{\sqrt{A_{24}}} \right).$$

Corollaries 5.9 and 5.10 show that

$$0.002\sigma_{24} < 0.002(\sigma_1 + \sigma_2 + \sigma_3),$$

$$\frac{0.0004}{\sqrt{A_{24}}} < 0.0004\sqrt{2}(\sigma_1 + \sigma_2 + \sigma_3) < 0.0006(\sigma_1 + \sigma_2 + \sigma_3).$$

Since  $(2\pi)^{-3/2} \cdot 2(0.002 + 0.0006) < 0.0004$ , this proves that

$$\left| \iiint_{\mathbb{R}^3 \setminus \Sigma} t_{24} G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \right| < \frac{(2\pi)^{3/2} G(\vec{\sigma})(\sigma_1 + \sigma_2 + \sigma_3)}{m^{3/2} \sqrt{P}} \cdot \frac{0.0004}{m}. \quad \square$$

It remains to consider the integral over  $\Sigma$ . The following lemma describes the behavior of the integrand for  $\vec{t} \in \Sigma$ .

**Lemma 5.12.** *Assume  $A_{23} = A_{\min}$ . Let  $\vec{t} \in \Sigma$  be given. Then*

$$|\operatorname{Re}(\rho_{23}(t_{23}))| \leq \frac{\alpha''_{k_{23}}(\sigma_{23})}{4} t_{23}^2,$$

and for  $(i, j) \neq (2, 3)$ , we have

$$|\operatorname{Re}(\rho_{ij}(t_{ij}))| \leq \frac{\alpha''_{k_{ij}}(\sigma_{ij})}{72} t_{ij}^2.$$

*Proof.* It follows from Lemma 5.8 that  $|t_{23}| < \sigma_{23}$ :

$$|t_{23}| = |t_{13} + t_{24} - t_{14}| \leq |t_{13}| + |t_{24}| + |t_{14}| \leq \frac{1}{3} \left( \frac{\sigma_{13}}{\sqrt{2}} + \frac{\sigma_{24}}{\sqrt{2}} + \frac{\sigma_{14}}{\sqrt{2}} \right) < \sigma_{23}.$$

Now use Lemma 4.7. □

Recall that  $\operatorname{Re}(\rho)$  is an even function and  $\operatorname{Im}(\rho)$  is an odd function, so

$$\begin{aligned} & \iiint_{\Sigma} (t_1 + t_2 + t_3) G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \\ &= \iiint_{\Sigma} (t_1 + t_2 + t_3) H(\vec{t}) e^{m\rho(\vec{t})} dt_3 dt_2 dt_1 \\ &= \iiint_{\Sigma} (t_1 + t_2 + t_3) H(\vec{t}) e^{m\operatorname{Re}(\rho)} [\cos(m\operatorname{Im}(\rho)) + i \sin(m\operatorname{Im}(\rho))] dt_3 dt_2 dt_1 \\ &= i \iiint_{\Sigma} (t_1 + t_2 + t_3) H(\vec{t}) e^{m\operatorname{Re}(\rho)} \sin(m\operatorname{Im}(\rho)) dt_3 dt_2 dt_1. \end{aligned} \quad (5.9)$$

By Lemma 5.5,

$$|\sin(m\operatorname{Im}(\rho(\vec{t})))| \leq |m\operatorname{Im}(\rho(\vec{t}))| \leq \frac{\sqrt{2}}{3} m \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} A_{ij}^{3/2} |t_{ij}|^3$$

Lemma 5.12 shows that for  $\vec{t} \in \Sigma$ ,

$$|\operatorname{Re}(\rho(\vec{t}))| \leq \frac{1}{72} (A_{13}t_{13}^2 + A_{14}t_{14}^2 + A_{24}t_{24}^2) + \frac{1}{4} A_{14}t_{14}^2.$$

Hence  $H(\vec{t})e^{m\operatorname{Re}(\rho)} \leq \tilde{H}(\vec{t})$ , where we define

$$\tilde{H}(\vec{t}) = G(\vec{\sigma}) \exp \left( -m \left( \frac{35}{72} A_{13}t_{13}^2 + \frac{35}{72} A_{14}t_{14}^2 + \frac{35}{72} A_{24}t_{24}^2 + \frac{1}{4} A_{23}t_{23}^2 \right) \right).$$

Plugging these inequalities into (5.9), we find that

$$\begin{aligned} & \left| \iiint_{\Sigma} (t_1 + t_2 + t_3) G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \right| \leq \\ & \frac{\sqrt{2}}{3} m \iiint_{\Sigma} |t_1 + t_2 + t_3| (A_{13}^{3/2} |t_{13}|^3 + A_{23}^{3/2} |t_{23}|^3 + A_{14}^{3/2} |t_{14}|^3 + A_{24}^{3/2} |t_{24}|^3) \tilde{H}(\vec{t}) dt_3 dt_2 dt_1. \end{aligned} \quad (5.10)$$

Note that  $\tilde{H}(\vec{t})$  is obtained from  $H(\vec{t})$  by replacing  $A_{23}$  with  $\frac{1}{2}A_{23}$  and the other  $A_{ij}$ 's with  $\frac{35}{36}A_{ij}$ , so our previous lemmas about  $H$  also apply to  $\tilde{H}$ . In particular,  $P$  is replaced by

$$\tilde{P} := P_3(\frac{35}{36}A_{13}, \frac{35}{36}A_{14}, \frac{35}{36}A_{24}, \frac{1}{2}A_{23}).$$

We will need to know how  $\tilde{P}$  compares to  $P$ .

**Lemma 5.13.** *Assume  $A_{23} = A_{\min}$ . Then  $\tilde{P} > 0.5841P$ .*

*Proof.* The assumption that  $A_{23} = A_{\min}$  ensures that  $\Pi_{23} = A_{13}A_{14}A_{24} \geq P/4$ . Thus

$$\begin{aligned} \tilde{P} &= \frac{1}{2} \left( \frac{35}{36} \right)^2 (\Pi_{13} + \Pi_{14} + \Pi_{24}) + \left( \frac{35}{36} \right)^3 \Pi_{23} \\ &= \frac{1}{2} \left( \frac{35}{36} \right)^2 P + \left( \left( \frac{35}{36} \right)^3 - \frac{1}{2} \left( \frac{35}{36} \right)^2 \right) \Pi_{23} \\ &\geq \left( \frac{1}{2} \left( \frac{35}{36} \right)^2 + \frac{1}{4} \left( \left( \frac{35}{36} \right)^3 - \frac{1}{2} \left( \frac{35}{36} \right)^2 \right) \right) P > 0.5841P. \end{aligned} \quad \square$$

Now we can bound the relevant integrals.

**Lemma 5.14.** *Assume  $A_{23} = A_{\min}$  and  $A_{24} \leq A_{13}$ . Then*

$$\begin{aligned} A_{23}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{23}^4 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{22.206}{m^2}, \\ A_{14}^{-1/2} A_{23}^2 \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{23}^4 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{13.008}{m^2}, \\ A_{13}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{13}^4 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{3.441}{m^2}, \\ A_{24}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{24}^4 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{5.874}{m^2}, \\ A_{13}^{-1/2} A_{24}^2 \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{24}^4 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{3.441}{m^2}. \end{aligned}$$

*Proof.* We prove the first claim; the rest are proven similarly. Lemma 5.2 shows that

$$\begin{aligned}
A_{23}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{23}^4 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< A_{23}^{3/2} \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\bar{P}}} \cdot \frac{3}{m^2 (\frac{1}{2} A_{23})^2} \\
&< A_{23}^{3/2} \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\bar{P}}} \cdot \frac{3 \cdot 2^2 / \sqrt{0.5841}}{m^2 A_{23}^2} \\
&< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\bar{P}}} \cdot \frac{15.7014}{m^2 \sqrt{A_{23}}}.
\end{aligned}$$

Now Corollary 5.10 shows that

$$A_{23}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{23}^4 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\bar{P}}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{15.7014 \sqrt{2}}{m^2}.$$

Note that  $15.7014 \sqrt{2} < 22.206$ . □

**Lemma 5.15.** *We have*

$$\begin{aligned}
A_{14}^{1/2} A_{23} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{14}^2 t_{23}^2 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\bar{P}}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{6.690}{m^2}, \\
A_{13}^{1/2} A_{24} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{13}^2 t_{24}^2 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\bar{P}}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{3.441}{m^2}.
\end{aligned}$$

*Proof.* Recall from the proof of Lemma 5.2 that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{23}^2 H(\vec{t}) dt_3 dt_2 dt_1 = \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{5/2} P^{3/2}} \frac{\partial P}{\partial A_{23}}.$$

It follows that

$$\begin{aligned}
&\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{14}^2 t_{23}^2 H(\vec{t}) dt_3 dt_2 dt_1 \\
&= -\frac{2}{m} \frac{\partial}{\partial A_{14}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{23}^2 H(\vec{t}) dt_3 dt_2 dt_1 \\
&= -\frac{2}{m} \frac{\partial}{\partial A_{14}} \left( \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{5/2} P^{3/2}} \frac{\partial P}{\partial A_{23}} \right) \\
&= 3 \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{7/2} P^{5/2}} \frac{\partial P}{\partial A_{14}} \frac{\partial P}{\partial A_{23}} - \frac{2}{m} \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{5/2} P^{3/2}} (A_{13} + A_{24}) \\
&< 3 \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{7/2} P^{5/2}} \frac{\partial P}{\partial A_{14}} \frac{\partial P}{\partial A_{23}}.
\end{aligned}$$



Now inequality (5.1) shows that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{14}^2 t_{23}^2 H(\vec{t}) dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} \cdot \frac{3}{m^2 A_{14} A_{23}}.$$

We conclude that

$$\begin{aligned} A_{14}^{1/2} A_{23} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} t_{14}^2 t_{23}^2 \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< A_{14}^{1/2} A_{23} \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{\tilde{P}}} \cdot \frac{3}{m^2 (\frac{35}{36} A_{14}) (\frac{1}{2} A_{23})} \\ &< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} \cdot \frac{8.075}{m^2 \sqrt{A_{23}}}, \end{aligned}$$

because  $3 \cdot \frac{36}{35} \cdot 2/\sqrt{0.5841} < 8.075$ . Corollary 5.10 completes the proof of the first claim,

because  $8.075(2\sqrt{2} - 2) < 6.690$ . The second claim is proven identically.  $\square$

**Lemma 5.16.** *If  $A_{23} = A_{min}$ , then*

$$A_{23}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |t_{14} t_{23}^3| \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{9.849}{m^2}.$$

*If furthermore  $A_{24} \leq A_{13}$ , then*

$$A_{24}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |t_{13} t_{24}^3| \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{3.441}{m^2}.$$

*Proof.* The claims follow from the previous two lemmas; by AM-GM,

$$A_{23}^{3/2} |t_{14} t_{23}^3| \leq \frac{1}{2} (A_{14}^{1/2} A_{23} t_{14}^2 t_{23}^2 + A_{14}^{-1/2} A_{23}^2 t_{23}^4).$$

$\square$

**Lemma 5.17.** *Assume  $A_{23} = A_{min}$ . Then*

$$A_{14}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |t_{24} t_{14}^3| \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{3.975}{m^2},$$

$$A_{13}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |t_{24} t_{13}^3| \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{3.975}{m^2},$$

$$A_{14}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |t_{13} t_{14}^3| \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{2.329}{m^2}.$$

*Proof.* We prove the first claim; the rest are proven similarly. Note that

$$\begin{aligned}
& A_{14}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |t_{24} t_{14}^3| \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 \\
& \leq A_{14}^{3/2} G(\vec{\sigma}) \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |t_{24} t_{14}^3| \exp\left(-\frac{35m}{72}(A_{13}t_{13}^2 + A_{14}t_{14}^2 + A_{24}t_{24}^2)\right) dt_3 dt_2 dt_1 \\
& = A_{14}^{3/2} G(\vec{\sigma}) \int_{-\infty}^{\infty} |t_{24}| e^{-\frac{35m}{72}A_{24}t_{24}^2} dt_{24} \int_{-\infty}^{\infty} |t_{14}^3| e^{-\frac{35m}{72}A_{14}t_{14}^2} dt_{14} \int_{-\infty}^{\infty} e^{-\frac{35m}{72}A_{13}t_{13}^2} dt_{13}.
\end{aligned}$$

We can use Lemma 4.6 to evaluate these integrals:

$$\begin{aligned}
\int_{-\infty}^{\infty} |t_{24}| \exp\left(-\frac{35m}{72}A_{24}t_{24}^2\right) dt_{24} &= 2 \int_0^{\infty} t_{24} \exp\left(-\frac{35m}{72}A_{24}t_{24}^2\right) dt_{24} = \frac{72/35}{mA_{24}}, \\
\int_{-\infty}^{\infty} |t_{14}^3| \exp\left(-\frac{35m}{72}A_{14}t_{14}^2\right) dt_{14} &= 2 \int_0^{\infty} t_{14}^3 \exp\left(-\frac{35m}{72}A_{14}t_{14}^2\right) dt_{14} = \frac{(72/35)^2}{m^2 A_{14}^2}, \\
\int_{-\infty}^{\infty} \exp\left(-\frac{35m}{72}A_{13}t_{13}^2\right) dt_{13} &= \frac{\sqrt{72\pi/35}}{\sqrt{mA_{13}}}.
\end{aligned}$$

Thus

$$A_{14}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |t_{24} t_{14}^3| \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 \leq \frac{22.131 G(\vec{\sigma})}{m^{3.5} \sqrt{A_{24} A_{14} A_{13}}} \frac{1}{\sqrt{A_{24}}}.$$

Using  $A_{24} A_{14} A_{13} \geq P/4$  and Corollary 5.10,

$$\begin{aligned}
A_{14}^{3/2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |t_{24} t_{14}^3| \tilde{H}(\vec{t}) dt_3 dt_2 dt_1 &< \frac{G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{62.596}{m^2} \\
&< \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{3.975}{m^2}.
\end{aligned}$$

The second claim is proven identically. The third claim is proven similarly, except that Corollary 5.10 contributes a factor of  $(2\sqrt{2} - 2)$  instead of  $\sqrt{2}$ . Note that  $3.975(2\sqrt{2} - 2)/\sqrt{2} < 2.329$ .  $\square$

Now we can estimate  $f'/f$  using inequality (5.10). Since  $t_1 + t_2 + t_3 = t_{13} + t_{24} = t_{14} + t_{23}$ ,

we have

$$|t_1 + t_2 + t_3|(A_{13}^{3/2}|t_{13}|^3 + A_{23}^{3/2}|t_{23}|^3 + A_{14}^{3/2}|t_{14}|^3 + A_{24}^{3/2}|t_{24}|^3) \leq$$

$$A_{13}^{3/2}(t_{13}^4 + |t_{24}t_{13}^3|) + A_{23}^{3/2}(|t_{14}t_{23}^3| + t_{23}^4) + A_{14}^{3/2}(|t_{13}t_{14}^3| + |t_{24}t_{14}^3|) + A_{24}^{3/2}(|t_{13}t_{24}^3| + t_{24}^4).$$

Now we can split the integral in (5.10) into eight separate integrals, which we bound using

Lemmas 5.14, 5.16, and 5.17 (still assuming that  $A_{23} = A_{\min}$  and  $A_{24} \leq A_{13}$ ):

$$\left| \iiint_{\Sigma} (t_1 + t_2 + t_3) G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \right| \left/ \left( \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \right) \right.$$

$$< \frac{1}{m} \frac{\sqrt{2}}{3} (3.441 + 3.975 + 9.849 + 22.206 + 2.329 + 3.975 + 3.441 + 5.874)$$

$$< \frac{25.9697}{m}.$$

Combining this with Lemma 5.11, we conclude that, for  $m \geq 1000$ ,

$$\left| \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (t_1 + t_2 + t_3) G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \right| < \frac{(2\pi)^{3/2} G(\vec{\sigma})}{m^{3/2} \sqrt{P}} (\sigma_1 + \sigma_2 + \sigma_3) \cdot \frac{25.971}{m}.$$

Now equation (5.7) and Lemma 5.7 show that, for  $m \geq 1000$ ,

$$\frac{1}{\sigma_1 + \sigma_2 + \sigma_3} \left| -\frac{1}{2m} \frac{f'}{f} - (\sigma_1 + \sigma_2 + \sigma_3) \right|$$

$$= \frac{1}{\sigma_1 + \sigma_2 + \sigma_3} \left| \frac{1}{(2\pi)^3 f} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (t_1 + t_2 + t_3) G(\vec{\sigma} + i\vec{t}) dt_3 dt_2 dt_1 \right|$$

$$\leq \frac{1}{|f|} \frac{G(\vec{\sigma})}{(2\pi)^{3/2} m^{3/2} \sqrt{P}} \cdot \frac{25.971}{m}$$

$$\leq \frac{1}{1 - \frac{378.1}{m}} \cdot \frac{25.971}{m}.$$

This proves the following lemma:

**Lemma 5.18.** *For  $m \geq 1000$ ,*

$$-\frac{1}{2m} \frac{f'}{f} = (\sigma_1 + \sigma_2 + \sigma_3) (1 + \beta),$$

$$\text{where } |\beta| < \frac{25.971}{m - 378.1}.$$

## Chapter 6

# Properties of $\Psi$

In this section, we prove some properties of the digamma function  $\Psi$  which will be needed when we study the saddle point  $\vec{\sigma}$  in the next section.

**Lemma 6.1.** *For any  $t > 0$ , we have the following inequalities:<sup>1</sup>*

$$\begin{aligned}\frac{t}{e^t - 1} - 1 + \frac{t}{2} &\geq 0, \\ \frac{t}{e^t - 1} - 1 + \frac{t}{2} - \frac{t^2}{12} &\leq 0, \\ \frac{t}{e^t - 1} - 1 + \frac{t}{2} - \frac{t^2}{12} + \frac{t^4}{720} &\geq 0.\end{aligned}$$

*Proof.* Since  $t/(e^t - 1) - 1 + t/2$  vanishes when  $t = 0$ , the first inequality follows from the fact that

$$\frac{d}{dt} \left( \frac{t}{e^t - 1} - 1 + \frac{t}{2} \right) = \frac{\sinh(t) - t}{2 \cosh(t) - 2} > 0 \quad \text{for } t > 0.$$

---

<sup>1</sup>These inequalities are all special cases of the conjecture that  $t/(e^t - 1)$  is enveloped by its Taylor series for  $t > 0$ .

To prove the second inequality, we consider instead  $t^2/(e^t - 1) - t + t^2/2 - t^3/12$ . Again, we need only check that its derivative is nonpositive:

$$\frac{d}{dt} \left( \frac{t^2}{e^t - 1} - t + \frac{t^2}{2} - \frac{t^3}{12} \right) = - \left( \frac{t}{e^t - 1} - 1 + \frac{t}{2} \right)^2 \leq 0.$$

The first two inequalities say that

$$\frac{t^2}{12} \geq \frac{t}{e^t - 1} - 1 + \frac{t}{2} \geq 0 \quad \text{for all } t > 0.$$

It follows that

$$\frac{d}{dt} \left( \frac{t^2}{e^t - 1} - t + \frac{t^2}{2} - \frac{t^3}{12} + \frac{t^5}{720} \right) = \left( \frac{t^2}{12} \right)^2 - \left( \frac{t}{e^t - 1} - 1 + \frac{t}{2} \right)^2 \geq 0 \quad \text{for all } t > 0,$$

which proves the last inequality.  $\square$

Recall the asymptotic series [1], 6.3.18, 6.4.12-14:

$$\begin{aligned} \Psi(x) &\sim \log x - \frac{1}{2x} - \frac{1}{12x^2} + \frac{1}{120x^4} - \frac{1}{252x^6} + \cdots, \\ \Psi'(x) &\sim \frac{1}{x} + \frac{1}{2x^2} + \frac{1}{6x^3} - \frac{1}{30x^5} + \frac{1}{42x^7} - \cdots, \\ \Psi''(x) &\sim -\frac{1}{x^2} - \frac{1}{x^3} - \frac{1}{2x^4} + \frac{1}{6x^6} - \frac{1}{6x^8} + \cdots, \\ \Psi^{(3)}(x) &\sim \frac{2}{x^3} + \frac{3}{x^4} + \frac{2}{x^5} - \frac{1}{x^7} + \frac{4}{3x^9} - \cdots. \end{aligned}$$

As these series are directly related to the Taylor series for  $t/(e^t - 1)$ , the previous lemma lets us turn these series into inequalities.

**Lemma 6.2.** *For any  $x > 0$ ,*

$$\begin{aligned} \log x - \frac{1}{2x} - \frac{1}{12x^2} &< \Psi(x) < \log x - \frac{1}{2x} - \frac{1}{12x^2} + \frac{1}{120x^4}, \\ \frac{1}{x} + \frac{1}{2x^2} + \frac{1}{6x^3} - \frac{1}{30x^5} &< \Psi'(x) < \frac{1}{x} + \frac{1}{2x^2} + \frac{1}{6x^3}, \end{aligned}$$

$$\Psi''(x) < -\frac{1}{x^2} - \frac{1}{x^3} - \frac{1}{2x^4} + \frac{1}{6x^6},$$

$$\Psi^{(3)}(x) < \frac{2}{x^3} + \frac{3}{x^4} + \frac{2}{x^5}.$$

*Proof.* Recall ([2], p. 18) that

$$\Psi(x) = \log x - \frac{1}{2x} - \int_0^\infty \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) e^{-tx} dt.$$

Differentiating, we get similar expressions for the derivatives of  $\Psi$ . We illustrate one proof; the rest are all analogous:

$$\begin{aligned} \Psi(x) &= \log x - \frac{1}{2x} - \int_0^\infty \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) e^{-tx} dt \\ &= \log x - \frac{1}{2x} - \frac{1}{12x^2} - \int_0^\infty \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} - \frac{t}{12} \right) e^{-tx} dt \\ &> \log x - \frac{1}{2x} - \frac{1}{12x^2}. \end{aligned} \quad \square$$

Recall ([1], 6.3.5) that  $\Psi(x) = -\frac{1}{x} + \Psi(x+1)$ ; taking derivatives, we get recurrence formulas for the derivatives of  $\Psi$  as well. Combining these formulas with the previous lemma, we obtain bounds for  $\Psi$  and its derivatives: for any  $x > 0$ ,

$$\Psi(x) < -\frac{1}{x} + \log(x+1) - \frac{1}{2(x+1)} - \frac{1}{12(x+1)^2} + \frac{1}{120(x+1)^4}, \quad (6.1)$$

$$\Psi(x + \frac{1}{2}) > -\frac{1}{x + \frac{1}{2}} + \log(x + \frac{3}{2}) - \frac{1}{2(x + \frac{3}{2})} - \frac{1}{12(x + \frac{3}{2})^2}, \quad (6.2)$$

$$0 < \Psi'(x) < \frac{1}{x^2} + \frac{1}{x+1} + \frac{1}{2(x+1)^2} + \frac{1}{6(x+1)^3}, \quad (6.3)$$

$$\Psi'(x + \frac{1}{2}) > \frac{1}{(x + \frac{1}{2})^2} + \frac{1}{x + \frac{3}{2}} + \frac{1}{2(x + \frac{3}{2})^2} + \frac{1}{6(x + \frac{3}{2})^3} - \frac{1}{30(x + \frac{3}{2})^5}, \quad (6.4)$$

$$\Psi''(x) < -\frac{2}{x^3} - \frac{1}{(x+1)^2} - \frac{1}{(x+1)^3} - \frac{1}{2(x+1)^4} + \frac{1}{6(x+1)^6} < 0, \quad (6.5)$$

$$0 < \Psi^{(3)}(x) < \frac{6}{x^4} + \frac{2}{(x+1)^3} + \frac{3}{(x+1)^4} + \frac{2}{(x+1)^5}. \quad (6.6)$$

**Lemma 6.3.** *For any  $x > 0$ ,*

$$\Psi(x + \frac{1}{2}) - \Psi(x) > -\frac{1}{x + \frac{1}{2}} - \frac{1}{2(x + \frac{3}{2})} - \frac{1}{12(x + \frac{3}{2})^2} + \frac{1}{x} + \frac{1}{x + 1} - \frac{1}{24(x + 1)^2} - \frac{1}{120(x + 1)^4} > 0.$$

*Proof.* Note that  $\lambda(t) := \log(1 + t) - t + \frac{1}{2}t^2 \geq 0$  for all  $t \geq 0$ ; this follows from  $\lambda(0) = 0$  and  $\lambda'(t) = t^2/(t + 1) \geq 0$  for all  $t \geq 0$ . Therefore

$$\log(x + \frac{3}{2}) - \log(x + 1) = \log\left(1 + \frac{1}{2(x + 1)}\right) \geq \frac{1}{2(x + 1)} - \frac{1}{8(x + 1)^2}.$$

Now the first inequality follows immediately from inequalities (6.1) and (6.2). To prove the second inequality, combine terms; we get a rational function with positive coefficients.  $\square$

**Lemma 6.4.** *For any  $x > 0$ ,*

$$\frac{3}{2}\Psi''(x)^2 < \Psi'(x)\Psi^{(3)}(x).$$

*Proof.* Let  $\zeta$  denote the Hurwitz zeta function. Recall from equation (4.1) that

$$\Psi^{(n)}(x) = (-1)^{n+1}n!\zeta(n + 1, x).$$

Thus the inequality is equivalent to  $\zeta(3, x)^2 < \zeta(2, x)\zeta(4, x)$ , which follows from strict log-convexity of  $n \mapsto \zeta(n, x)$ .  $\square$

**Lemma 6.5.** *For a given  $x > 0$ , the function  $\kappa \mapsto |\alpha_\kappa^{(3)}(x)|/\alpha_\kappa''(x)$  is increasing for  $\kappa \in [0, 1]$ .*

*Proof.* Observe that

$$\begin{aligned} \frac{d}{d\kappa} \frac{|\alpha_\kappa^{(3)}(x)|}{\alpha_\kappa''(x)} &= \frac{d}{d\kappa} \frac{-\alpha_\kappa^{(3)}(x)}{\alpha_\kappa''(x)} = \frac{\alpha_\kappa''(x)(\Psi''(x + \frac{1}{2}) - \Psi''(x)) + (\Psi'(x) - \Psi'(x + \frac{1}{2}))\alpha_\kappa^{(3)}(x)}{\alpha_\kappa''(x)^2} \\ &= \frac{\Psi'(x)\Psi''(x + \frac{1}{2}) - \Psi'(x + \frac{1}{2})\Psi''(x)}{\alpha_\kappa''(x)^2}, \end{aligned}$$

so we want to prove that  $\Psi'(x)\Psi''(x + \frac{1}{2}) - \Psi'(x + \frac{1}{2})\Psi''(x) > 0$ . We do so by showing that  $\Psi''/\Psi'$  is an increasing function: by Lemma 6.4,

$$\frac{d}{dx} \frac{\Psi''(x)}{\Psi'(x)} = \frac{\Psi'(x)\Psi^{(3)}(x) - \Psi''(x)^2}{\Psi'(x)^2} > 0. \quad \square$$

**Lemma 6.6.** *For any  $x > 0$  and any  $\kappa \in [\frac{1}{2}, 1]$ ,*

$$\frac{|\alpha_\kappa^{(3)}(x)|}{\alpha_\kappa''(x)} > \frac{\Psi'(x) - \Psi'(x + \frac{1}{2})}{\Psi(x + \frac{1}{2}) - \Psi(x)}.$$

*Proof.* Thanks to Lemma 6.5, we may assume  $\kappa = \frac{1}{2}$ . Recall the duplication formula for  $\Psi$ , which says that  $\alpha'_{1/2}(x) = \Psi(2x) - \log 2$ . Thus we want to show that

$$\frac{2|\Psi''(2x)|}{\Psi'(2x)} - \frac{\Psi'(x) - \Psi'(x + \frac{1}{2})}{\Psi(x + \frac{1}{2}) - \Psi(x)} > 0$$

for  $x > 0$ . By Lemma 6.3 and inequalities (6.3), (6.4), and (6.5), this function is bounded below by a rational function with positive coefficients.  $\square$



## Chapter 7

# Properties of the Saddle Point

Now we are prepared to consider the saddle point  $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ . If  $i \in \{1, 2\}$ , let  $i'$  denote the other element of  $\{1, 2\}$ ; define  $j'$  similarly for  $j \in \{3, 4\}$ . Thus

$$\sigma_{ij} + \sigma_{i'j'} = \sigma_1 + \sigma_2 + \sigma_3 \quad \text{for any } i \in \{1, 2\} \text{ and } j \in \{3, 4\}.$$

Recall from Lemma 3.2 that, for a nonzero  $a \in \mathcal{O}_L$  and  $y \in \mathbb{R}$ , the corresponding saddle point is the unique  $\vec{\sigma} \in \mathbb{R}^3$  with all  $\sigma_{ij} > 0$  which satisfies

$$a_{w_1} + y = \frac{1}{2}\alpha'_{k_{13}}(\sigma_{13}) + \frac{1}{2}\alpha'_{k_{14}}(\sigma_{14}) \tag{7.1}$$

$$a_{w_2} + y = \frac{1}{2}\alpha'_{k_{23}}(\sigma_{23}) + \frac{1}{2}\alpha'_{k_{24}}(\sigma_{24}) \tag{7.2}$$

$$a_{w_3} + y = \frac{1}{2}\alpha'_{k_{13}}(\sigma_{13}) + \frac{1}{2}\alpha'_{k_{23}}(\sigma_{23}). \tag{7.3}$$

In order for our estimates to be useful, we need to choose a  $y$  which gives a good lower bound on  $\sigma_1 + \sigma_2 + \sigma_3$ , independent of  $a$  and  $\vec{k}$ .

**Lemma 7.1.** *Let  $y_0 \in \mathbb{R}$  be given. For any  $y \geq y_0$ ,  $\vec{k} \in [\frac{1}{2}, 1]^4$ , and  $a \in L$ , the corresponding*

saddle point  $\vec{\sigma}$  satisfies

$$\sigma_1 + \sigma_2 + \sigma_3 \geq 2(\alpha'_{1/2})^{-1}(y_0).$$

In particular, if  $y \geq -1.18$ , then  $\sigma_1 + \sigma_2 + \sigma_3 \geq 1.0572$ .

*Proof.* Using the fact that  $a$  is an algebraic integer, we add equations (7.1) and (7.2) to obtain

$$2y \leq a_{w_1} + a_{w_2} + 2y = \frac{1}{2} \sum_{j=3}^4 \left( \alpha'_{k_{1j}}(\sigma_{1j}) + \alpha'_{k_{2j}}(\sigma_{2j'}) \right).$$

Thus we can choose  $j \in \{3, 4\}$  such that

$$2y \leq \alpha'_{k_{1j}}(\sigma_{1j}) + \alpha'_{k_{2j}}(\sigma_{2j'}) \leq \alpha'_{1/2}(\sigma_{1j}) + \alpha'_{1/2}(\sigma_{2j'}).$$

Since  $\alpha'_{1/2}$  is a concave function, it follows that  $y \leq \alpha'_{1/2}((\sigma_{1j} + \sigma_{2j'})/2)$ . Thus

$$\sigma_1 + \sigma_2 + \sigma_3 = \sigma_{1j} + \sigma_{2j'} \geq 2(\alpha'_{1/2})^{-1}(y) \geq 2(\alpha'_{1/2})^{-1}(y_0). \quad \square$$

Recall that, once we choose  $y$  such that  $-\frac{f'}{f}(y, a) \geq 2m$  for all  $a$ , we can ignore all terms in inequality (3.3) except for the  $a = 1$  term. It remains to understand the saddle point corresponding to  $a = 1$ . When  $a = 1$ , equations (7.1)-(7.3) say that the saddle point  $\vec{\sigma}$  satisfies

$$2y = \alpha'_{k_{13}}(\sigma_{13}) + \alpha'_{k_{14}}(\sigma_{14}) = \alpha'_{k_{23}}(\sigma_{23}) + \alpha'_{k_{24}}(\sigma_{24}) = \alpha'_{k_{13}}(\sigma_{13}) + \alpha'_{k_{23}}(\sigma_{23}),$$

or equivalently,

$$\alpha'_{k_{24}}(\sigma_{24}) + \alpha'_{k_{14}}(\sigma_{14}) = 2y, \tag{7.4}$$

$$\alpha'_{k_{13}}(\sigma_{13}) - \alpha'_{k_{24}}(\sigma_{24}) = 0, \tag{7.5}$$

$$\alpha'_{k_{14}}(\sigma_{14}) - \alpha'_{k_{23}}(\sigma_{23}) = 0. \tag{7.6}$$

Now we regard  $a = 1$  and  $y \in \mathbb{R}$  as fixed, and consider properties of  $\sigma_{ij}$  as a function of  $\vec{k}$ , defined by equations (7.4), (7.5), and (7.6). For any  $\vec{k} \in [0, 1]^4$  and any  $s_{ij} > 0$ , define

$$P_{\vec{k}}(s_{13}, s_{14}, s_{23}, s_{24}) = P_3(\alpha''_{k_{13}}(s_{13}), \alpha''_{k_{14}}(s_{14}), \alpha''_{k_{23}}(s_{23}), \alpha''_{k_{24}}(s_{24})),$$

where  $P_3$  is the degree-3 elementary symmetric polynomial in four variables. Then we wish to find an upper bound for

$$P = P(\vec{k}) = P_{\vec{k}}(\sigma_{13}(\vec{k}), \sigma_{14}(\vec{k}), \sigma_{23}(\vec{k}), \sigma_{24}(\vec{k})),$$

for  $\vec{k} \in [\frac{1}{2}, 1]^4$ . I claim that  $P(\vec{k})$  is maximized when  $\vec{k} = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ . Essentially, we will prove this by showing that  $\partial P / \partial k_{ij} < 0$ . To simplify notation, define

$$A_{ij} = \alpha''_{k_{ij}}(\sigma_{ij}) > 0,$$

$$A_{ij}^{(3)} = \alpha^{(3)}_{k_{ij}}(\sigma_{ij}) < 0,$$

$$\Delta_{ij} = \Psi(\sigma_{ij} + \frac{1}{2}) - \Psi(\sigma_{ij}) > 0,$$

$$\Delta'_{ij} = \Psi'(\sigma_{ij}) - \Psi'(\sigma_{ij} + \frac{1}{2}) > 0.$$

Note that  $\Delta_{ij} = -\partial \alpha'_{k_{ij}}(\sigma_{ij}) / \partial k_{ij}$  if we take the derivative with  $\sigma_{ij}$  fixed (not a function of  $k_{ij}$ ).

Finally, let  $Q$  denote the degree-2 elementary symmetric polynomial in 3 variables. For  $i \in \{1, 2\}$  and  $j \in \{3, 4\}$ , define

$$Q_{ij} = Q(A_{ij'}, A_{i'j}, A_{i'j'});$$

e.g.,  $Q_{13} = A_{14}A_{23} + A_{14}A_{24} + A_{23}A_{24}$ .

**Lemma 7.2.** For any  $i \in \{1, 2\}$  and  $j \in \{3, 4\}$ ,

$$\begin{aligned}\frac{\partial \sigma_{ij}}{\partial k_{ij}} &= \frac{\Delta_{ij}}{P} Q_{ij}, & \frac{\partial \sigma_{ij'}}{\partial k_{ij}} &= \frac{\Delta_{ij}}{P} A_{i'1} A_{i'2}, \\ \frac{\partial \sigma_{i'j}}{\partial k_{ij}} &= \frac{\Delta_{ij}}{P} A_{1j'} A_{2j'}, & \frac{\partial \sigma_{i'j'}}{\partial k_{ij}} &= -\frac{\Delta_{ij}}{P} A_{ij'} A_{i'j}.\end{aligned}$$

*Proof.* Applying the implicit function theorem to equations (7.4)-(7.6),

$$\begin{aligned}\begin{pmatrix} \partial \sigma_1 / \partial k_{13} \\ \partial \sigma_2 / \partial k_{13} \\ \partial \sigma_3 / \partial k_{13} \end{pmatrix} &= - \begin{pmatrix} A_{14} & A_{24} & 0 \\ A_{13} & -A_{24} & A_{13} \\ A_{14} & -A_{23} & -A_{23} \end{pmatrix}^{-1} \begin{pmatrix} 0 \\ -\Delta_{13} \\ 0 \end{pmatrix} \\ &= \frac{1}{P} \begin{pmatrix} (A_{13} + A_{24})A_{23} & A_{23}A_{24} & A_{13}A_{24} \\ A_{13}(A_{14} + A_{23}) & -A_{14}A_{23} & -A_{13}A_{14} \\ A_{14}A_{24} - A_{13}A_{23} & A_{14}(A_{23} + A_{24}) & -A_{24}(A_{13} + A_{14}) \end{pmatrix} \begin{pmatrix} 0 \\ \Delta_{13} \\ 0 \end{pmatrix} \\ &= \frac{\Delta_{13}}{P} \begin{pmatrix} A_{23}A_{24} \\ -A_{14}A_{23} \\ A_{14}(A_{23} + A_{24}) \end{pmatrix}.\end{aligned}$$

This proves the lemma for derivatives with respect to  $k_{13}$ ; the rest follow by symmetry.  $\square$

**Lemma 7.3.** For any  $i, j \in \{1, 2\}$ ,

$$\frac{\partial P}{\partial k_{ij}} = \frac{\Delta_{ij}}{P} \left( \frac{\Delta'_{ij}}{\Delta_{ij}} P Q_{ij} - |A_{ij'}^{(3)}| A_{i'1} A_{i'2} Q_{ij'} - |A_{i'j}^{(3)}| A_{1j'} A_{2j'} Q_{i'j} - |A_{ij}^{(3)}| Q_{ij}^2 + |A_{i'j'}^{(3)}| A_{ij'} A_{i'j} Q_{i'j'} \right).$$

*Proof.* By symmetry, it suffices to prove the case  $(i, j) = (1, 4)$ . Lemma 7.2 shows that

$$\begin{aligned}\frac{\partial}{\partial k_{14}} A_{13} A_{14} A_{23} &= \left( \frac{\Delta_{14}}{P} A_{13}^{(3)} A_{23} A_{24} \right) A_{14} A_{23} + A_{13} \left( \frac{\Delta_{14}}{P} A_{14}^{(3)} Q_{14} + \Delta'_{14} \right) A_{23} + \\ &\quad A_{13} A_{14} \left( -\frac{\Delta_{14}}{P} A_{23}^{(3)} A_{13} A_{24} \right), \\ \frac{\partial}{\partial k_{14}} A_{13} A_{14} A_{24} &= \left( \frac{\Delta_{14}}{P} A_{13}^{(3)} A_{23} A_{24} \right) A_{14} A_{24} + A_{13} \left( \frac{\Delta_{14}}{P} A_{14}^{(3)} Q_{14} + \Delta'_{14} \right) A_{24} + \\ &\quad A_{13} A_{14} \left( \frac{\Delta_{14}}{P} A_{24}^{(3)} A_{13} A_{23} \right),\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial k_{14}} A_{13} A_{23} A_{24} &= \left( \frac{\Delta_{14}}{P} A_{13}^{(3)} A_{23} A_{24} \right) A_{23} A_{24} + A_{13} \left( -\frac{\Delta_{14}}{P} A_{23}^{(3)} A_{13} A_{24} \right) A_{24} + \\
&\quad A_{13} A_{23} \left( \frac{\Delta_{14}}{P} A_{24}^{(3)} A_{13} A_{23} \right), \\
\frac{\partial}{\partial k_{14}} A_{14} A_{23} A_{24} &= \left( \frac{\Delta_{14}}{P} A_{14}^{(3)} Q_{14} + \Delta'_{14} \right) A_{23} A_{24} + A_{14} \left( -\frac{\Delta_{14}}{P} A_{23}^{(3)} A_{13} A_{24} \right) A_{24} + \\
&\quad A_{14} A_{23} \left( \frac{\Delta_{14}}{P} A_{24}^{(3)} A_{13} A_{23} \right).
\end{aligned}$$

Summing these equalities and simplifying yields the desired result.  $\square$

We now have formulas for  $\partial P / \partial k_{ij}$ . If we could prove that these derivatives are always negative, that would complete the proof that  $P$  is maximized when  $\vec{k} = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ . But we are not able to prove that, so we proceed in two steps. First, the next lemma narrows down the possibilities for where the maximum could occur. Then it is feasible to check by brute force that the necessary derivatives are negative.

**Lemma 7.4.** *Then there exist  $\kappa_1, \kappa_2 \in [\frac{1}{2}, 1]$  such that*

$$\max_{\vec{k} \in [\frac{1}{2}, 1]^4} P(\vec{k}) = P(\kappa_1, \frac{1}{2}, \frac{1}{2}, \kappa_2) = P(\frac{1}{2}, \kappa_1, \kappa_2, \frac{1}{2}).$$

*Proof.* First observe that  $P(\kappa_1, \frac{1}{2}, \frac{1}{2}, \kappa_2) = P(\frac{1}{2}, \kappa_1, \kappa_2, \frac{1}{2})$  is automatic, by symmetry. Now we prove that if  $P$  is maximized at  $\vec{k}$ , then  $\vec{k}$  has the form  $(\kappa_1, \frac{1}{2}, \frac{1}{2}, \kappa_2)$  or  $(\frac{1}{2}, \kappa_1, \kappa_2, \frac{1}{2})$ . This is immediate from the following claim: for every  $\vec{k} \in [\frac{1}{2}, 1]^4$ ,

$$\frac{\partial P}{\partial k_{13}}, \frac{\partial P}{\partial k_{24}} < 0 \quad \text{or} \quad \frac{\partial P}{\partial k_{14}}, \frac{\partial P}{\partial k_{23}} < 0. \quad (7.7)$$

Lemma 6.6 says that  $\Delta'_{14} / \Delta_{14} < |A_{14}^{(3)}| / A_{14}$ . Thus

$$-|A_{14}^{(3)}| Q_{14}^2 + \frac{\Delta'_{14}}{\Delta_{14}} P Q_{14} < |A_{14}^{(3)}| Q_{14} \left( \frac{P}{A_{14}} - Q_{14} \right) = |A_{14}^{(3)}| Q_{14} \frac{A_{13} A_{23} A_{24}}{A_{14}}.$$

Plugging this into the previous lemma,

$$\frac{\partial P}{\partial k_{14}} < \frac{\Delta_{14}}{P} \left( -|A_{13}^{(3)}| A_{23} A_{24} Q_{13} - |A_{24}^{(3)}| A_{13} A_{23} Q_{24} + |A_{14}^{(3)}| Q_{14} \frac{A_{13} A_{23} A_{24}}{A_{14}} + |A_{23}^{(3)}| A_{13} A_{24} Q_{23} \right).$$

Consequently,

$$\begin{aligned} \frac{P}{\Delta_{14} A_{13}^2 A_{14} A_{23}^2 A_{24}^2} \frac{\partial P}{\partial k_{14}} &< -\frac{|A_{13}^{(3)}|}{A_{13}^2} \frac{Q_{13}}{A_{14} A_{23} A_{24}} - \frac{|A_{24}^{(3)}|}{A_{24}^2} \frac{Q_{24}}{A_{13} A_{14} A_{23}} + \\ &\quad \frac{|A_{14}^{(3)}|}{A_{14}^2} \frac{Q_{14}}{A_{13} A_{23} A_{24}} + \frac{|A_{23}^{(3)}|}{A_{14}^2} \frac{Q_{23}}{A_{13} A_{14} A_{24}}. \end{aligned}$$

Hence  $\partial P / \partial k_{14}$  is negative if

$$\frac{|A_{13}^{(3)}|}{A_{13}^2} \frac{Q_{13}}{A_{14} A_{23} A_{24}} + \frac{|A_{24}^{(3)}|}{A_{24}^2} \frac{Q_{24}}{A_{13} A_{14} A_{23}} \geq \frac{|A_{14}^{(3)}|}{A_{14}^2} \frac{Q_{14}}{A_{13} A_{23} A_{24}} + \frac{|A_{23}^{(3)}|}{A_{14}^2} \frac{Q_{23}}{A_{13} A_{14} A_{24}}.$$

Similarly,  $\partial P / \partial k_{23}$  is negative if this inequality holds;  $\partial P / \partial k_{13}$  and  $\partial P / \partial k_{24}$  are negative if this inequality holds in the reverse direction. This proves claim (7.7).  $\square$

Now we need only check that  $\partial P / \partial k_{13}$  and  $\partial P / \partial k_{24}$  are negative when  $\vec{k}$  lies in the square

$$S = [\tfrac{1}{2}, 1] \times \{\tfrac{1}{2}\} \times \{\tfrac{1}{2}\} \times [\tfrac{1}{2}, 1].$$

By symmetry, it suffices to check  $\partial P / \partial k_{13} < 0$ . If  $\vec{k}$  lies in a subset of  $S$  of the form

$$S' = [k_{13}^{\min}, k_{13}^{\max}] \times \{\tfrac{1}{2}\} \times \{\tfrac{1}{2}\} \times [k_{24}^{\min}, k_{24}^{\max}],$$

we want to find an upper bound for  $\partial P / \partial k_{13}$  as a function of  $k_{13}^{\min}$ ,  $k_{13}^{\max}$ ,  $k_{24}^{\min}$ , and  $k_{24}^{\max}$ .

Then we partition the interval  $[\tfrac{1}{2}, 1]$  into  $N$  equal subintervals of length  $\frac{1}{2N}$ , thereby partitioning  $S$  into  $N^2$  equal subsquares. By choosing  $N$  sufficiently large, we will see that  $\partial P / \partial k_{13} < 0$ .

In order to bound  $\partial P / \partial k_{13}$  on  $S'$ , we will need upper/lower bounds for the  $\sigma_{ij}$ .

**Lemma 7.5.** *Let  $y \in \mathbb{R}$  be fixed. Let  $\frac{1}{2} \leq k_{13}^{min} \leq k_{13}^{max} \leq 1$  and  $\frac{1}{2} \leq k_{24}^{min} \leq k_{24}^{max} \leq 1$  be given. Define*

$$\sigma_{13}^{min} = \sigma_{13}(k_{13}^{min}, \frac{1}{2}, \frac{1}{2}, k_{24}^{max}), \quad \sigma_{13}^{max} = \sigma_{13}(k_{13}^{max}, \frac{1}{2}, \frac{1}{2}, k_{24}^{min}),$$

$$\sigma_{24}^{min} = \sigma_{24}(k_{13}^{max}, \frac{1}{2}, \frac{1}{2}, k_{24}^{min}), \quad \sigma_{24}^{max} = \sigma_{24}(k_{13}^{min}, \frac{1}{2}, \frac{1}{2}, k_{24}^{max}),$$

*Suppose  $k_{13} \in [k_{13}^{min}, k_{13}^{max}]$  and  $k_{24} \in [k_{24}^{min}, k_{24}^{max}]$ , and let  $\vec{k} = (k_{13}, \frac{1}{2}, \frac{1}{2}, k_{24})$ . Then*

$$\sigma_{13}^{min} \leq \sigma_{13}(\vec{k}) \leq \sigma_{13}^{max},$$

$$\sigma_{24}^{min} \leq \sigma_{24}(\vec{k}) \leq \sigma_{24}^{max},$$

$$\sigma_{14}(\vec{k}) = \sigma_{23}(\vec{k}) = \frac{\sigma_{13}(\vec{k}) + \sigma_{24}(\vec{k})}{2}.$$

*Proof.* The inequalities are immediate from Lemma 7.2:  $\partial\sigma_{13}/\partial k_{13}$  and  $\partial\sigma_{24}/\partial k_{24}$  are positive, while  $\partial\sigma_{13}/\partial k_{24}$  and  $\partial\sigma_{24}/\partial k_{13}$  are negative. Also,  $\sigma_{14} = \sigma_{23}$  follows from  $\alpha'_{1/2}(\sigma_{14}) = \alpha'_{1/2}(\sigma_{23})$ , and then the last claim follows from  $\sigma_{13} + \sigma_{24} = \sigma_{14} + \sigma_{23}$ .  $\square$

The lemma implies that, for  $\vec{k} \in S'$  as above,

$$A_{13}(\vec{k}) \leq \alpha''_{k_{13}^{max}}(\sigma_{13}^{min}).$$

(Recall that  $\alpha''_{\kappa}(x)$  is strictly decreasing as a function of  $x$ , and strictly increasing as a function of  $\kappa$ .) Similarly, the  $A_{ij}$ , the  $|A_{ij}^{(3)}|$ , the  $\Delta_{ij}$ , and the  $\Delta'_{ij}$  are clearly all monotone in the  $\sigma_{ij}$  and  $k_{ij}$ , so this technique gives us upper/lower bounds for every term in Lemma 7.3.

**Lemma 7.6.** *Suppose  $y = -1.18$  and  $\vec{k} \in [\frac{1}{2}, 1]^4$ . Then*

$$P(\vec{k}) \leq P(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}) < 111.78.$$

*Proof.* Use the upper bound described in the previous paragraph, partitioning  $S$  into  $30^2$  subsquares.  $\square$

For  $\vec{k} \in [0, 1]^4$ , recall that the function  $\alpha_{\vec{k}}$  is defined by

$$\alpha_{\vec{k}}(s_1, s_2, s_3) = \alpha_{k_{13}}(s_1 + s_3) + \alpha_{k_{23}}(s_2 + s_3) + \alpha_{k_{14}}(s_1) + \alpha_{k_{24}}(s_2).$$

Define

$$\begin{aligned} g(\vec{k}) &= \alpha_{\vec{k}}(\vec{\sigma}(\vec{k})) - 2y(\sigma_1(\vec{k}) + \sigma_2(\vec{k}) + \sigma_3(\vec{k})) \\ &= \alpha_{\vec{k}}(\vec{\sigma}(\vec{k})) - \vec{y} \cdot \vec{\sigma}(\vec{k}), \end{aligned}$$

where  $\vec{y} = (2y, 2y, 2y)$ .

**Lemma 7.7.** *The function  $g$  is concave on  $[0, 1]^4$ .*

*Proof.* First observe that

$$\frac{\partial g}{\partial k_{13}} = (\log \Gamma(\sigma_{13}) - \log \Gamma(\sigma_{13} + \tfrac{1}{2})) + \left( \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \alpha'_{k_{ij}}(\sigma_{ij}) \frac{\partial \sigma_{ij}}{\partial k_{13}} \right) - 2y \frac{\partial(\sigma_1 + \sigma_2 + \sigma_3)}{\partial k_{13}}.$$

Recall that  $\sigma_{13} + \sigma_{24} = \sigma_1 + \sigma_2 + \sigma_3 = \sigma_{23} + \sigma_{14}$ . Along with equations (7.4)-(7.6), this yields

$$\begin{aligned} \sum_{\substack{i \in \{1,2\} \\ j \in \{3,4\}}} \alpha'_{k_{ij}}(\sigma_{ij}) \frac{\partial \sigma_{ij}}{\partial k_{13}} &= \alpha'_{k_{13}}(\sigma_{13}) \frac{\partial \sigma_{13}}{\partial k_{13}} + \alpha'_{k_{24}}(\sigma_{24}) \frac{\partial \sigma_{24}}{\partial k_{13}} + \alpha'_{k_{14}}(\sigma_{14}) \frac{\partial \sigma_{14}}{\partial k_{13}} + \alpha'_{k_{23}}(\sigma_{23}) \frac{\partial \sigma_{23}}{\partial k_{13}} \\ &= \alpha'_{k_{13}}(\sigma_{13}) \frac{\partial \sigma_{13}}{\partial k_{13}} + \alpha'_{k_{13}}(\sigma_{13}) \frac{\partial \sigma_{24}}{\partial k_{13}} + \alpha'_{k_{14}}(\sigma_{14}) \frac{\partial \sigma_{14}}{\partial k_{13}} + \alpha'_{k_{14}}(\sigma_{14}) \frac{\partial \sigma_{23}}{\partial k_{13}} \\ &= [\alpha'_{k_{13}}(\sigma_{13}) + \alpha'_{k_{14}}(\sigma_{14})] \frac{\partial(\sigma_1 + \sigma_2 + \sigma_3)}{\partial k_{13}} \\ &= 2y \frac{\partial(\sigma_1 + \sigma_2 + \sigma_3)}{\partial k_{13}}. \end{aligned}$$



Thus we have  $\partial g / \partial k_{13} = \log \Gamma(\sigma_{13}) - \log \Gamma(\sigma_{13} + \frac{1}{2})$ . Identical arguments show that

$$\frac{\partial g}{\partial k_{ij}} = \log \Gamma(\sigma_{ij}) - \log \Gamma(\sigma_{ij} + \frac{1}{2})$$

for all  $i \in \{1, 2\}$  and  $j \in \{3, 4\}$ . It follows that, for any  $i_0, i_1 \in \{1, 2\}$  and  $j_0, j_1 \in \{3, 4\}$ ,

$$\frac{\partial}{\partial k_{i_1 j_1}} \frac{\partial g}{\partial k_{i_0 j_0}} = -\Delta_{i_0 j_0} \frac{\partial \sigma_{i_0 j_0}}{\partial k_{i_1 j_1}}.$$

Recall that  $\Delta_{i_0 j_0} > 0$ . Hence, to prove that  $g$  is convex, we must prove that the matrix

$$M = \begin{pmatrix} \frac{\partial \sigma_{13}}{\partial k_{13}} & \frac{\partial \sigma_{13}}{\partial k_{14}} & \frac{\partial \sigma_{13}}{\partial k_{23}} & \frac{\partial \sigma_{13}}{\partial k_{24}} \\ \frac{\partial \sigma_{14}}{\partial k_{13}} & \frac{\partial \sigma_{14}}{\partial k_{14}} & \frac{\partial \sigma_{14}}{\partial k_{23}} & \frac{\partial \sigma_{14}}{\partial k_{24}} \\ \frac{\partial \sigma_{23}}{\partial k_{13}} & \frac{\partial \sigma_{23}}{\partial k_{14}} & \frac{\partial \sigma_{23}}{\partial k_{23}} & \frac{\partial \sigma_{23}}{\partial k_{24}} \\ \frac{\partial \sigma_{24}}{\partial k_{13}} & \frac{\partial \sigma_{24}}{\partial k_{14}} & \frac{\partial \sigma_{24}}{\partial k_{23}} & \frac{\partial \sigma_{24}}{\partial k_{24}} \end{pmatrix}$$

is positive semidefinite. We do so by checking that the principal minors of  $M$  are nonnegative. Since  $\sigma_{13} + \sigma_{24} = \sigma_{14} + \sigma_{23}$ , we have  $\det(M) = 0$ . Lemma 7.2 shows that the  $1 \times 1$  principal minors are all positive.

Now we consider the  $2 \times 2$  principal minors. By symmetry, it suffices to consider

$$\det \begin{pmatrix} \frac{\partial \sigma_{13}}{\partial k_{13}} & \frac{\partial \sigma_{13}}{\partial k_{14}} \\ \frac{\partial \sigma_{14}}{\partial k_{13}} & \frac{\partial \sigma_{14}}{\partial k_{14}} \end{pmatrix} \quad \text{and} \quad \det \begin{pmatrix} \frac{\partial \sigma_{13}}{\partial k_{13}} & \frac{\partial \sigma_{13}}{\partial k_{24}} \\ \frac{\partial \sigma_{24}}{\partial k_{13}} & \frac{\partial \sigma_{24}}{\partial k_{24}} \end{pmatrix}.$$

By Lemma 7.2, the first is

$$\frac{\Delta_{13}\Delta_{14}}{P^2} [(A_{14}A_{23} + A_{14}A_{24} + A_{23}A_{24})(A_{13}A_{24} + A_{13}A_{23} + A_{23}A_{24}) - (A_{23}A_{24})^2] > 0,$$

and the second is

$$\frac{\Delta_{13}\Delta_{24}}{P^2} [(A_{14}A_{23} + A_{14}A_{24} + A_{23}A_{24})(A_{14}A_{23} + A_{13}A_{23} + A_{13}A_{14}) - (-A_{14}A_{23})^2] > 0.$$

It remains to consider the  $3 \times 3$  principal minors; by symmetry, we need only consider the leading  $3 \times 3$  minor. Thus, we need to check that

$$\det \begin{pmatrix} A_{14}A_{23} + A_{14}A_{24} + A_{23}A_{24} & A_{23}A_{24} & A_{14}A_{24} \\ A_{23}A_{24} & A_{13}A_{24} + A_{13}A_{23} + A_{23}A_{24} & -A_{13}A_{24} \\ A_{14}A_{24} & -A_{13}A_{24} & A_{13}A_{14} + A_{13}A_{24} + A_{14}A_{24} \end{pmatrix}$$

is positive. This is trivial: expand the determinant, and all the negative terms will be canceled by positive terms.  $\square$

**Lemma 7.8.** *Let  $y = -1.18$ . Then the minimum value of  $g(\vec{k})$  for  $\vec{k} \in [\frac{1}{2}, 1]^4$  occurs when  $\vec{k} = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ .*

*Proof.* By concavity of  $g$ , it suffices to check the vertices of  $[\frac{1}{2}, 1]^4$ . By symmetry, we need only consider

$$g(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}) \approx 3.49963,$$

$$g(1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}) \approx 3.72,$$

$$g(1, 1, \frac{1}{2}, \frac{1}{2}) \approx 3.92,$$

$$g(1, \frac{1}{2}, \frac{1}{2}, 1) \approx 3.97,$$

$$g(1, 1, 1, \frac{1}{2}) \approx 4.15,$$

$$g(1, 1, 1, 1) \approx 4.35. \quad \square$$

## Chapter 8

# Conclusion

Lemmas 5.18 and 7.1 show that, whenever  $m \geq 1000$  and  $y \geq -1.18$ , we have  $-\frac{1}{2m} \frac{f'}{f}(y, 1) >$

1. Hence inequality (3.4) holds:

$$\frac{\text{Reg}_{K_1, K_2}(L)}{\#\mu_L} \geq 2^{3-|\mathcal{A}_L|} \pi^{-r_2(L)/2} \left( -1 - \frac{1}{2m} \frac{f'}{f}(y, 1) \right) f(y, 1).$$

We also have, in the notation of the previous section,

$$\begin{aligned} f(y, 1) &\geq \frac{\exp(mg(\vec{k}))}{(2\pi)^{3/2} m^{3/2} \sqrt{P(\vec{k})}} \left( 1 - \frac{378.1}{m} \right), \\ -\frac{1}{2m} \frac{f'}{f}(y, 1) &\geq (\sigma_1 + \sigma_2 + \sigma_3) \left( 1 - \frac{25.971}{m - 378.1} \right). \end{aligned}$$

Thus Lemmas 7.1, 7.6, and 7.8 show that, for  $m \geq 1000$ ,

$$\begin{aligned} f(-1.18, 1) &> \frac{e^{3.49962m}}{(2\pi)^{3/2} m^{3/2} \sqrt{111.78}} \left( 1 - \frac{378.1}{m} \right) > (1.866 \times 10^{-8}) e^{3.4995m}, \\ -1 - \frac{1}{2m} \frac{f'}{f}(-1.18, 1) &\geq -1 + 1.0572 \left( 1 - \frac{25.971}{m - 378.1} \right) > 0.0287. \end{aligned}$$

Hence for  $m \geq 1000$ ,

$$\frac{\text{Reg}_{K_1, K_2}(L)}{\#\mu_L} > (4.28 \times 10^{-9}) e^{3.4995m} 2^{-|\mathcal{A}_L|} \pi^{-r_2(L)/2}.$$

Note that  $|\mathcal{A}_L| + r_2(L) = [L : \mathbb{Q}] = 4m$ . Thus,

$$2^{-|\mathcal{A}_L|} \pi^{-r_2(L)/2} = 2^{-4m+r_2(L)} \pi^{-r_2(L)/2} \geq 16^{-m},$$

as  $2^{-4m+r_2} \pi^{-r_2/2}$  is minimized (as a function of  $r_2$ ) when  $r_2 = 0$ . We conclude that, for

$$[L : K] \geq 1000,$$

$$\frac{\text{Reg}_{K_1, K_2}(L)}{\#\mu_L} \geq (4.28 \times 10^{-9}) e^{3.4995m} 16^{-m} > (4.28 \times 10^{-9}) \cdot 2.0686^m.$$

# Appendices

# Appendix A

## Proof of Lemma 5.8

Friedman and Skoruppa made an error in the proof of their Lemma 5.6, while bounding

$$J_1 := \left| \int_{-\sigma}^{\sigma} t e^{-m\alpha''(\sigma)t^2/2+m\rho} dt \right|.$$

(See their paper for the notation.) They claim that

$$J_1 \leq \frac{\sqrt{2\pi}}{2m^{3/2}\alpha''(\sigma)} \frac{|\alpha^{(3)}(\sigma)|}{(\alpha''(\sigma))^{3/2}} \leq \frac{\sqrt{2\pi}}{\sqrt{\kappa}m^{3/2}\alpha''(\sigma)}. \quad (\text{A.1})$$

However, the first inequality is incorrect. In fact, they found the asymptotic behavior of  $J_1$ , not a bound for  $J_1$ :

$$J_1 \sim \frac{\sqrt{2\pi}}{2m^{3/2}\alpha''(\sigma)} \frac{|\alpha^{(3)}(\sigma)|}{(\alpha''(\sigma))^{3/2}} \quad \text{as } m \rightarrow \infty.$$

Lemma 5.6 is used only to prove Lemma 5.7, which is used only to prove Lemma 5.8. As such, we do not attempt to prove Lemmas 5.6 or 5.7 here, only Lemma 5.8. Furthermore, we prove it only in the cases which are needed for their paper, namely  $\kappa \in [\frac{1}{2}, 1]$ ,  $\sigma > 0$ , and integers  $m \geq 40$ . We proceed in three steps:

- Prove Lemma 5.6 (and thus Lemma 5.8) for  $\kappa \in [\frac{1}{2}, 1]$ ,  $\sigma \geq 0.65$ , and  $m \geq 40$ .

- Prove Lemma 5.8 for  $\kappa \in [\frac{1}{2}, 1]$ ,  $\sigma > 0$ , and  $m \geq 140$ .
- It remains only to consider  $\kappa \in [\frac{1}{2}, 1]$ ,  $\sigma \in (0, 0.65]$ , and  $m \in \mathbb{Z} \cap [40, 139]$ . A brute-force computer search can check that Lemma 5.8 holds in this region.

## A.1 Proof for $\sigma \geq 0.65$

We prove that Friedman and Skoruppa's bound for  $J_1$  in the proof of their Lemma 5.6 is valid for  $\sigma \geq 0.65$ , for all  $m \geq 40$  and all  $\kappa \in [\frac{1}{2}, 1]$ . We start by collecting some lemmas.

**Lemma A.1.** *For any  $0 \leq t \leq \sigma$ ,*

$$\begin{aligned} 0 \leq \operatorname{Re}(\rho(t)) &\leq t^4 \frac{\alpha^{(4)}(\sigma)}{4!}, \\ 0 \leq \operatorname{Im}(\rho(t)) &\leq -t^3 \frac{\alpha^{(3)}(\sigma)}{3!} = t^3 \frac{|\alpha^{(3)}(\sigma)|}{3!}. \end{aligned}$$

*Furthermore,  $\operatorname{Re}(\rho(t))$  is an even function and  $\operatorname{Im}(\rho(t))$  is an odd function.*

*Proof.* See Friedman and Skoruppa's Lemma 5.1. □

**Lemma A.2.** *Let an integer  $n \geq 2$  be given. The function*

$$\frac{|\Psi^{(n-1)}(s)|}{\Psi'(s)^{n/2}}$$

*is strictly decreasing for  $s > 0$ .*

*Proof.* Recall that  $\Psi^{(k)}(s) = (-1)^{k+1} k! \zeta(k+1, s)$ , where  $\zeta$  is the Hurwitz zeta function.

Thus we need to show that  $\zeta(n, s)/\zeta(2, s)^{n/2}$  is a decreasing function. We have

$$\frac{d}{ds} \frac{\zeta(n, s)}{\zeta(2, s)^{n/2}} = -n \frac{\zeta(n+1, s)\zeta(2, s) - \zeta(n, s)\zeta(3, s)}{\zeta(2, s)^{\frac{n}{2}+1}} < 0,$$

where the inequality follows from log-convexity of  $k \mapsto \zeta(k, s)$ . □

**Lemma A.3.** *Let  $\sigma_0 > 0$  and an integer  $n \geq 2$  be given. Define*

$$E = \frac{|\Psi^{(n-1)}(\sigma_0)|}{(n-1)!\Psi'(\sigma_0)^{n/2}}.$$

*Then for all  $\sigma \geq \sigma_0$ ,*

$$\frac{|\alpha^{(n)}(\sigma)|}{\alpha''(\sigma)^{n/2}} \leq \frac{(n-1)!E}{\kappa^{\frac{n}{2}-1}}.$$

*Proof.* Set

$$\begin{aligned} t &= \frac{1-\kappa}{\kappa} \geq 0, \quad A = \Psi'(\sigma), \quad B = \Psi'(\sigma + \tfrac{1}{2}), \\ C &= \frac{|\Psi^{(n-1)}(\sigma)|}{(n-1)!}, \quad D = \frac{|\Psi^{(n-1)}(\sigma + \tfrac{1}{2})|}{(n-1)!}. \end{aligned}$$

The lemma is equivalent to the positivity of

$$f(t) := (A + tB)^{n/2}E - C - tD.$$

We shall prove that  $f(t) \geq 0$  by showing  $f(0) \geq 0$ ,  $f'(0) \geq 0$ , and  $f''(t) \geq 0$  for all  $t \geq 0$ .

Lemma A.2 implies that

$$E \geq \frac{|\Psi^{(n-1)}(s)|/(n-1)!}{\Psi'(s)^{n/2}} \quad \text{for all } s \geq \sigma_0.$$

Taking  $s = \sigma$  shows that  $A^{\frac{n}{2}}E \geq C$ . Taking  $s = \sigma + \frac{1}{2}$  shows that  $B^{\frac{n}{2}}E \geq D$ .

We have  $f(0) = A^{n/2}E - C \geq 0$ . Next observe that  $A > B$  because  $\Psi'$  is decreasing, so

$$f'(0) = \tfrac{n}{2}A^{\frac{n}{2}-1}BE - D \geq A^{\frac{n}{2}-1}BE - D > B^{\frac{n}{2}}E - D \geq 0.$$

Finally,  $f''(t) = \tfrac{n}{2}(\tfrac{n}{2}-1)(A+tB)^{\frac{n}{2}-2}B^2E \geq 0$ . □

**Corollary A.4.** *For all  $\sigma \geq 0.65$ ,*

$$\frac{|\alpha^{(3)}(\sigma)|}{\alpha''(\sigma)^{3/2}} < \frac{1.39154}{\sqrt{\kappa}} \quad \text{and} \quad \frac{\alpha^{(4)}(\sigma)}{\alpha''(\sigma)^2} < \frac{3.3975}{\kappa}.$$



*Proof.* Note that

$$\frac{|\Psi^{(2)}(0.65)|}{\Psi'(0.65)^{3/2}} < 1.39154, \quad \frac{\Psi^{(3)}(0.65)}{\Psi'(0.65)^2} < 3.3975.$$

□

**Lemma A.5.** *Let  $s_0 \in (0, 0.65]$  be given. Then the function  $f(s) = s^2\Psi^{(3)}(s)/\Psi'(s)$  is maximized for  $s \in [s_0, \infty)$  at  $s = s_0$ .*

*Proof.* We can replace  $f$  with  $g(s) = s^2\zeta(4, s)/\zeta(2, s)$ . We prove the claim in two steps:

- The function  $g$  is decreasing on the interval  $(0, 0.85]$ . [We could also prove the claim simply by showing that  $g$  is decreasing on  $(0, \infty)$ .]
- If  $s \geq 0.85$ , then  $g(s) < g(s_0)$ . We prove this by finding a decreasing function  $h$  with  $h(0.85) < g(0.65)$  and  $g(s) < h(s)$  for all  $s$ . Hence  $g(s) < h(s) \leq h(0.85) < g(0.65) \leq g(s_0)$  for all  $s \geq 0.85$ .

We want to show that

$$g'(s) = \frac{2s\zeta(4, s)\zeta(2, s) - 4s^2\zeta(5, s)\zeta(2, s) + 2s^2\zeta(4, s)\zeta(3, s)}{\zeta(2, s)^2}$$

is negative for  $s \in (0, 0.85]$ . We estimate the zeta functions by integrals:

$$\begin{aligned} \frac{1}{s^2} + \frac{1}{(s+1)^2} + \frac{1}{s+2} &< \zeta(2, s) < \frac{1}{s^2} + \frac{1}{(s+1)^2} + \frac{1}{s+1}, \\ \zeta(3, s) &< \frac{1}{s^3} + \frac{1}{(s+1)^3} + \frac{1}{2(s+1)^2}, \\ \zeta(4, s) &< \frac{1}{s^4} + \frac{1}{(s+1)^4} + \frac{1}{3(s+1)^3}, \\ \frac{1}{s^5} + \frac{1}{(s+1)^5} + \frac{1}{4(s+2)^4} &< \zeta(5, s). \end{aligned}$$

Plugging in these estimates, we find that  $2s\zeta(4, s)\zeta(2, s) - 4s^2\zeta(5, s)\zeta(2, s) + 2s^2\zeta(4, s)\zeta(3, s)$  is bounded above by a rational function, which is easily checked to be negative on the interval  $(0, 0.85]$ . Thus  $g' < 0$  on  $(0, 0.85]$ .

Again using the above estimates for the zeta function, we have

$$g(s) < s^2 \frac{\frac{1}{s^4} + \frac{1}{(s+1)^4} + \frac{1}{3(s+1)^3}}{\frac{1}{s^2} + \frac{1}{(s+1)^2} + \frac{1}{s+2}} =: h(s).$$

Note that

$$h'(s) = -\frac{3s^9 + 17s^8 + 52s^7 + 154s^6 + 361s^5 + 535s^4 + 455s^3 + 207s^2 + 36s}{3(s+1)^3(s^4 + 4s^3 + 7s^2 + 5s + 2)^2} < 0$$

and that  $h(0.85) < g(0.65)$ , so  $h$  has the claimed properties.  $\square$

**Lemma A.6.** *Let  $\sigma_0 \in (0, 0.65]$  be given. For all  $\sigma \geq \sigma_0$ ,*

$$\sigma^2 \alpha^{(4)}(\sigma) \leq \frac{\sigma_0^2 \Psi^{(3)}(\sigma_0)}{\Psi'(\sigma_0)} \alpha''(\sigma).$$

*Proof.* Lemma A.5 says that

$$\sigma^2 \Psi^{(3)}(\sigma) \leq \frac{\sigma_0^2 \Psi^{(3)}(\sigma_0)}{\Psi'(\sigma_0)} \Psi'(\sigma),$$

$$\sigma^2 \Psi^{(3)}(\sigma + \tfrac{1}{2}) < (\sigma + \tfrac{1}{2})^2 \Psi^{(3)}(\sigma + \tfrac{1}{2}) \leq \frac{\sigma_0^2 \Psi^{(3)}(\sigma_0)}{\Psi'(\sigma_0)} \Psi'(\sigma + \tfrac{1}{2}),$$

and combining these inequalities yields the lemma.  $\square$

**Corollary A.7.** *For all  $\sigma > 0$ ,  $\sigma^2 \alpha^{(4)}(\sigma) \leq 6\alpha''(\sigma)$ . For all  $\sigma \geq 0.65$ ,  $\sigma^2 \alpha^{(4)}(\sigma) < 4.58111\alpha''(\sigma)$ .*

*Proof.* Note that  $\sigma^2 \Psi^{(3)}(\sigma)/\Psi'(\sigma) \rightarrow 6$  as  $\sigma \rightarrow 0$ . Also,  $0.65^2 \Psi^{(3)}(0.65)/\Psi'(0.65) < 4.58111$ .  $\square$

Now we are prepared to bound  $J_1$ . Recall that, by Lemma A.1,

$$\begin{aligned} \frac{J_1}{2} &= \left| \int_0^\sigma t e^{-m\alpha''(\sigma)t^2/2 + \operatorname{Re}(\rho(t))} \sin(m \operatorname{Im}(\rho(t))) dt \right| \\ &\leq m \frac{|\alpha^{(3)}(\sigma)|}{3!} \int_0^\sigma t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt. \end{aligned}$$

**Lemma A.8.** *Suppose  $0 < L < U$  and  $\sigma \geq 0.65$ . Let*

$$\beta_U = \left( \frac{1}{2} - \frac{3.3975U}{480} \right).$$

*Then*

$$\int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt \leq \frac{1}{(m\alpha''(\sigma))^{5/2}} \frac{1}{2\beta_U^{5/2}} \int_{\beta_U L}^{\beta_U U} s^{3/2} e^{-s} ds.$$

*Proof.* By Corollary A.4, for any  $0 \leq t \leq \sqrt{U/(m\alpha''(\sigma))}$ ,

$$\frac{\alpha^{(4)}(\sigma)t^2}{24} \leq \frac{U}{m\alpha''(\sigma)} \frac{\alpha^{(4)}(\sigma)}{24} \leq \frac{3.3975U}{24m\kappa} \alpha''(\sigma) \leq \frac{3.3975U}{480} \alpha''(\sigma).$$

It follows that

$$-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24 \leq -\left( \frac{1}{2} - \frac{3.3975U}{480} \right) m\alpha''(\sigma)t^2 = -\beta_U m\alpha''(\sigma)t^2.$$

Thus, making the substitution  $s = \beta_U m\alpha''(\sigma)t^2$ ,

$$\begin{aligned} \int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt &\leq \int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t^4 e^{-\beta_U m\alpha''(\sigma)t^2} dt \\ &= \frac{1}{(m\alpha''(\sigma))^{5/2}} \frac{1}{2\beta_U^{5/2}} \int_{\beta_U L}^{\beta_U U} s^{3/2} e^{-s} ds. \quad \square \end{aligned}$$

**Lemma A.9.** *Let  $L \geq 0$  be given, and suppose  $\sigma \geq 0.65$ . Then*

$$\int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sigma} t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt \leq \frac{1}{(m\alpha''(\sigma))^{5/2}} \cdot \frac{1}{2 \cdot 0.309^{5/2}} \int_{0.309L}^{\infty} s^{3/2} e^{-s} ds.$$

*Proof.* By Corollary A.7, for any  $0 \leq t \leq \sigma$ ,

$$\frac{\alpha^{(4)}(\sigma)t^2}{24} \leq \frac{\sigma^2\alpha^{(4)}(\sigma)}{24} \leq \frac{4.58111\alpha''(\sigma)}{24}.$$

It follows that

$$-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24 \leq -\left(\frac{1}{2} - \frac{4.58111}{24}\right)m\alpha''(\sigma)t^2 \leq -0.309m\alpha''(\sigma)t^2.$$

The rest of the argument is identical to the previous proof, with 0.309 replacing  $\beta_U$ .  $\square$

Let  $T = \sqrt{26/(m\alpha''(\sigma))}$ . Now we estimate  $J_1$  by considering separately the integral over  $[0, T]$  and the integral over  $[T, \sigma]$ . We use Lemma A.8 for the first integral, splitting  $[0, T]$  into 100 separate intervals:

$$\begin{aligned} \int_0^T t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt &= \sum_{k=1}^{100} \int_{\sqrt{\frac{.26(k-1)}{m\alpha''(\sigma)}}}^{\sqrt{\frac{.26k}{m\alpha''(\sigma)}}} t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt \\ &\leq \frac{1}{(m\alpha''(\sigma))^{5/2}} \sum_{k=1}^{100} \frac{1}{2\beta_{.26k}^{5/2}} \int_{.26(k-1)\beta_{.26k}}^{.26k\beta_{.26k}} s^{3/2} e^{-s} ds \\ &< \frac{5.318}{(m\alpha''(\sigma))^{5/2}}. \end{aligned}$$

We use Lemma A.9 for the second integral:

$$\begin{aligned} \int_T^\sigma t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt &\leq \frac{1}{(m\alpha''(\sigma))^{5/2}} \cdot \frac{1}{2 \cdot 0.309^{5/2}} \int_{0.309 \cdot 26}^\infty s^{3/2} e^{-s} ds \\ &< \frac{0.0834}{(m\alpha''(\sigma))^{5/2}}. \end{aligned}$$

Combining these bounds and applying Corollary A.4, we conclude that

$$\begin{aligned}
\frac{J_1}{2} &< m \frac{|\alpha^{(3)}(\sigma)|}{3!} \frac{5.4014}{(m\alpha''(\sigma))^{5/2}} \\
&< \frac{5.4014 \cdot 1.39154/3!}{\sqrt{\kappa} m^{3/2} \alpha''(\sigma)} \\
&< \frac{1.253}{\sqrt{\kappa} m^{3/2} \alpha''(\sigma)} \\
&< \frac{\sqrt{2\pi}/2}{\sqrt{\kappa} m^{3/2} \alpha''(\sigma)},
\end{aligned}$$

which is the desired inequality.

## A.2 Proof for $m \geq 140$

Now we prove the inequality for all  $\kappa \in [\frac{1}{2}, 1]$ ,  $\sigma > 0$ , and  $m \geq 140$ . We follow Friedman and Skoruppa's proof closely, giving stronger versions of Lemmas 5.5 and 5.7 to compensate for the weaker, corrected Lemma 5.6. This involves few new ideas; mostly we just plug in  $m = 140$  where they plugged in  $m = 40$ .

### Lemma 5.5

First, we give an improved version of Lemma 5.5 for  $m \geq 140$ .

**Lemma A.10.** *If  $m \geq 140$ , then*

$$F_\kappa(y, m) = \frac{e^{m\alpha(\sigma) - my\sigma}}{\sqrt{2\pi m\alpha''(\sigma)}} \left(1 + \frac{\gamma}{m}\right) \quad |\gamma| < 5.23.$$

*Proof.* We have

$$\frac{2^{\frac{3}{2}} \sigma \sqrt{m\alpha''(\sigma)}}{\sqrt{\pi}(m\kappa - 2) 1.25^{m\kappa[\sigma]/2} 2^{m\kappa/2}} = \frac{4}{m\kappa^{3/2} \sqrt{\pi}} \frac{\sqrt{m\kappa/2}}{2^{m\kappa/2}} \frac{m\kappa}{m\kappa - 2} \frac{\sqrt{\sigma^2 \alpha''(\sigma)}}{1.25^{m\kappa[\sigma]/2}} < \frac{10^{-8}}{m}.$$

Also, if we define  $q(m)$  as in the proof of Lemma 5.5, then  $\max_{m \geq 140} q(m) = q(140) < 5.224$ .

It follows that  $|\gamma| \leq 5.224 + 10^{-8} < 5.23$ .  $\square$

### Lemma 5.6

We will need a version of Lemma 5.6 that is valid for  $m \geq 140$ . Since the estimate for  $J_2$  in the paper is fine, we need only estimate  $J_1$ . First, some lemmas:

**Lemma A.11.** *For any  $\sigma > 0$ ,*

$$\sigma^2 \alpha^{(4)}(\sigma) < 6\alpha''(\sigma).$$

*Proof.* Clearly it suffices to check that  $\sigma^2 \Psi^{(3)}(\sigma) < 6\Psi'(\sigma)$ . Recall that

$$\Psi^{(k)}(\sigma) = (-1)^{k+1} k! \zeta(k+1, \sigma),$$

where  $\zeta$  is the Hurwitz zeta function. Estimating the zeta functions by integrals, we obtain the inequalities

$$\begin{aligned} \zeta(4, \sigma) &< \frac{1}{\sigma^4} + \frac{1}{(\sigma+1)^4} + \frac{1}{3(\sigma+1)^3}, \\ \zeta(2, \sigma) &> \frac{1}{\sigma^2} + \frac{1}{(\sigma+1)^2} + \frac{1}{\sigma+2}. \end{aligned}$$

It follows that, for any  $\sigma > 0$ ,

$$\frac{\sigma^2 \Psi^{(3)}(\sigma)}{\Psi'(\sigma)} = \frac{6\sigma^2 \zeta(4, \sigma)}{\zeta(2, \sigma)} < 6 \frac{\frac{1}{\sigma^2} + \frac{\sigma^2}{(\sigma+1)^4} + \frac{\sigma^2}{3(\sigma+1)^3}}{\frac{1}{\sigma^2} + \frac{1}{(\sigma+1)^2} + \frac{1}{\sigma+2}} < 6. \quad \square$$

Now we are prepared to bound  $J_1$ . Recall that, using Lemma 5.1,

$$\begin{aligned} \frac{J_1}{2} &= \left| \int_0^\sigma t e^{-m\alpha''(\sigma)t^2/2 + m \operatorname{Re}(\rho(t))} \sin(m \operatorname{Im}(\rho(t))) dt \right| \\ &\leq m \frac{|\alpha^{(3)}(\sigma)|}{3!} \int_0^\sigma t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt. \end{aligned}$$

**Lemma A.12.** Suppose  $0 < L < U < 140$ . Let

$$\beta_U = \frac{1}{2} - \frac{U}{280}.$$

If  $m \geq 140$ , then

$$\int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt \leq \frac{1}{(m\alpha''(\sigma))^{5/2}} \frac{1}{2\beta_U^{5/2}} \int_{\beta_U L}^{\beta_U U} s^{3/2} e^{-s} ds.$$

*Proof.* By Lemma 5.2, for any  $0 \leq t \leq \sqrt{U/(m\alpha''(\sigma))}$ ,

$$\frac{\alpha^{(4)}(\sigma)t^2}{24} \leq \frac{U}{m\alpha''(\sigma)} \frac{\alpha^{(4)}(\sigma)}{24} \leq \frac{6U}{24m\kappa} \alpha''(\sigma) \leq \frac{U}{280} \alpha''(\sigma).$$

It follows that

$$-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24 \leq -\left(\frac{1}{2} - \frac{U}{280}\right) m\alpha''(\sigma)t^2 = -\beta_U m\alpha''(\sigma)t^2.$$

Thus, making the substitution  $s = \beta_U m\alpha''(\sigma)t^2$ ,

$$\begin{aligned} \int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt &\leq \int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t^4 e^{-\beta_U m\alpha''(\sigma)t^2} dt \\ &= \frac{1}{(m\alpha''(\sigma))^{5/2}} \frac{1}{2\beta_U^{5/2}} \int_{\beta_U L}^{\beta_U U} s^{3/2} e^{-s} ds. \quad \square \end{aligned}$$

**Lemma A.13.** Let  $L \geq 0$  be given. For any  $m > 0$ ,

$$\int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sigma} t^4 e^{-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24} dt \leq \frac{16}{(m\alpha''(\sigma))^{5/2}} \int_{L/4}^{\infty} s^{3/2} e^{-s} ds.$$

*Proof.* By Lemma A.11, for any  $0 \leq t \leq \sigma$ ,

$$\frac{\alpha^{(4)}(\sigma)t^2}{24} \leq \frac{\sigma^2 \alpha^{(4)}(\sigma)}{24} \leq \frac{\alpha''(\sigma)}{4}.$$

It follows that

$$-m\alpha''(\sigma)t^2/2 + m\alpha^{(4)}(\sigma)t^4/24 \leq -\left(\frac{1}{2} - \frac{1}{4}\right) m\alpha''(\sigma)t^2 = -\frac{1}{4} m\alpha''(\sigma)t^2.$$

The rest of the argument is identical to the previous proof, with  $1/4$  replacing  $\beta_U$ .  $\square$

Let  $T = \sqrt{50/(m\alpha''(\sigma))}$ . Now we estimate  $J_1$  by considering separately the integral over  $[0, T]$  and the integral over  $[T, \sigma]$ . We use Lemma A.12 for the first integral, splitting  $[0, T]$  into 200 separate intervals:

$$\begin{aligned} \int_0^T t^4 e^{-m\alpha''(\sigma)t^2/2+m\alpha^{(4)}(\sigma)t^4/24} dt &= \sum_{k=1}^{200} \int_{\sqrt{\frac{(k-1)/4}{m\alpha''(\sigma)}}}^{\sqrt{\frac{k/4}{m\alpha''(\sigma)}}} t^4 e^{-m\alpha''(\sigma)t^2/2+m\alpha^{(4)}(\sigma)t^4/24} dt \\ &\leq \frac{1}{(m\alpha''(\sigma))^{5/2}} \sum_{k=1}^{200} \frac{1}{2\beta_{k/4}^{5/2}} \int_{\frac{1}{4}(k-1)\beta_{k/4}}^{\frac{1}{4}k\beta_{k/4}} s^{3/2} e^{-s} ds \\ &< \frac{4.353}{(m\alpha''(\sigma))^{5/2}}. \end{aligned}$$

We use Lemma A.13 for the second integral:

$$\begin{aligned} \int_T^\sigma t^4 e^{-m\alpha''(\sigma)t^2/2+m\alpha^{(4)}(\sigma)t^4/24} dt &\leq \frac{16}{(m\alpha''(\sigma))^{5/2}} \int_{50/4}^\infty s^{3/2} e^{-s} ds \\ &< \frac{0.003}{(m\alpha''(\sigma))^{5/2}}. \end{aligned}$$

Combining these bounds and applying Lemma 5.2, we conclude that

$$\begin{aligned} \frac{J_1}{2} &< m \frac{|\alpha^{(3)}(\sigma)|}{3!} \frac{4.36}{(m\alpha''(\sigma))^{5/2}} \\ &< \frac{4.36 \cdot 2/3!}{\sqrt{\kappa} m^{3/2} \alpha''(\sigma)} \\ &< \frac{1.455}{\sqrt{\kappa} m^{3/2} \alpha''(\sigma)}. \end{aligned}$$

Combining this with the existing bound for  $J_2$  gives us our new version of Lemma 5.6:

**Lemma A.14.** *Suppose  $m \geq 140$ . Then*

$$\left| \int_{-\infty}^\infty t e^{m(\alpha(\sigma+it)-iyt)} dt \right| \leq \frac{4e^{m\alpha(\sigma)} \sigma^2}{(m\kappa - 2) 1.25^{m\kappa[\sigma]/2} 2^{m\kappa/2}} + \frac{2.91e^{m\alpha(\sigma)}}{\sqrt{\kappa} m^{3/2} \alpha''(\sigma)}.$$

**Lemma 5.7**

Our new version of Lemma 5.6 translates directly into a new version of Lemma 5.7:



**Lemma A.15.** *Assume  $m \geq 140$ . Then*

$$\left| \int_{-\infty}^{\infty} t e^{m(\alpha(\sigma+it)-iyt)} dt \right| \leq \frac{2.322\sigma}{m} \frac{\sqrt{2\pi} e^{m\alpha(\sigma)}}{\sqrt{m\alpha''(\sigma)}}.$$

*Proof.* Lemma A.14 says that

$$\left| \int_{-\infty}^{\infty} t e^{m(\alpha(\sigma+it)-iyt)} dt \right| \leq \frac{\sigma \sqrt{2\pi} e^{m\alpha(\sigma)}}{\sqrt{m\alpha''(\sigma)}} \left( \frac{2^{\frac{3}{2}} \sigma \sqrt{m\alpha''(\sigma)}}{\sqrt{\pi}(m\kappa - 2) 1.25^{m\kappa[\sigma]/2} 2^{m\kappa/2}} + \frac{2.91/\sqrt{2\pi}}{m \sqrt{\kappa \sigma^2 \alpha''(\sigma)}} \right).$$

We saw in the proof of Lemma A.10 that

$$\frac{2^{\frac{3}{2}} \sigma \sqrt{m\alpha''(\sigma)}}{\sqrt{\pi}(m\kappa - 2) 1.25^{m\kappa[\sigma]/2} 2^{m\kappa/2}} \leq \frac{10^{-8}}{m}.$$

We know that  $\sigma^2 \alpha''(\sigma) > \kappa$ , so

$$\frac{2.91/\sqrt{2\pi}}{m \sqrt{\kappa \sigma^2 \alpha''(\sigma)}} < \frac{2.91/\sqrt{2\pi}}{m\kappa} < \frac{5.82/\sqrt{2\pi}}{m} < \frac{2.3219}{m}. \quad \square$$

## Lemma 5.8

**Lemma A.16.** *For  $y \in \mathbb{R}$ ,  $\kappa \in [\frac{1}{2}, 1]$ , and  $m \geq 140$ , with  $\sigma = \sigma_{\kappa}(y)$ ,*

$$-\frac{1}{m} \frac{\partial F_{\kappa}/\partial y}{F_{\kappa}}(y, m) = \sigma_{\kappa}(y) \left( 1 + \frac{\beta}{m} \right), \quad |\beta| = |\beta(y, m, \kappa)| < 2.42.$$

*Proof.* The proof is identical to the original proof, with our Lemmas A.10 and A.15 replacing the original Lemmas 5.5 and 5.7. Only the last line of the proof changes:

$$\left| \frac{i \int_{-\infty}^{+\infty} t e^{m(\alpha(\sigma+it)-iyt)} dt}{2\pi e^{my\sigma} F_{\kappa}(y, m)} \right| < \sigma \frac{2.322}{m} \frac{1}{1 - \frac{5.23}{m}} \leq \sigma \frac{2.322}{m} \frac{1}{1 - \frac{5.23}{140}} < \sigma \frac{2.42}{m}.$$

$\square$

## A.3 Proof for remaining cases

In the remaining cases, as in the proof for  $m \geq 140$  above, we give stronger versions of Lemmas 5.5 and 5.7 to compensate for the weaker Lemma 5.6. As we have already seen

how Lemmas 5.5 and 5.7 can be strengthened by restricting to particular cases, the only remaining problem is how to prove a correct version of Lemma 5.6 that is not too much weaker than the original version. We use a computer to bound  $J_1$  by brute force, by splitting the integral up over several small intervals and estimating each one separately.

To simplify notation, define

$$\tilde{\rho}(t) = \alpha(\sigma + it) - \alpha(\sigma) - iyt = -\frac{1}{2}\alpha''(\sigma)t^2 + \rho(t).$$

Let  $0 < m_{\min} \leq m_{\max}$ ,  $0 \leq \kappa_{\min} \leq \kappa_{\max} \leq 1$ ,  $0 \leq \sigma_{\min} \leq \sigma_{\max} \leq 0.65$ , and  $0 < U_0 \leq m_{\min}\sigma_{\min}^2\alpha''_{\kappa_{\min}}(\sigma_{\min})$  be given.<sup>1</sup> We would like to find an upper bound for the integral<sup>2</sup>

$$\int_0^{\sqrt{\frac{U_0}{m\alpha''(\sigma)}}} te^{-m\operatorname{Re}(\tilde{\rho})} \sin(m\operatorname{Im}(\tilde{\rho})) dt, \quad (\text{A.2})$$

where  $m_{\min} \leq m \leq m_{\max}$ ,  $\kappa_{\min} \leq \kappa \leq \kappa_{\max}$ , and  $\sigma_{\min} \leq \sigma \leq \sigma_{\max}$ .

Note that  $m\sigma^2\alpha''_{\kappa}(\sigma)$  is increasing as a function of  $\kappa$ ,  $\sigma$ , and  $m$ ; this follows from the formula

$$\sigma^2\alpha''_{\kappa}(\sigma) = \kappa \sum_{j=0}^{\infty} \left( \frac{\sigma}{\sigma + j} \right)^2 + (1 - \kappa) \sum_{j=0}^{\infty} \left( \frac{\sigma}{\sigma + \frac{1}{2} + j} \right)^2.$$

Hence the assumption on  $U_0$  ensures that

$$\sqrt{\frac{U_0}{m\alpha''(\sigma)}} \leq \sigma. \quad (\text{A.3})$$

---

<sup>1</sup>The assumption  $\sigma_{\max} \leq 0.65$  is not necessary, and is made only to simplify the proofs. It makes sense to permit  $\sigma_{\min} = 0$ , even though  $\alpha(0)$  is not defined, because every function which we evaluate at  $\sigma_{\min}$  in the following argument will approach a limit as  $\sigma \rightarrow 0^+$ .

<sup>2</sup>More precisely, we would like to find an upper bound for the absolute value of the integral. The methods described below could be adapted in the obvious way to provide a lower bound for the integral, and therefore bound the absolute value. In practice, it's fairly clear that these integrals are always positive – at least in the cases that we care about – so we will not bother with the lower bounds here.

Once we have a bound for (A.2), we can combine it with an easy upper bound for

$$\left| \int_{\sqrt{\frac{U_0}{m\alpha''(\sigma)}}}^{\sigma} t e^{-m \operatorname{Re}(\tilde{\rho})} \sin(m \operatorname{Im}(\rho)) dt \right|$$

to obtain an upper bound for

$$\int_0^{\sigma} t e^{-m \operatorname{Re}(\tilde{\rho})} \sin(m \operatorname{Im}(\rho)) dt.$$

## Algorithm

We split the interval  $[0, \sqrt{U_0/(m\alpha''(\sigma))}]$  into a number of subintervals of the form

$$\left[ \sqrt{\frac{L}{m\alpha''(\sigma)}}, \sqrt{\frac{U}{m\alpha''(\sigma)}} \right],$$

and bound the integral over each subinterval separately. For each subinterval, we find an inequality of the form

$$\int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t e^{-m \operatorname{Re}(\tilde{\rho})} \sin(m \operatorname{Im}(\rho)) dt \leq \frac{C_{L,U}}{m\alpha''(\sigma)}, \quad (\text{A.4})$$

Adding the inequalities for each subinterval, we get an inequality

$$\int_0^{\sqrt{\frac{U_0}{m\alpha''(\sigma)}}} t e^{-m \operatorname{Re}(\tilde{\rho})} \sin(m \operatorname{Im}(\rho)) dt \leq \frac{C}{m\alpha''(\sigma)}.$$

The first step in determining  $C_{L,U}$  is to obtain upper and lower bounds for  $m \operatorname{Im}(\rho)$ .

The proof of Friedman and Skoruppa's Lemma 5.1 shows that  $m \operatorname{Im}(\rho)$  is given by the alternating series

$$m \operatorname{Im}(\rho) = m \sum_{j=1}^{\infty} (-1)^{j+1} \frac{|\alpha^{(2j+1)}(\sigma)|}{(2j+1)!} t^{2j+1},$$

and that the absolute values of the summands converge monotonically to zero.<sup>3</sup> Therefore we can obtain upper or lower bounds for  $m \operatorname{Im}(\rho)$  by truncating the series, e.g.,

$$m \operatorname{Im}(\rho) \leq m \frac{|\alpha^{(3)}(\sigma)|}{3!} t^3 - m \frac{|\alpha^{(5)}(\sigma)|}{5!} t^5 + m \frac{|\alpha^{(7)}(\sigma)|}{7!} t^7.$$

---

<sup>3</sup>This is why inequality A.3 was necessary.

Of course, we need bounds in terms of the given constants  $L$ ,  $U$ ,  $m_{\min}$ ,  $m_{\max}$ , etc. It is clear how to eliminate  $t$  and  $m$ : to continue the above example,

$$\begin{aligned} m \operatorname{Im}(\rho) &\leq \frac{|\alpha^{(3)}(\sigma)|}{\alpha''(\sigma)^{3/2}} \frac{U^{3/2}}{3!m^{1/2}} - \frac{|\alpha^{(5)}(\sigma)|}{\alpha''(\sigma)^{5/2}} \frac{L^{5/2}}{5!m^{3/2}} + \frac{|\alpha^{(7)}(\sigma)|}{\alpha''(\sigma)^{7/2}} \frac{U^{7/2}}{7!m^{5/2}} \\ &\leq \frac{|\alpha^{(3)}(\sigma)|}{\alpha''(\sigma)^{3/2}} \frac{U^{3/2}}{3!m_{\min}^{1/2}} - \frac{|\alpha^{(5)}(\sigma)|}{\alpha''(\sigma)^{5/2}} \frac{L^{5/2}}{5!m_{\max}^{3/2}} + \frac{|\alpha^{(7)}(\sigma)|}{\alpha''(\sigma)^{7/2}} \frac{U^{7/2}}{7!m_{\min}^{5/2}}. \end{aligned}$$

To handle the terms involving  $\sigma$ , we use the following lemma:

**Lemma A.17.** *Let  $0 \leq \sigma_{\min} \leq \sigma_{\max} \leq 0.65$  and  $0 \leq \kappa_{\min} \leq \kappa_{\max} \leq 1$  be given. Let  $n \geq 3$  be an integer. If  $\sigma \in [\sigma_{\min}, \sigma_{\max}]$  and  $\kappa \in [\kappa_{\min}, \kappa_{\max}]$ , then*

$$\left( \frac{\kappa_{\min}}{\kappa_{\max}} \right)^{\frac{n}{2}-1} \frac{|\alpha_{\kappa_{\min}}^{(n)}(\sigma_{\max})|}{\alpha_{\kappa_{\min}}''(\sigma_{\max})^{n/2}} \leq \frac{|\alpha_{\kappa}^{(n)}(\sigma)|}{\alpha_{\kappa}''(\sigma)^{n/2}} \leq \left( \frac{\kappa_{\max}}{\kappa_{\min}} \right)^{\frac{n}{2}-1} \frac{|\alpha_{\kappa_{\max}}^{(n)}(\sigma_{\min})|}{\alpha_{\kappa_{\max}}''(\sigma_{\min})^{n/2}}.$$

*Proof.* This is immediate from Lemmas A.20 and A.21 in the next section.  $\square$

Using the lemma to replace each  $|\alpha_{\kappa}^{(n)}(\sigma)|/\alpha_{\kappa}''(\sigma)^{n/2}$  with an upper or lower bound, we get constant bounds  $\theta_1 \leq m \operatorname{Im}(\rho) \leq \theta_2$  as desired. We can use the exact same argument to find bounds  $\eta_1 \leq -m \operatorname{Re}(\tilde{\rho}) \leq \eta_2$ .

Define

$$S = \max_{\theta_1 \leq \theta \leq \theta_2} \sin(\theta).$$

We consider two cases separately:

- If  $S \geq 0$ , then

$$\int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t e^{-m \operatorname{Re}(\tilde{\rho})} \sin(m \operatorname{Im}(\rho)) dt \leq S e^{\eta_2} \int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t dt = \frac{S e^{\eta_2} (U - L)/2}{m\alpha''(\sigma)}.$$

- If  $S < 0$ , then

$$\int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t e^{-m \operatorname{Re}(\tilde{\rho})} \sin(m \operatorname{Im}(\rho)) dt \leq S e^{\eta_1} \int_{\sqrt{\frac{L}{m\alpha''(\sigma)}}}^{\sqrt{\frac{U}{m\alpha''(\sigma)}}} t dt = \frac{S e^{\eta_1} (U - L)/2}{m\alpha''(\sigma)}.$$

In either case, we have an inequality of the form (A.4), as claimed.

It remains only to bound the integral over  $[\sqrt{U_0/(m\alpha''(\sigma))}, \sigma]$ . The estimate relies on the following bound, which will be proven in the last section.

**Lemma A.18.** *Assume<sup>4</sup>  $\sigma_{\min} \leq 0.65$ . For all  $|t| \leq \sigma$ , we have*

$$-m \operatorname{Re}(\tilde{\rho}(t)) \leq -m\beta\alpha''(\sigma)t^2,$$

where

$$\beta = \frac{1}{2} - \frac{\sigma_{\min}^2 \Psi^{(3)}(\sigma_{\min})}{24\Psi'(\sigma_{\min})}.$$

Define  $V_0 = m_{\max}\sigma_{\max}^2\alpha''_{\kappa_{\max}}(\sigma_{\max})$ . Recall that  $m\sigma^2\alpha''_{\kappa}(\sigma)$  is strictly increasing as a function of  $m$ ,  $\kappa$ , and  $\sigma$ , so

$$\sigma \leq \sqrt{\frac{m_{\max}\sigma_{\max}^2\alpha''_{\kappa_{\max}}(\sigma_{\max})}{m\alpha''(\sigma)}} = \sqrt{\frac{V_0}{m\alpha''(\sigma)}}.$$

Hence

$$\begin{aligned} \left| \int_{\sqrt{\frac{U_0}{m\alpha''(\sigma)}}}^{\sigma} t e^{-m \operatorname{Re}(\tilde{\rho})} \sin(m \operatorname{Im}(\rho)) dt \right| &\leq \int_{\sqrt{\frac{U_0}{m\alpha''(\sigma)}}}^{\sqrt{\frac{V_0}{m\alpha''(\sigma)}}} t e^{-m\beta\alpha''(\sigma)t^2} dt \\ &= \frac{.5(e^{-\beta U_0} - e^{-\beta V_0})\sqrt{m\kappa}/\beta}{\sqrt{\kappa}m^{3/2}\alpha''(\sigma)} \\ &\leq \frac{.5(e^{-\beta U_0} - e^{-\beta V_0})\sqrt{m_{\max}\kappa_{\max}}/\beta}{\sqrt{\kappa}m^{3/2}\alpha''(\sigma)}. \end{aligned}$$

## Proof of Lemma A.17

Let  $\zeta$  denote the Hurwitz zeta function.

**Lemma A.19.** *For any integer  $n \geq 3$  and any  $\sigma \in (0, 0.65]$ ,*

$$\zeta(n+1, \sigma)\zeta(2, \sigma + \tfrac{1}{2}) + \zeta(n+1, \sigma + \tfrac{1}{2})\zeta(2, \sigma) - \zeta(n, \sigma)\zeta(3, \sigma + \tfrac{1}{2}) - \zeta(n, \sigma + \tfrac{1}{2})\zeta(3, \sigma) > 0.$$

---

<sup>4</sup>Again, this assumption is unnecessary.

*Proof.* Estimating the zeta functions by integrals, we obtain the following bounds:

$$\begin{aligned}
\zeta(n+1, \sigma) &> \frac{1}{\sigma^{n+1}} + \frac{1}{(\sigma+1)^{n+1}}, \\
\zeta(2, \sigma + \tfrac{1}{2}) &> \frac{1}{(\sigma + \tfrac{1}{2})^2} + \frac{1}{(\sigma + \tfrac{3}{2})^2} + \frac{1}{(\sigma + \tfrac{5}{2})^2}, \\
\zeta(n+1, \sigma + \tfrac{1}{2}) &> \frac{1}{(\sigma + \tfrac{1}{2})^{n+1}} + \frac{1}{(\sigma + \tfrac{3}{2})^{n+1}}, \\
\zeta(2, \sigma) &> \frac{1}{\sigma^2} + \frac{1}{(\sigma+1)^2}, \\
\zeta(n, \sigma) &< \frac{1}{\sigma^n} + \frac{1}{(n-1)\sigma^{n-1}}, \\
\zeta(3, \sigma + \tfrac{1}{2}) &< \frac{1}{(\sigma + \tfrac{1}{2})^3} + \frac{1}{(\sigma + \tfrac{3}{2})^3} + \frac{1}{2(\sigma + \tfrac{5}{2})^2}, \\
\zeta(n, \sigma + \tfrac{1}{2}) &< \frac{1}{(\sigma + \tfrac{1}{2})^n} + \frac{1}{(\sigma + \tfrac{3}{2})^n} + \frac{1}{(n-1)(\sigma + \tfrac{5}{2})^{n-1}}, \\
\zeta(3, \sigma) &< \frac{1}{\sigma^3} + \frac{1}{(\sigma+1)^3} + \frac{1}{2(\sigma+1)^2}.
\end{aligned}$$

Combining these bounds, we obtain a lower bound for the target function. When  $n = 3$ , the bound is

$$\frac{p(\sigma)}{\sigma^4(\sigma+1)^4(2\sigma+1)^4(2\sigma+3)^4(5+2\sigma)^2},$$

where

$$\begin{aligned}
p(\sigma) = & -512\sigma^{14} - 10240\sigma^{13} - 80640\sigma^{12} - 354048\sigma^{11} - \\
& 995520\sigma^{10} - 1914176\sigma^9 - 2571008\sigma^8 - 2348240\sigma^7 - 1255858\sigma^6 - \\
& 35300\sigma^5 + 591167\sigma^4 + 558684\sigma^3 + 275655\sigma^2 + 75852\sigma + 9324.
\end{aligned}$$

We can see that this is positive for  $0 \leq \sigma \leq 0.65$ . [Proof: By Descartes' rule of signs,  $p(\sigma)$  changes sign only once for  $\sigma > 0$ , and  $p(0), p(0.65) > 0$ .] This proves the claim when  $n = 3$ .

We can prove the claim similarly when  $n = 4, 5, 6, 7, 8$ .

Now assume  $n \geq 9$ . In this case, we will use the bounds

$$\begin{aligned}
\zeta(n+1, \sigma) &> \frac{1}{\sigma^{n+1}}, & \zeta(2, \sigma + \tfrac{1}{2}) &> \frac{1}{(\sigma + \frac{1}{2})^2}, \\
\zeta(n+1, \sigma + \tfrac{1}{2}) &> \frac{1}{(\sigma + \frac{1}{2})^{n+1}}, & \zeta(2, \sigma) &> \frac{1}{\sigma^2}, \\
\zeta(n, \sigma) &< \frac{1}{\sigma^n} + \frac{1}{(n-1)\sigma^{n-1}} \leq \frac{1}{\sigma^n} + \frac{1}{8\sigma^{n-1}}, & \zeta(3, \sigma + \tfrac{1}{2}) &< \frac{1}{(\sigma + \frac{1}{2})^3} + \frac{1}{2(\sigma + \frac{1}{2})^2}, \\
\zeta(n, \sigma + \tfrac{1}{2}) &< \frac{1}{(\sigma + \frac{1}{2})^n} + \frac{1}{8(\sigma + \frac{1}{2})^{n-1}}, & \zeta(3, \sigma) &< \frac{1}{\sigma^3} + \frac{1}{2\sigma^2}.
\end{aligned}$$

We obtain a lower bound of

$$\frac{(1 + \frac{1}{2\sigma})^{n-2} (32 - 16\sigma - 42\sigma^2 - 4\sigma^3) - (34 + 25\sigma + 44\sigma^2 + 4\sigma^3)}{64\sigma^3(\sigma + \frac{1}{2})^{n+1}}. \quad (\text{A.5})$$

For  $\sigma \in (0, 0.65]$  and  $n \geq 9$ , we have

$$\left(1 + \frac{1}{2\sigma}\right)^{n-2} \geq \left(1 + \frac{1}{2 \cdot 0.65}\right)^7 > 54.$$

Note that  $32 - 16\sigma - 42\sigma^2 - 4\sigma^3 > 0$  for  $0 \leq \sigma \leq 0.65$ ; thus the numerator in (A.5) is bounded below by

$$54(32 - 16\sigma - 42\sigma^2 - 4\sigma^3) - (34 + 25\sigma + 44\sigma^2 + 4\sigma^3) = 1694 - 889\sigma - 2312\sigma^2 - 220\sigma^3.$$

We can check that  $1694 - 889\sigma - 2312\sigma^2 - 220\sigma^3 > 0$  for  $0 \leq \sigma \leq 0.65$ . This completes the proof.  $\square$

**Lemma A.20.** *For any  $n \geq 3$  and  $\kappa \in [0, 1]$ , the function*

$$\frac{|\alpha_\kappa^{(n)}(\sigma)|}{\alpha_\kappa''(\sigma)^{n/2}}$$

*is decreasing for  $\sigma \in (0, 0.65]$ .*

*Proof.* We have

$$\frac{d}{d\sigma} \frac{|\alpha_\kappa^{(n)}(\sigma)|}{\alpha_\kappa''(\sigma)^{n/2}} = \frac{d}{d\sigma} (-1)^n \frac{\alpha_\kappa^{(n)}(\sigma)}{\alpha_\kappa''(\sigma)^{n/2}} = (-1)^n \frac{\alpha_\kappa^{(n+1)}(\sigma) \alpha_\kappa''(\sigma) - \frac{n}{2} \alpha_\kappa^{(n)}(\sigma) \alpha_\kappa^{(3)}(\sigma)}{\alpha_\kappa''(\sigma)^n},$$

so we need to show that

$$(-1)^n [\alpha_\kappa^{(n+1)}(\sigma) \alpha_\kappa''(\sigma) - \frac{n}{2} \alpha_\kappa^{(n)}(\sigma) \alpha_\kappa^{(3)}(\sigma)] \leq 0.$$

Plugging in the definition of  $\alpha_\kappa$  and using the formula  $\Psi^{(k)}(s) = (-1)^{k+1} k! \zeta(k+1, s)$ , this becomes

$$\begin{aligned} & -[\kappa \zeta(n+1, \sigma) + (1-\kappa) \zeta(n+1, \sigma + \tfrac{1}{2})][\kappa \zeta(2, \sigma) + (1-\kappa) \zeta(2, \sigma + \tfrac{1}{2})] + \\ & [\kappa \zeta(n, \sigma) + (1-\kappa) \zeta(n, \sigma + \tfrac{1}{2})][\kappa \zeta(3, \sigma) + (1-\kappa) \zeta(3, \sigma + \tfrac{1}{2})] \leq 0. \end{aligned}$$

The left-hand side is the sum of the following three expressions:

$$\begin{aligned} & \kappa^2 [-\zeta(n+1, \sigma) \zeta(2, \sigma) + \zeta(n, \sigma) \zeta(3, \sigma)], \\ & (1-\kappa)^2 [-\zeta(n+1, \sigma + \tfrac{1}{2}) \zeta(2, \sigma + \tfrac{1}{2}) + \zeta(n, \sigma + \tfrac{1}{2}) \zeta(3, \sigma + \tfrac{1}{2})], \\ & -\kappa(1-\kappa) [\zeta(n+1, \sigma) \zeta(2, \sigma + \tfrac{1}{2}) + \zeta(n+1, \sigma + \tfrac{1}{2}) \zeta(2, \sigma) - \zeta(n, \sigma) \zeta(3, \sigma + \tfrac{1}{2}) - \zeta(n, \sigma + \tfrac{1}{2}) \zeta(3, \sigma)]. \end{aligned}$$

I claim that all three are non-positive. For the first two, this follows from log-convexity of  $k \mapsto \zeta(k, \sigma)$ ; the third claim is Lemma A.19.  $\square$

**Lemma A.21.** *Let  $0 < \sigma \leq 0.65$  and  $n \geq 3$  be given. Then*

$$\kappa^{\frac{n}{2}-1} \frac{|\alpha_\kappa^{(n)}(\sigma)|}{\alpha_\kappa''(\sigma)^{n/2}}.$$

*is increasing for  $\kappa \in [0, 1]$ .*

*Proof.* Set

$$\begin{aligned} t &= \frac{1-\kappa}{\kappa} \geq 0, \quad A = \Psi'(\sigma), \quad B = \Psi'(\sigma + \tfrac{1}{2}), \\ C &= |\Psi^{(n-1)}(\sigma)|, \quad D = |\Psi^{(n-1)}(\sigma + \tfrac{1}{2})|. \end{aligned}$$



With this notation, we want to show that

$$\frac{C + tD}{(A + tB)^{n/2}}$$

is decreasing as a function of  $t$  for all  $t \geq 0$ . We have

$$\frac{d}{dt} \frac{C + tD}{(A + tB)^{n/2}} = \frac{AD - \frac{n}{2}BC - (\frac{n}{2} - 1)BDt}{(A + tB)^{\frac{n}{2}+1}}.$$

Clearly  $-(\frac{n}{2} - 1)BDt \leq 0$ , so we need only check that  $AD - \frac{n}{2}BC < 0$ . Rewriting  $A$ ,  $B$ ,  $C$ , and  $D$  in terms of  $\zeta$ , we want to show that

$$\zeta(2, \sigma)\zeta(n, \sigma + \frac{1}{2}) - \frac{n}{2}\zeta(2, \sigma + \frac{1}{2})\zeta(n, \sigma) < 0.$$

We have

$$\begin{aligned} & \zeta(2, \sigma)\zeta(n, \sigma + \frac{1}{2}) - \frac{n}{2}\zeta(2, \sigma + \frac{1}{2})\zeta(n, \sigma) \\ & < \left(\frac{1}{\sigma^2} + \frac{1}{\sigma}\right) \left(\frac{1}{(\sigma + \frac{1}{2})^n} + \frac{1}{(n-1)(\sigma + \frac{1}{2})^{n-1}}\right) - \frac{n}{2} \frac{1}{(\sigma + \frac{1}{2})^2} \cdot \frac{1}{\sigma^n} \\ & = \frac{1}{(\sigma + \frac{1}{2})^{n+2}} \left[ \left(\frac{1}{\sigma^2} + \frac{1}{\sigma}\right) (\sigma + \frac{1}{2})^2 \left(1 + \frac{\sigma + \frac{1}{2}}{n-1}\right) - \frac{n}{2} \left(1 + \frac{1}{2\sigma}\right)^n \right]. \end{aligned}$$

The quantity in brackets is decreasing as a function of  $n$ , so it suffices to check the case  $n = 3$ . When  $n = 3$ , this quantity is

$$\frac{(2\sigma + 1)^2(2\sigma^3 + 7\sigma^2 - \sigma - 3)}{16\sigma^3}.$$

We can see that this is negative when  $0 \leq \sigma \leq 0.65$ . □

## Proof of Lemma A.18

**Lemma A.22.** *Let  $\sigma_{min} \in (0, 0.65]$  be given. For all  $\sigma \geq \sigma_{min}$ ,*

$$\sigma^2 \alpha^{(4)}(\sigma) \leq \frac{\sigma_{min}^2 \Psi^{(3)}(\sigma_{min})}{\Psi'(\sigma_{min})} \alpha''(\sigma).$$

*Proof.* Lemma A.5 says that

$$\sigma^2 \Psi^{(3)}(\sigma) \leq \frac{\sigma_{\min}^2 \Psi^{(3)}(\sigma_{\min})}{\Psi'(\sigma_{\min})} \Psi'(\sigma),$$

$$\sigma^2 \Psi^{(3)}(\sigma + \tfrac{1}{2}) < (\sigma + \tfrac{1}{2})^2 \Psi^{(3)}(\sigma + \tfrac{1}{2}) \leq \frac{\sigma_{\min}^2 \Psi^{(3)}(\sigma_{\min})}{\Psi'(\sigma_{\min})} \Psi'(\sigma + \tfrac{1}{2}),$$

and combining these inequalities yields the lemma.  $\square$

Lemma A.18 is an immediate consequence:

$$\begin{aligned} \operatorname{Re}(\tilde{\rho}(t)) &\leq -\frac{1}{2} \alpha''(\sigma) t^2 + \frac{1}{24} \alpha^{(4)}(\sigma) t^4 \\ &= \left( -\frac{1}{2} + \frac{1}{24} \frac{\alpha^{(4)}(\sigma)}{\alpha''(\sigma)} t^2 \right) \alpha''(\sigma) t^2 \\ &\leq \left( -\frac{1}{2} + \frac{1}{24} \frac{\sigma^2 \alpha^{(4)}(\sigma)}{\alpha''(\sigma)} \right) \alpha''(\sigma) t^2 \\ &\leq \left( -\frac{1}{2} + \frac{\sigma_{\min}^2 \Psi^{(3)}(\sigma_{\min})}{24 \Psi'(\sigma_{\min})} \right) \alpha''(\sigma) t^2. \end{aligned}$$

# Bibliography

- [1] Milton Abramowitz and Irene A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [2] Harry Bateman and Arthur Erdlyi. *Higher transcendental functions*. Bateman Manuscript Project. McGraw-Hill, New York, NY, 1955.
- [3] A.-M. Bergé and J. Martinet. Minorations de hauteurs et petits régulateurs relatifs. In *Séminaire de Théorie des Nombres, 1987–1988 (Talence, 1987–1988)*, pages Exp. No. 11, 28. Univ. Bordeaux I, Talence, 1987.
- [4] A.-M. Bergé and J. Martinet. Notions relatives de régulateurs et de hauteurs. *Acta Arith.*, 54(2):155–170, 1989.
- [5] A.-M. Bergé and J. Martinet. Sur les minorations géométriques des régulateurs. In *Séminaire de Théorie des Nombres, Paris 1987–88*, volume 81 of *Progr. Math.*, pages 23–50. Birkhäuser Boston, Boston, MA, 1990.

- [6] Antone Costa and Eduardo Friedman. Ratios of regulators in totally real extensions of number fields. *J. Number Theory*, 37(3):288–297, 1991.
- [7] N. G. de Bruijn. *Asymptotic methods in analysis*. Bibliotheca Mathematica. Vol. 4. North-Holland Publishing Co., Amsterdam; P. Noordhoff Ltd., Groningen; Interscience Publishers Inc., New York, 1958.
- [8] Eduardo Friedman and Nils-Peter Skoruppa. Relative regulators of number fields. *Invent. Math.*, 135(1):115–144, 1999.