

That Little Matter of Consciousness

Martha Farah, University of Pennsylvania

Examples of all three approaches can be found in Table 1 of the Target Article. One approach is to show preserved high-level cognitive processing by patients' brains, of the kind normally accompanied by consciousness. This approach is exemplified by the study of Schiff et al. (2005). Patients in MCS were scanned while recordings of either meaningful speech or backwards speech were played to them. The difference in brain activation in response to forward and backward speech was used as a measure of their brains' processing of meaningful speech per se; that is, without auditory processing that is common to both meaningful and meaningless speech sounds. Surprisingly, the MCS patients

showed patterns of brain activity that were qualitatively similar to those evoked in normal healthy subjects, suggesting that large-scale networks underlying language comprehension were to some degree preserved in these severely brain-damaged individuals. Although the authors of the study were careful to interpret these results as indicating preserved language function but not necessarily consciousness, the results have been described elsewhere as indicative of consciousness (Carey, 2005) and the finding is discussed in most reviews of consciousness in VS and MCS. The implicit assumption here is that higher cognitive processes, such as those involved in speech recognition, cannot be carried out unconsciously, and so evidence of these cognitive processes is evidence of consciousness. This assumption is supported by everyday experience, but it is not true under all circumstances. As pointed out by Levy (2008), the cognitive psychology literature contains many examples of dissociated cognition and awareness. More to the point, brain damage can lead to just this type of dissociation (Farah et al. 1993). In short, *cognition* is not the same as *consciousness*. To demonstrate consciousness, other approaches are needed.

A second approach to detecting consciousness in severely brain-damaged patients makes use of previous cognitive neuroscience research on the neural correlates of consciousness. It involves showing preservation of patterns of activity that have been demonstrated, in previous research, to distinguish conscious from unconscious processing. This approach is exemplified by Boly and colleagues (2004) listed in Table 1 of the target article. In this study, investigators used simple clicks as probes to activate the brain and looked to see what parts of the brain became active and whether the activity in these different parts was intercorrelated. The implicit assumption here is that consciousness arises with certain patterns of brain activity — perhaps activity in certain regions, such as medial frontal and parietal areas, or activity that is correlated across certain brain areas. Such an association between patterns of activation and consciousness would have to be demonstrated initially by experimental manipulations of consciousness in normal brains and then used as a signature of consciousness in the damaged brain. Boly et al. sought patterns of activation like those demonstrated in other studies to mark conscious auditory perception and found them in MCS but not VS patients. A weakness of this approach comes from the paucity of brain imaging studies in which conscious and unconscious processing have been directly compared. To advance research using this approach, we need more such studies, using different subject populations and different methods of manipulating conscious awareness of stimuli, to provide a more general and reliable “brain signature” of consciousness.

The third and final approach that has so far been taken is to use brain activity as a surrogate for overt behavior in examining patients for consciousness. This was used by Owen and colleagues (2006) to demonstrate command following in a patient who met criteria for VS. The commands in this case were to imagine playing tennis or taking a walk through the rooms of one’s home, and the patient’s compliance with these commands was demonstrated by patterns

of brain activity characteristic of imagined motor activity and imagined navigation. The implicit assumption for this approach is that such commands cannot be followed without conscious awareness. If it were the case that hearing the request to imagine playing tennis could automatically and unconsciously trigger motor imagery, then Owen et al.’s findings would not be evidence of consciousness. However, this seems implausible. After all, when a patient squeezes the examiner’s hand on request, we take that as evidence of consciousness and do not ask whether the squeeze could have been triggered unconsciously. Nevertheless, this approach could be strengthened by devising commands that result in recognizable patterns of brain activity but are even less plausibly attributed to automatic and unconscious associations with specific words or phrases.

In sum, three research strategies that have been used so far to detect consciousness in severely brain-damaged patients. One is based on an incorrect assumption about the relation between consciousness and cognition, and these studies should not be used for evidence concerning patients’ conscious awareness. In contrast, the other two strategies hold promise. Their potential can be more fully realized by further research on the neural correlates of consciousness in the healthy brain and the development of new protocols using brain activity as a surrogate for behavioral responses.

REFERENCES

- Bauer, G., Gerstenbrand F., and Rimpl E. 1979. Varieties of the locked-in syndrome. *Journal of Neurology* 221: 77–91.
- Boly, M., Faymonville M. E., Peigneux P., Lambermont B., Damas P., Del Fiore G., Degueldre C., Franck G., Luxen A., Lamy M., Moonen G., Maquet P., and Laureys S. 2004. Auditory processing in severely brain injured patients: Differences between the minimally conscious state and the persistent vegetative state. *Archives of Neurology* 6(2): 233–238.
- Carey, B. February 8, 2005. New signs of awareness seen in some brain-injured patients. *New York Times*.
- Farah, M. J. 2008. Neuroethics and the problem of other minds: Implications of neuroscience evidence for the moral status of brain-damaged patients and nonhuman animals. *Neuroethics* 1: 9–18.
- Farah, M. J., O’Reilly, R. C., and Vecera, S. P. 1993. Dissociated overt and covert recognition as an emergent property of a lesioned neural network. *Psychological Review* 100: 571–588.
- Fins, J. J., Illes, J., Bernat, J. L., Hirsch, J., Laureys, S., and Murphy, E. R. 2008. Consciousness, imaging, ethics, and the injured brain. *American Journal of Bioethics (AJOB- Neuroscience)* 8(9): x–xx.
- Levy, N. 2008. Going beyond the evidence. *American Journal of Bioethics (AJOB-Neuroscience)* 8(9): x–xx.
- Owen, A. M., Coleman, M. R., Boly, M., Davis, M. H., Laureys, S., and Pickard, J. D. 2006. Detecting awareness in the vegetative state. *Science* 313(5792): 1402.
- Schiff, N. D., Rodriguez-Moreno, D., Kamal, A., Kim, K. H., Giacino, J. T., Plum, F., and Hirsch, J. 2005. fMRI reveals large-scale network activation in minimally conscious patients. *Neurology* 64(3): 514–523.

Queries

- Q1. Au: Please provide pages for quotes.
- Q2. Au: Pls. provide Table 1.
- Q3. Au: of the target article or of Boly et al.?
- Q4. Au: Please provide page.