



2015

Trivalent Influenza Vaccine Adverse Event Analysis Based On MedDRA System Organ Classes Using VAERS Data

Cui Tao

Jingcheng Du

Yi Cai

Yong Chen

University of Pennsylvania, ychen123@mail.med.upenn.edu

Follow this and additional works at: <https://repository.upenn.edu/epidemiology>

 Part of the [Epidemiology Commons](#)

Tao, Cui; Du, Jingcheng; Cai, Yi; and Chen, Yong, "Trivalent Influenza Vaccine Adverse Event Analysis Based On MedDRA System Organ Classes Using VAERS Data" (2015). *Department of Biostatistics, Epidemiology and Informatics*. 7.

<https://repository.upenn.edu/epidemiology/7>

Cui Tao, Jingcheng Du, Yi Cai, Yong Chen. (2015). Trivalent Influenza Vaccine Adverse Event Analysis Based On MedDRA System Organ Classes Using VAERS Data. In I.N. Sarkar et al. (Eds.), *MEDINFO 2015: eHealth-enabled Health* (p. 1076). [dx.doi.org/10.3233/978-1-61499-564-7-1076](https://doi.org/10.3233/978-1-61499-564-7-1076)

This article is published online with Open Access by IOS Press and distributed under the terms of the Creative Commons Attribution Non-Commercial License.

This paper is posted at ScholarlyCommons. <https://repository.upenn.edu/epidemiology/7>

For more information, please contact repository@pobox.upenn.edu.

Trivalent Influenza Vaccine Adverse Event Analysis Based On MedDRA System Organ Classes Using VAERS Data

Abstract

We studied serious reports following influenza vaccine from VAERS database in year 2011. Our statistical analyses revealed differences of reactions among different age groups and between genders. The results may lead to additional studies to uncover factors contributing to the individual differences in susceptibility to influenza infection.

Keywords

Influnza vaccine adverse event, MedDRA, VAERS

Disciplines

Epidemiology | Medicine and Health Sciences | Public Health

Comments

Cui Tao, Jingcheng Du, Yi Cai, Yong Chen. (2015). Trivalent Influenza Vaccine Adverse Event Analysis Based On MedDRA System Organ Classes Using VAERS Data. In I.N. Sarkar et al. (Eds.), *MEDINFO 2015: eHealth-enabled Health* (p. 1076). [dx.doi.org/10.3233/978-1-61499-564-7-1076](https://doi.org/10.3233/978-1-61499-564-7-1076)

This article is published online with Open Access by IOS Press and distributed under the terms of the Creative Commons Attribution Non-Commercial License.

Trivalent Influenza Vaccine Adverse Event Analysis Based On MedDRA System Organ Classes Using VAERS Data

Cui Tao^a, Jingcheng Du^a, Yi Cai^b, Yong Chen^b

^aSchool of Biomedical Informatics, ^bSchool of Public Health
The University of Texas Health Science Center at Houston, TX, USA

Abstract

We studied serious reports following influenza vaccine from VAERS database in year 2011. Our statistical analyses revealed differences of reactions among different age groups and between genders. The results may lead to additional studies to uncover factors contributing to the individual differences in susceptibility to influenza infection.

Keywords:

Influnza vaccine adverse event, MedDRA, VAERS

Introduction

Seasonal influenza is a common vaccine-preventable disease with substantial morbidity. The social burden of seasonal flu can be substantial [1]. Annual vaccination is the most effective strategy to prevent influenza [2]. While their benefits far overweigh their risks and costs, influenza vaccines are accompanied with specific adverse events. Post-approval surveillance of vaccine adverse events is critically needed to assess the vaccine safety throughout its life on the market. The Vaccine Adverse Event Reporting System (VAERS) is a passive surveillance system to monitor vaccine safety after the administration of vaccines licensed in the United States [3]. In this study, we explored statistical analysis on annotated symptoms in the VAERS reports for patients with different genders and ages.

Materials and Methods

We searched the VAERS for US reports after Trivalent Influenza Vaccine (FLU3) in year 2011 and extracted serious reports (i.e., death, life-threatening illness, hospitalization, prolonged hospitalization, or permanent disability). For each report, the VAERS provides annotations for post-vaccination symptoms in Medical Dictionary for Regulatory Activities (MedDRA) terms [4]. To facilitate further statistical analysis, we further grouped these symptoms based on the MedDRA System Organ Class (SOC) using the NCBO Web Services [5].

To model the total number of serious symptoms per subject, we fit a zero-truncated Poisson regression on age groups and gender because a subject has to have at least one symptom to be included in the VAERS database. To study the risks of having serious symptom for each SOC type, we use logistic regression on age groups and gender.

Results

During the study period, VAERS received 7986 FLU3 reports, 638 were serious. Out of the 638 reports, 324 were for female patients, 295 were for male patients, and 134, 156, 110, 185 were for patients in age groups 0.5-17, 17-49, 49-64, and >64 respectively. 5407 symptoms were grouped into 26 SOCs. The most frequent SOCs in the 638 reports are nervous system disorders, general disorders, and administration site condition and investigations.

Analysis using zero-truncated Poisson model indicated that the average number of symptoms per subjects in the study cohort is 8.74 (95% CI 6.76, 10.73). There are statistically significant differences in number of symptoms among four age groups and between different genders. The youngest age group (0.5 - 17 years) has the smallest number of symptoms per year, followed by age group 2 (17-49), age group 4 (>64), and finally age group 3 (49 - 64). The average number of symptoms for subjects of 17-49 years old is 13% higher than the average number of symptoms for subjects of 0.5-17 years old with the same gender ($p=0.003$).

Conclusion

This poster reports our preliminary analyses on influenza vaccine adverse events using VAERS data and MedDRA SOCs, which revealed differences of reactions among different age groups and between genders. The results may lead to additional studies to uncover factors contributing to the individual differences in susceptibility to influenza infection.

Acknowledgements

This research is partially supported by the National Library Of Medicine of the National Institutes of Health under Award Number R01LM011829 to Dr. Cui Tao.

References

1. Poehling KA, Edwards KM, Griffin MR, Szilagyi PG, Staat MA, Iwane MK, et al. The burden of influenza in young children, 2004-2009. *Pediatrics*. 2013; **131**(2): 207-16.
2. CDC: Children, the Flu, and the Flu Vaccine. [cited; Available from: <http://www.cdc.gov/flu/protect/children.htm>
3. The U.S. FDA/CDC Vaccine Adverse Event Reporting System (VAERS) [cited; Available from: <http://vaers.hhs.gov/index>
4. Brown EG, Wood L, Wood S. The medical dictionary for regulatory activities (MedDRA). *Drug safety*. 1999; **20**(2): 109-17.
5. Whetzel PL. NCBO Technology: Powering semantically aware applications. *Journal of biomedical semantics*. 2013; **4** **Suppl 1**: S8.