



April 2005

## Hypoxia-inducible Factor Regulates $\alpha v \beta 3$ Integrin Cell Surface Expression

Karen D. Cowden Dahl  
*University of Pennsylvania*

Sarah E. Robertson  
*University of Pennsylvania*

Valerie M. Weaver  
*University of Pennsylvania, vmweaver@seas.upenn.edu*

M. Celeste Simon  
*University of Pennsylvania, celeste2@mail.med.upenn.edu*

Follow this and additional works at: [https://repository.upenn.edu/ime\\_papers](https://repository.upenn.edu/ime_papers)

---

Cowden Dahl, Karen D.; Robertson, Sarah E.; Weaver, Valerie M.; and Simon, M. Celeste, "Hypoxia-inducible Factor Regulates  $\alpha v \beta 3$  Integrin Cell Surface Expression" (2005). *Institute for Medicine and Engineering Papers*. 31.

[https://repository.upenn.edu/ime\\_papers/31](https://repository.upenn.edu/ime_papers/31)

Reprinted from *Molecular Biology of the Cell*, Volume 16, Issue 4, April 2005, pages 1901-12.  
Publisher URL: <http://www.molbiolcell.org/cgi/reprint/16/4/1901.pdf>

This paper is posted at ScholarlyCommons. [https://repository.upenn.edu/ime\\_papers/31](https://repository.upenn.edu/ime_papers/31)  
For more information, please contact [repository@pobox.upenn.edu](mailto:repository@pobox.upenn.edu).

---

## Hypoxia-inducible Factor Regulates $\alpha v \beta 3$ Integrin Cell Surface Expression

### Abstract

Hypoxia-inducible factor (HIF)-deficient placentas exhibit a number of defects, including changes in cell fate adoption, lack of fetal angiogenesis, hypocellularity, and poor invasion into maternal tissue. HIF is a heterodimeric transcription factor consisting of  $\alpha$  and  $\beta$  aryl hydrocarbon receptor nuclear translocator or ARNT) subunits. We used undifferentiated trophoblast stem (TS) cells to characterize HIF-dependent adhesion, migration, and invasion.  $Arnt^{-/-}$  and  $Hif\alpha^{-/-}$  TS cells exhibit reduced adhesion and migration toward vitronectin compared with wild-type cells. Furthermore, this defect is associated with decreased cell surface expression of integrin  $\alpha v \beta 3$  and significantly decreased expression of this integrin in focal adhesions. Because of the importance of adhesion and migration in tumor progression (in addition to placental development), we examined the affect of culturing B16F0 melanoma cells in 1.5% oxygen ( $O_2$ ). Culturing B16F0 melanoma cells at 1.5%  $O_2$  resulted in increased  $\alpha v \beta 3$  integrin surface expression and increased adhesion to and migration toward vitronectin. Together, these data suggest that HIF and  $O_2$  tension influence placental invasion and tumor migration by increasing cell surface expression of  $\alpha v \beta 3$  integrin.

### Comments

Reprinted from *Molecular Biology of the Cell*, Volume 16, Issue 4, April 2005, pages 1901-12.  
Publisher URL: <http://www.molbiolcell.org/cgi/reprint/16/4/1901.pdf>

# Hypoxia-inducible Factor Regulates $\alpha\beta3$ Integrin Cell Surface Expression

Karen D. Cowden Dahl,\* Sarah E. Robertson,<sup>†</sup> Valerie M. Weaver,<sup>‡§</sup> and M. Celeste Simon\*<sup>||</sup>

\*Abramson Family Cancer Research Institute; Departments of <sup>†</sup>Cell and Developmental Biology, <sup>‡</sup>Pathology and Laboratory Medicine, and <sup>§</sup>Bioengineering, Institute for Medicine and Engineering; and <sup>||</sup>Howard Hughes Medical Institute, School of Medicine, University of Pennsylvania, Philadelphia, PA 19104

Submitted December 15, 2004; Revised January 24, 2005; Accepted January 25, 2005  
Monitoring Editor: Marianne Bronner-Fraser

Hypoxia-inducible factor (HIF)-deficient placentas exhibit a number of defects, including changes in cell fate adoption, lack of fetal angiogenesis, hypocellularity, and poor invasion into maternal tissue. HIF is a heterodimeric transcription factor consisting of  $\alpha$  and  $\beta$  aryl hydrocarbon receptor nuclear translocator or ARNT) subunits. We used undifferentiated trophoblast stem (TS) cells to characterize HIF-dependent adhesion, migration, and invasion. *Arnt*<sup>-/-</sup> and *Hif1 $\alpha$* <sup>-/-</sup> TS cells exhibit reduced adhesion and migration toward vitronectin compared with wild-type cells. Furthermore, this defect is associated with decreased cell surface expression of integrin  $\alpha\beta3$  and significantly decreased expression of this integrin in focal adhesions. Because of the importance of adhesion and migration in tumor progression (in addition to placental development), we examined the effect of culturing B16F0 melanoma cells in 1.5% oxygen (O<sub>2</sub>). Culturing B16F0 melanoma cells at 1.5% O<sub>2</sub> resulted in increased  $\alpha\beta3$  integrin surface expression and increased adhesion to and migration toward vitronectin. Together, these data suggest that HIF and O<sub>2</sub> tension influence placental invasion and tumor migration by increasing cell surface expression of  $\alpha\beta3$  integrin.

## INTRODUCTION

Oxygen (O<sub>2</sub>) deprivation or “hypoxia” acting through hypoxia-inducible factors (HIFs) influences cell adhesion and migration in a number of biological contexts. For example, HIF regulates progenitor cell adhesion and migration in ischemic tissue by inducing expression of SDF-1 and promotes tumor invasion through the tyrosine kinase receptor MET (Pennacchietti *et al.*, 2003; Ceradini *et al.*, 2004). *Hif1 $\alpha$* <sup>-/-</sup> macrophages exhibit poor migration and Matrigel invasion (Cramer *et al.*, 2003), and directed migration of renal cell carcinoma cells is regulated by HIF activation of the chemokine receptor CXCR4 (Staller *et al.*, 2003). HIF activity also is clearly important during placentation (Kozak *et al.*, 1997; Adelman *et al.*, 2000): placentas from embryos lacking HIF exhibit a variety of defects, including poor invasion of the maternal decidua (Adelman *et al.*, 2000; Cowden Dahl, Mack, Compennolle, Adelman, Carmeliet, and Simon, unpublished data). Furthermore, tumor hypoxia is often clinically correlated with increased metastatic potential (Vaupel *et al.*, 2001). The integrin adhesion receptors promote placental invasion, macrophage adhesion and migration, and metastatic progression of neoplasms (Zhou *et al.*, 1997b; Ding *et al.*, 1999; Jin and Varner, 2004), and hypoxia has been shown to increase mRNA levels of multiple integrins ( $\beta3$ ,  $\beta2$ , and  $\alpha5$ ) (Walton *et al.*, 2000; Iwaki *et al.*, 2004; Koike *et al.*, 2004; Kong *et al.*, 2004). Therefore, HIF

regulates cell motility in a variety of developmental and pathological situations.

The transcriptional response to hypoxia is mediated by the dimeric basic-helix-loop-helix-PAS (bHLH-PAS) transcription factor HIF. HIF is composed of an  $\alpha$  subunit (HIF1 $\alpha$  or HIF2 $\alpha$ ) and  $\beta$  subunit (aryl hydrocarbon receptor nuclear translocator; ARNT) and activates transcription by binding to hypoxic response elements (HREs) scattered throughout the genome (Gu *et al.*, 2000). HIF $\alpha$  is labile under normoxic conditions, but stabilized by hypoxia, enabling dimerization with ARNT to induce genes such as angiogenic factors, glycolytic enzymes, glucose transporters, and erythropoietin (Wang and Semenza, 1993; Firth *et al.*, 1994; Liu *et al.*, 1995; Forsythe *et al.*, 1996; Semenza *et al.*, 1996; Ema *et al.*, 1997; Okino *et al.*, 1998). *Arnt*<sup>-/-</sup> mice die by embryonic day (E) 10.5 with defects in angiogenesis, cardiogenesis, hematopoiesis, and placentation (Kozak *et al.*, 1997; Maltepe *et al.*, 1997; Adelman *et al.*, 1999, 2000; Abbott and Buckalew, 2000). *Hif1 $\alpha$* <sup>-/-</sup> embryos also die by E10.5, exhibiting angiogenic and cardiac defects (Iyer *et al.*, 1998; Ryan *et al.*, 1998; Compennolle *et al.*, 2003). Mutations in *Hif2 $\alpha$*  also result in embryonic lethality, with *Hif2 $\alpha$* <sup>-/-</sup> animals exhibiting defects in cardiac function, angiogenesis, and lung maturation (Tian *et al.*, 1998; Peng *et al.*, 2000; Compennolle *et al.*, 2002). Therefore, HIF activity is essential for viability and cardiovascular development during embryogenesis.

*Arnt*<sup>-/-</sup> placentas are avascular, hypocellular, and exhibit aberrant cell fate adoption in addition to poor invasion. To study mechanisms underlying HIF regulation of multiple aspects of placentation, we used undifferentiated trophoblast stem (TS) cells from wild-type, *Arnt*<sup>-/-</sup>, *Hif1 $\alpha$* <sup>+/-</sup>–*Hif2 $\alpha$* <sup>+/-</sup>, and *Hif1 $\alpha$* <sup>-/-</sup>–*Hif2 $\alpha$* <sup>-/-</sup> embryos. TS cells are non-transformed, immortal precursor stem cells derived from blastocysts capable of differentiating into multiple placental

This article was published online ahead of print in *MBC in Press* (<http://www.molbiolcell.org/cgi/doi/10.1091/mbc.E04-12-1082>) on February 2, 2005.

Address correspondence to: M. Celeste Simon (celeste2@mail.med.upenn.edu).

lineages (e.g., trophoblast giant cells, spongiotrophoblasts, and syncytiotrophoblasts).

Because HIF-deficient placentas exhibit poor maternal invasion, we determined whether HIF regulates trophoblast invasion through integrin-mediated adhesion to the extracellular matrix (ECM). We show here that undifferentiated *Arnt*<sup>-/-</sup> and *Hif1* $\alpha$ <sup>-/-</sup>*Hif2* $\alpha$ <sup>-/-</sup> TS cells exhibit reduced  $\alpha$ v $\beta$ 3 integrin-mediated adhesion and migration due to decreased cell surface expression of this integrin. These studies explain in part the poorly invasive phenotype observed in *Arnt*<sup>-/-</sup> and *Hif1* $\alpha$ <sup>-/-</sup>*Hif2* $\alpha$ <sup>-/-</sup> placentas in vivo. Moreover, because HIF regulates  $\alpha$ v $\beta$ 3 integrin-mediated adhesion, we reasoned that hypoxia also may up-regulate  $\alpha$ v $\beta$ 3 integrin expression in tumors, leading to increased metastasis. Melanoma metastases contain hypoxic regions, and metastatic progression is associated with increased  $\alpha$ v $\beta$ 3 integrin expression (Gehlsen *et al.*, 1992; Nip *et al.*, 1992; Lartigau *et al.*, 1997; Felding-Habermann *et al.*, 2002). Poorly and highly metastatic melanoma cells (B16F0 and B16F10, respectively) were used to study hypoxic regulation of adhesion pathways that may be relevant to tumor metastasis. We show that hypoxia induces  $\alpha$ v $\beta$ 3 integrin cell surface expression in poorly metastatic B16F0 melanoma cells, resulting in increased adhesion and migration.

## MATERIALS AND METHODS

### Antibodies

The Novus HIF1 $\alpha$  antibody (clone H1 $\alpha$ 67) was used for HIF1 $\alpha$  electrophoretic mobility shift assay (EMSA) "supershift." PE-RMV-7 from BD Biosciences PharMingen (San Diego, CA) was used for fluorescence-activated cell sorting (FACS) of  $\alpha$ v integrin, and the Chemicon International (Temecula, CA) polyclonal  $\alpha$ v integrin antibody was used for immunoprecipitation and Western blot assays. For  $\beta$ 3 integrin, 2C9.G2 subclone from BD Biosciences PharMingen was used for blocking function, FACS, immunoprecipitation, immunoelectron microscopy (immuno-EM), and immunofluorescence. For  $\beta$ 1 integrin, MAB1997 was used for immunoprecipitation. Additionally, the Chemicon International  $\beta$ 3 integrin and  $\beta$ 5 integrin antibodies were used in Western blot assays. For FACS analysis, BD Biosciences PharMingen's Ha2/5 was used for  $\beta$ 1 integrin, the Santa Cruz Biotechnology (Santa Cruz, CA) (H-96) for  $\beta$ 5 integrin, or BD Biosciences PharMingen's GoH3 for  $\alpha$ 6 integrin. For immunofluorescence, 58K (Abcam, Cambridge, MA), anti-calreticulin (StressGen Biotechnologies, San Diego, CA), phospho-Y397FAK (BD Transduction Laboratories, Lexington, KY), and anti-mouse-Alexa-fluor 488, anti-rabbit-Alexa-fluor 488, and streptavidin Alexa-fluor 568 from Molecular Probes (Eugene, OR) were used.

### EMSAs

EMSAs were performed in binding buffer consisting of 10 mM Tris-HCl, pH 7.5, 50 mM NaCl, 50 mM KCl, 1 mM MgCl<sub>2</sub>, 5 mM dithiothreitol, 1 mM EDTA, and 5% glycerol to which 0.1 mg/ml bovine serum albumin (BSA), 5  $\mu$ g of nuclear extract, and 10<sup>5</sup> cpm of probe were added per 20  $\mu$ l of reaction. The binding site sequence of the probe (E24F) is 5'-GCCCTACGTGCTGCCTCGCATGGC-3'. Cold cAMP response element-binding protein (CREB) competitor oligo (100 $\times$  excess) containing the CREB binding site from the somatostatin promoter was added: 5'-GATCGCCTCCTGGCTGACGTCA-GAGAGCTAG-3'.

### Northern Blot Analysis

Undifferentiated TS and embryonic stem (ES) cells were cultured for 16 h at 20% or 1.5% O<sub>2</sub> and then washed with Dulbecco's phosphate-buffered saline (DPBS). RNA was extracted with TRIzol reagent (Invitrogen, Carlsbad, CA) according to manufacturer's protocol. Fifteen micrograms of total RNA was electrophoresed in 1% agarose containing formaldehyde. The RNAs were transferred to Hybond N<sup>+</sup> membranes (Amersham Biosciences, Piscataway, NJ). Full-length HIF1 $\alpha$  and  $\beta$ -tubulin cDNAs were labeled as probes.

### Cell Culture

*Arnt*<sup>+/+</sup> and *Arnt*<sup>-/-</sup> TS were described previously (Adelman *et al.*, 2000) and derived by standard protocols (Tanaka *et al.*, 1998). *Hif1* $\alpha$ <sup>+/+</sup>*Hif2* $\alpha$ <sup>+/+</sup>, and *Hif1* $\alpha$ <sup>-/-</sup>*Hif2* $\alpha$ <sup>-/-</sup> TS cells are described by Cowden Dahl, Mack, Comperolle, Adelman, Carmeliet, and Simon (unpublished data). Briefly, TS cells were maintained in an undifferentiated state on mouse embryo fibroblasts (MEFs) in the presence of FGF4 and heparin. For all experiments, TS cells

were cultured in the absence of MEFs but supplemented with MEF-conditioned media containing FGF4 and heparin. TS cells were plated on glass coverslips for immunofluorescence assays. B16F0 and B16F10 cells were obtained from the American Type Culture Collection (Manassas, VA) and grown in DMEM supplemented with 10% serum, HEPES, penicillin, streptomycin, and L-glutamine. B16 melanoma cells were cultured either under standard culture conditions (20% O<sub>2</sub>) or at 1.5% O<sub>2</sub> (in an IG750 3 gas incubator; Jouan, Winchester, VA) for 24 h. CO<sub>2</sub> is maintained at 5% for all experiments.

### Immunoprecipitation of Biotin-labeled Cell Surface Proteins

TS or melanoma cells were cultured to 75% confluence, placed on ice, and washed with ice cold DPBS with calcium and magnesium. Cells were then incubated in 1 mg/ml NHS-LC-Biotin (Pierce Chemical, Rockford, IL) for 90 min on ice and washed three times with 50 mM glycine in DPBS. Cell lysates were prepared in NP-40 buffer, pH 7.5 (0.1 M Tris-Base, 1% NP-40, 10  $\mu$ g/ml Pefabloc, 1 mM E-64, 5  $\mu$ g/ml leupeptin, 1  $\mu$ g/ml pepstatin, 0.02 mg/ml aproptinin, and 0.5M EDTA). Debris was pelleted, and supernates were transferred to new tubes. Lysates were precleared overnight with protein A/G beads (Pierce Chemical). Biotinylated acid assays were performed on precleared lysates to determine protein concentration. Immunoprecipitations were performed on 250  $\mu$ g of protein by using protein A/G beads and anti- $\alpha$ v integrin (Chemicon International), anti- $\beta$ 3 integrin (BD Biosciences PharMingen) or anti- $\beta$ 1 integrin (Chemicon International). After immunoprecipitation, beads were washed thrice with MDB buffer (50 mM Tris, pH 7.5, 0.1% SDS, 0.05% NP-40, and 0.3 M NaCl) and thrice with Tris-wash buffer (10 mM Tris, pH 7.5). Lysates were then resuspended in nonreducing sample buffer and run on an 8% SDS-polyacrylamide gel.

### FACS Analysis

Cell surface expression of integrins was determined by FACS analysis. Cells were trypsinized and washed in DPBS and blocked with anti-CD16/32 for 15 min at room temperature. After washing in 1% BSA in DPBS, cells were incubated with primary antibody for 30 min on ice, washed with 1% BSA/DPBS, pelleted, and resuspended in isotonic buffer. Propidium iodide was used to exclude dead cells. FACS data were analyzed using CellQuest software, and error bars represent  $\pm$ SE. Student's *t* tests were performed to determine statistical significance.

### Adhesion, Migration, and Invasion Assays

Adhesion assays were performed on CytoMatrix cell adhesion strips containing fibronectin (FN), vitronectin (VN), collagen IV, and laminin (LN) according to manufacturer's instructions (Chemicon International). The concentration of ECM proteins and BSA for the adhesion and migration assays is not provided by Chemicon International. Briefly, strips were rehydrated with DPBS. Cells were prepared in a single cell suspension in serum-free media, and 1  $\times$  10<sup>5</sup> cells were added to each well of the adhesion strips. Strips were incubated for 1 h at 37°C and washed twice with DPBS, stained with 0.2% crystal violet in 10% ethanol (EtOH), washed again, and solubilized in 50% 0.1 M NaH<sub>2</sub>PO<sub>4</sub> in 50% EtOH. Absorbance was read at 560 nm on a microplate reader. Quantitative cell migration assays (QCM-FN and QCM-VN; Chemicon International) were performed on feeder-free TS cells according to manufacturer's instructions. Briefly, 2.5  $\times$  10<sup>5</sup> cells were placed in each well of FN-, VN-, or BSA-coated Boyden chambers. Cells were allowed to migrate for 16 h, and chambers were stained with crystal violet. Dye was then solubilized and read on a microplate reader. Absorbance was normalized to migration toward BSA. Invasion was measured on BD BioCoat Matrigel (BD Biosciences, San Jose, CA) invasion chambers according to manufacturer's instructions. Briefly, inserts were rehydrated with RPMI 1640 medium. Cells (4  $\times$  10<sup>5</sup>) were added to chambers and incubated for 22 h. Chambers were then cleaned and stained with crystal violet. Solubilized dye was measured on a microplate reader.

### Western Blots

Cells were washed in DPBS and lysed in NP-40 buffer (described above). Lysates were cleared by centrifugation, and equal amounts of protein were separated on a 10% SDS-polyacrylamide gel. Proteins were transferred to nitrocellulose, and the membrane was blocked with 3% BSA in Tris-buffered saline/Tween 20 and incubated with antibodies before development with ECL.

### Immunoelectron Microscopy

Briefly, samples were rinsed in buffer and fixed in 2% paraformaldehyde + 0.2% glutaraldehyde in DPBS for 30 min and permeabilized with 0.1% saponin. Fixative was removed, cells washed, and residual aldehyde was inactivated by treatment with 0.1% NaBH for 10 min. Cells were incubated with 1% ovalbumin + 5% normal horse or donkey serum and 0.2% cold water fish skin gelatin for 60 min at room temperature to prevent nonspecific binding. Primary antibody incubation was performed overnight at a dilution of 1:100

cells washed and treated with ultrasmall gold secondary overnight. After washing with DPBS cells were fixed in 1% glutaraldehyde in DPBS for 15 min. Cells were washed with DPBS and then with deionized water before enhancing with silver stain. After silver staining cells were osmicated with 1% osmium and dehydrated in graded alcohol and infiltration aided by propylene oxide and embedded in Epon. Ultrathin sections were mounted on Formvar-coated slotted copper grids and imaged in JEOL JEM 1010 aided by AMT HR-12 software and Hamamatsu Charge-coupled device camera.

### Fluorescence Confocal Microscopy

TS cells were plated on uncoated coverslips and then fixed for 10 min at room temperature in  $-20^{\circ}\text{C}$  acetone, washed in DPBS, permeabilized with DPBS/0.2% Triton X-100, and blocked in 0.2% BSA in DPBS-Tween 20. Cells were then incubated with primary and conjugated-secondary antibodies at room temperature for 1 h each. Coverslips were mounted on slides. Samples were viewed by confocal microscopy. Images were acquired for both red and green lasers and merged in Photoshop.


### Statistical Analysis

Each experiment was performed at least three times. Error bars represent  $\pm$  SE. Student's *t* tests were performed to determine statistical significance for the following assays: adhesion, migration, invasion, FACS, and immuno-EM quantitation.

## RESULTS

### TS Cells Exhibit High Levels of HIF Activity When Cultured in 20% $\text{O}_2$

*Arnt*<sup>-/-</sup> TS cells exhibit a number of phenotypes reminiscent of defects observed in *Arnt*<sup>-/-</sup> placentas, including aberrant cell fate adoption upon differentiation and poor adhesion to ECM (see below). Because differentiated *Arnt*<sup>-/-</sup> TS cells exhibit defects in cell fate adoption, we chose to confine the experiments exclusively to *undifferentiated* TS cells. We determined that the degree of differentiation in both wild-type and *Arnt*<sup>-/-</sup> TS cells was very low based on trophoblast lineage-specific gene expression, morphology, and DNA content (our unpublished data). Interestingly, defects in *Arnt*<sup>-/-</sup> TS cells are detected under both normoxic and hypoxic conditions. To demonstrate that the *Arnt*<sup>+/+</sup> and *Arnt*<sup>-/-</sup> TS cell phenotypes observed under normoxic conditions are mediated by the HIF pathway, we assessed the degree of HIF activity in normoxic and hypoxic TS cells. First, we examined mRNA expression of HIF1 $\alpha$  and HIF2 $\alpha$  in TS cells. For most experiments, we used two independent *Arnt*<sup>+/+</sup> (0 and 2) and *Arnt*<sup>-/-</sup> (4 and 10) TS cell lines. In Figure 1A, both *Arnt*<sup>+/+</sup> (0 and 2) and *Arnt*<sup>-/-</sup> (4 and 10) TS cell lines were examined for gene expression at 20% and 1.5%  $\text{O}_2$ . Because tubulin expression is regulated by  $\text{O}_2$  tension, the 28S RNA band serves as an important second loading control. Even though the abundance of HIF1 $\alpha$  mRNA somewhat varied between TS cell lines, HIF1 $\alpha$  mRNA was present at two- to threefold higher levels under both normoxia (20%  $\text{O}_2$ ) and hypoxia (1.5%  $\text{O}_2$ ) in all TS cell lines compared with wild-type ES cells (Figure 1A). Although HIF2 $\alpha$  is present in ES cells (our unpublished data), HIF2 $\alpha$  RNA was not detected in undifferentiated TS cells, indicating that all TS HIF activity is therefore due to HIF1 $\alpha$ /ARNT dimers. We then performed an EMSA to detect HIF DNA binding to an HRE residing in the erythropoietin (EPO) 3' enhancer. Low levels of HIF binding activity were observed in normoxic ES cells, which increased upon exposure to 1.5%  $\text{O}_2$  (Figure 1B, lanes 1 and 2). Importantly, antibody against HIF1 $\alpha$  completely shifted the hypoxic DNA binding complex, indicating that the inducible complex contains HIF (Figure 1B, lane 3). In comparison with ES cells, HIF DNA binding activity was detected in TS cells at relatively high levels at 20%  $\text{O}_2$ ; DNA binding activity was further enhanced by hypoxia and supershifted with  $\alpha$ -HIF1 $\alpha$  antibody (Figure 1B, lanes 4–6). These data suggested that HIF target




**Figure 1.** TS cells exhibit high levels of HIF activity when cultured in 20%  $\text{O}_2$ . (A) Northern blot analysis for HIF1 $\alpha$  mRNA levels in two independent *Arnt*<sup>+/+</sup> TS cell lines, two independent *Arnt*<sup>-/-</sup> TS cell lines, and wild-type ES cells. Tubulin and 28S RNA serve as loading controls. As expected, HIF1 $\alpha$  mRNA levels are not increased by culture at 1.5%  $\text{O}_2$  in either TS or ES cells. Each of these experiments was performed three times. (B) EMSA analysis showing HIF DNA complex formation. *Arnt*<sup>+/+</sup> ES cells demonstrate low levels of the HIF complex under normoxic conditions, but complex formation is enhanced under hypoxia. In addition to the hypoxia-inducible DNA complex, ES cells harbor a constitutive nonspecific complex as indicated. *Arnt*<sup>+/+</sup> TS cells exhibit HIF complex formation under normoxia and hypoxia, but no complex forms in the absence of ARNT. HIF complexes are supershifted with an anti-HIF1 $\alpha$  antibody, as indicated by the asterisk.

genes might be regulated by HIF in TS cells even under normoxic conditions. As expected, no HIF DNA binding activity was present in *Arnt*<sup>-/-</sup> TS cells (Figure 1B, lanes 7–9). Furthermore, we observed decreased expression of the HIF target genes vascular endothelial growth factor (VEGF), ERO1-L $\alpha$ , phosphofructose kinase (PFK), and Glut-1 (our unpublished data) in normoxic TS cells. Therefore, we concluded that HIF is partially active under normoxic conditions in *Arnt*<sup>+/+</sup> TS cells and conducted all subsequent experiments at 20%  $\text{O}_2$  to study a homogenous stem cell population.

### *Arnt*<sup>-/-</sup> TS Cells Exhibit Poor Migration and Adhesion


Because *Arnt*<sup>-/-</sup> trophoblasts exhibit poor invasion into maternal tissue, we investigated whether HIF activity plays a



**Figure 2.** *Arnt*<sup>-/-</sup> TS cells exhibit impaired migration toward vitronectin and fibronectin and reduced invasion into Matrigel. (A) Boyden chamber assays for migration toward vitronectin. (B) Boyden chamber assays for migration toward fibronectin. (C) Invasion of Matrigel. (D) Photographs of migration toward vitronectin (top) and migration toward fibronectin (bottom). Original magnifications, 100 $\times$ . Arrows indicate stained cells that have migrated through Boyden chamber pores. Each experiment was performed three times in duplicate. Error bars represent  $\pm$  SEM. Student's *t* tests were used for statistical significance.

role in undifferentiated TS cell migration. Migration toward the ECM proteins vitronectin and fibronectin was analyzed using a Boyden chamber assay, and migration was normalized to the wild-type control +/+ (2) TS cells for each experiment. Migration toward vitronectin was decreased by 5- and 5.9-fold, respectively, in two independent *Arnt*<sup>-/-</sup> TS cell lines compared with wild-type cells ( $p < 0.01$  and  $p < 0.01$ ) (Figure 2A). Additionally, migration toward fibronectin was decreased by 16.7- and 6.8-fold in *Arnt*<sup>-/-</sup> TS cell lines (Figure 2B). Photographs of the Boyden chambers showing decreased migration toward vitronectin (top) and fibronectin (bottom) are presented in Figure 2D. We also measured the ability of TS cells to invade basement membrane (Matrigel), which contains collagens, laminin, proteoglycans, matrix degrading enzymes and their inhibitors, and growth factors. Both *Arnt*<sup>+/+</sup> cell lines exhibited roughly equivalent invasion. In contrast, *Arnt*<sup>-/-</sup> TS cell lines displayed decreased invasion (3.2- and 2.1-fold, respectively) into Matrigel compared with the control +/+ (2) cells ( $p < 0.01$ ) (Figure 2C). We concluded that trophoblasts lacking HIF activity exhibit poor migration and invasion.


ECM is composed of a number of proteins that serve as ligands for integrins, including fibrous proteins (e.g., collagen), linker proteins (e.g., fibronectin, laminin, and vitronectin), and glycosaminoglycans. We reasoned that the decreased migration exhibited by *Arnt*<sup>-/-</sup> TS cells might be due to a defect in ECM adhesion. TS cells were plated on vitronectin-, fibronectin-, collagen IV-, or laminin-coated plates for 1 h to measure binding to different substrates. Binding to vitronectin was decreased by approximately five-fold in both *Arnt*<sup>-/-</sup> TS cell lines ( $p < 0.01$ ) (Figure 3A), whereas adhesion to fibronectin was decreased by  $\sim$ 1.8-fold ( $p < 0.001$ ) (Figure 3B). Because integrin  $\alpha$ v $\beta$ 3 is the prototypic receptor for vitronectin and has affinity for fibronectin, we determined whether the ECM adhesion defects were due to a decrease in  $\alpha$ v $\beta$ 3 activity. We treated *Arnt*<sup>+/+</sup> and *Arnt*<sup>-/-</sup> TS cells with a  $\beta$ 3 integrin-blocking antibody and then performed adhesion assays. Adhesion to vitronectin was greatly decreased in *Arnt*<sup>+/+</sup> and *Arnt*<sup>-/-</sup> TS cells by blocking  $\beta$ 3 integrin activity ( $p < 0.01$ ); however, adhesion to fibronectin was not (Figure 3, C and D). This suggested that inhibition of  $\beta$ 3 activity is sufficient to decrease adhe-



**Figure 3.** *Arnt*<sup>-/-</sup> TS cells adhere poorly to both vitronectin and fibronectin. *Arnt*<sup>+/+</sup> and *Arnt*<sup>-/-</sup> TS cells were plated on strips coated with vitronectin (A) or fibronectin (B) and stained with crystal violet. Solubilized crystal violet incorporated by adherent cells was measured on a plate reader at 560 nm. Two independent wild-type and mutant lines were used. Treatment of TS cells with a blocking antibody for  $\beta 3$  integrin reduces vitronectin (C) but not fibronectin (D). *Arnt*<sup>-/-</sup> TS cells adhere to collagen IV (E) and laminin (F) as well as *Arnt*<sup>+/+</sup> TS line 0. *Hif1 $\alpha$* <sup>-/-</sup> TS cells demonstrate a substantial reduction in adhesion to vitronectin ( $p < 0.01$ ) and are somewhat reduced in adhesion to fibronectin coated-strips compared with *Hif1 $\alpha$* <sup>+/-</sup> TS cells (G). Each experiment was performed three times in triplicate. Error bars represent  $\pm$  SEM. Student's *t* tests were used for statistical significance.

sion to vitronectin, but blocking  $\beta 3$  activity alone does not inhibit adhesion to fibronectin. Adhesion to collagen IV and laminin was equivalent between *Arnt*<sup>-/-</sup> cell lines and one of the wild-type cell lines +/+ (0) (Figure 3, E and F). The second wild-type cell line [+/+(2)] exhibited more adhesion to laminin and collagen IV than the other lines, indicating that this effect was not genotype specific. To demonstrate that the decrease in adhesion to vitronectin and fibronectin

was due to inactivation of the HIF pathway, ECM adhesion for TS cells lacking both *Hif1 $\alpha$*  and *Hif2 $\alpha$*  (designated *Hif1 $\alpha$* <sup>-/-</sup>) was examined. *Hif1 $\alpha$* <sup>-/-</sup> TS cells exhibited a fivefold decrease in adhesion to vitronectin compared with *Hif1 $\alpha$* <sup>+/-</sup> *Hif2 $\alpha$* <sup>+/-</sup> (*Hif1 $\alpha$* <sup>+/-</sup>) TS cells ( $p < 0.01$ ) (Figure 3G). Adhesion to fibronectin was also slightly decreased in *Hif1 $\alpha$* <sup>-/-</sup> TS cells (1.5-fold), but this change was not significant. Importantly, adhesion of *Hif1 $\alpha$* <sup>+/-</sup> TS cells to laminin was equivalent to



**Figure 4.** TS cells lacking *Arnt* exhibit reduced cell surface  $\alpha v\beta 3$  integrin. (A) Immunoblots showed that integrins  $\alpha v$ ,  $\beta 3$ , and  $\beta 5$  are expressed equivalently in wild-type and mutant TS whole cell lysates. Cyclin E is used as a loading control. (B) TS cells were surface labeled with biotin, and lysates were immunoprecipitated with antibodies for  $\alpha v$  integrin or  $\beta 3$  integrin and separated by nonreducing SDS-PAGE, which was subsequently blotted with streptavidin-HRP. Positions of the individual integrins are indicated. Lanes 1, 2, 5, 6, 9, and 10 are *Arnt*<sup>+/+</sup> cell lysates; lanes 3, 4, 7, 8, 11, and 12 are *Arnt*<sup>-/-</sup> cell lysates; lane 13 is *Hif* $\alpha$ <sup>+/-</sup> cell lysate, whereas lane 14 is *Hif* $\alpha$ <sup>-/-</sup> cell lysate. (C) *Arnt*<sup>+/+</sup> and *Arnt*<sup>-/-</sup> cells were analyzed by FACS for surface expression of integrins  $\alpha v$ ,  $\beta 1$ ,  $\beta 3$ , and  $\beta 5$ . Representative histograms are shown as well as bar graphs for the mean fluorescence intensity of at least three independent experiments for each integrin. Error bars represent  $\pm$  SEM. Student's *t* tests were performed to determine statistical significance. Of note, only one *Arnt*<sup>-/-</sup> TS line exhibited reduced surface integrin  $\alpha v$  expression.

*Hif* $\alpha$ <sup>-/-</sup> TS cells (our unpublished data). We concluded that *Arnt*<sup>-/-</sup> and *Hif* $\alpha$ <sup>-/-</sup> TS cells exhibit striking defects in adhesion to vitronectin and fibronectin.

**In the Absence of ARNT,  $\alpha v\beta 3$  Integrin Does Not Localize to the Cell Surface**

Because the adhesion and migration defects were specific for the ECM proteins vitronectin and fibronectin, we evaluated a role for HIF in the regulation of adhesion receptors. As stated above, integrin  $\alpha v\beta 3$  is a receptor for both vitronectin and fibronectin. Furthermore,  $\alpha v\beta 3$  integrin activity has been shown to be important in cytotrophoblast invasion in

the human placenta (Zhou *et al.*, 1997a). First, we determined whether HIF regulates expression of  $\alpha v\beta 3$  integrin. We performed Western blots on TS cell extracts to examine protein levels for  $\alpha v$ ,  $\beta 3$ , and  $\beta 5$  integrins. We found comparable levels of each of these integrins in whole cell lysates prepared from *Arnt*<sup>+/+</sup> and *Arnt*<sup>-/-</sup> TS cells (Figure 4A). Cyclin E was used as a loading control. These data suggest that the  $\alpha v$  and  $\beta 3$  integrins are not regulated at the level of transcription or translation and therefore are not likely to be direct HIF target genes in TS cells.

Next, we examined the cell surface expression of  $\alpha v\beta 3$  integrin on *Arnt*<sup>+/+</sup>, *Arnt*<sup>-/-</sup>, *Hif* $\alpha$ <sup>+/-</sup>, and *Hif* $\alpha$ <sup>-/-</sup> TS cells.



TS cell surfaces were biotinylated, and whole cell lysates were immunoprecipitated with antibodies to either  $\alpha v$  integrin or  $\beta 3$  integrin. Immunoprecipitations (IPs) were fractionated by nonreducing SDS-PAGE, blotted, and probed with streptavidin-horseradish peroxidase (HRP). The top left panel of Figure 4B presents an experiment where IPs were washed under high salt conditions (0.6 M NaCl), causing  $\beta 1$  integrin to dissociate from  $\alpha v$  integrin. The bottom panel represents IPs washed less stringently (0.3 M NaCl), and the position of  $\beta 1$  integrin is indicated.  $Arnt^{+/+}$  TS cells lysates immunoprecipitated with antibody to  $\alpha v$  integrin exhibited expression of  $\alpha v\beta 1$ ,  $\alpha v\beta 5$ , and  $\alpha v\beta 3$  integrin dimers on their cell surface (Figure 4B, lanes 1, 2, 9, and 10). Additionally, when  $Arnt^{+/+}$  extracts were immunoprecipitated with  $\beta 3$  integrin antibody, substantial  $\alpha v\beta 3$  integrin was detected on the cell surface (Figure 4B, lanes 5 and 6). In contrast,  $Arnt^{-/-}$  TS cells showed dramatically decreased  $\alpha v\beta 3$  integrin on the cell surface when lysates were immunoprecipitated with antibody against  $\alpha v$  integrin (lanes 3, 4, 11, and 12) or  $\beta 3$  integrin (lanes 7 and 8). The top right panel of Figure 4B depicts  $Hif\alpha^{+/+}$  and  $Hif\alpha^{-/-}$  TS cells that have been biotinylated and immunoprecipitated with integrin  $\alpha v$ .  $Hif\alpha^{-/-}$  TS cells exhibit reduced surface  $\alpha v\beta 3$  integrin compared with  $Hif\alpha^{+/+}$  TS cells (Figure 4B, lanes 13 and 14). Importantly,  $Arnt^{-/-}$  TS cells expressed comparable levels of  $\alpha v\beta 1$  integrin and  $\alpha v\beta 5$  integrin on their cell surface compared with  $Arnt^{+/+}$  TS cells. We also used FACS analysis to determine the extent of cell surface expression for integrins  $\alpha v$ ,  $\beta 1$ ,  $\beta 3$ , and  $\beta 5$ . Only one of the  $Arnt^{-/-}$  TS cell lines expressed less  $\alpha v$  integrin on the cell surface than wild-type cells, so this is not a genotype-specific phenomenon. Instead, we found that  $\beta 3$  integrin cell surface expression was decreased in both  $Arnt^{-/-}$  TS cell lines (Figure 4C) whereas integrins  $\beta 1$  and  $\beta 5$  were expressed at wild-type levels. In conclusion, HIF regulation of integrin  $\alpha v\beta 3$  cell surface expression seems to mediate adhesion to vitronectin and fibronectin.


Because  $\beta 3$  integrin seemed to be translated normally in  $Arnt^{-/-}$  TS cells, we examined cellular localization of  $\beta 3$  integrin by performing immunofluorescence confocal microscopy analysis. TS cells grow as epithelial sheets, so the micrographs shown consist of undifferentiated cells within a sheet and not individual cells. We first determined whether  $\beta 3$  integrin localized to the Golgi apparatus by analyzing colocalization between  $\beta 3$  integrin and 58K, a microtubule-binding Golgi protein (Bashour and Bloom, 1998).  $Arnt^{+/+}$  and  $Arnt^{-/-}$  TS cells were initially stained for  $\beta 3$  integrin (in red) and 58K (in green) by using fluorescently conjugated antibodies (Figure 5A). The yellow color present in the merged image of both  $Arnt^{+/+}$  and  $Arnt^{-/-}$  TS cells indicates colocalization between  $\beta 3$  integrin and 58K, suggesting that  $\beta 3$  integrin is present in the Golgi (Figure 5A). We subsequently assessed colocalization between  $\beta 3$  integrin (red) and the ER protein calreticulin (green).  $\beta 3$  integrin was detected in the ER in both  $Arnt^{+/+}$  and  $Arnt^{-/-}$  TS cells (Figure 5B). When integrins are activated, they cluster at focal adhesions to mediate attachment to the ECM (Giancotti and Ruoslahti, 1999). Focal adhesions are composed of structural proteins, cytoskeletal proteins, and signaling molecules. On integrin clustering, focal adhesion kinase (FAK) becomes phosphorylated at tyrosine 397 (Rodriguez-Fernandez, 1999). To determine whether  $\beta 3$  was present in TS cell focal adhesions, colocalization between  $\beta 3$  integrin and phosphorylated FAK was examined. Although integrin  $\beta 3$  was localized with phosphorylated FAK in wild-type cells, very little  $\beta 3$  integrin was associated with focal adhesions in  $Arnt^{-/-}$  TS cells (Figure 5C). In agreement with the de-

creased  $\beta 3$  integrin observed by FACS and immunoprecipitation assays,  $\beta 3$  integrin transits through the Golgi and ER in  $Arnt^{-/-}$  TS cells but is not transported to focal adhesions at the cell surface.

To further analyze  $\beta 3$  subcellular localization, we performed immuno-EM for  $\beta 3$  integrin. Immuno-EM is an unbiased approach used to examine the association of a particular protein with ultrastructural features or subcellular compartments. In  $Arnt^{+/+}$  TS cells,  $\beta 3$  integrin-specific gold particles were localized near the cell surface and in small intracellular clusters (Figure 5E). In direct contrast,  $\beta 3$  integrin was never present on the cell surface in  $Arnt^{-/-}$  cells. Instead, there were large clusters of intracellular  $\beta 3$ -specific gold particles, indicating that  $\beta 3$  integrin was localized in the cell interior (Figure 5F). These gold particle aggregates seemed to be in membrane-bound compartments, which may be some type of vesicle. To quantitate these findings, we counted small gold particle clusters (1 gold particle) or large gold particle clusters ( $\geq 3$  gold particles) for at least nine electron micrographs for  $Arnt^{+/+}$  and  $Arnt^{-/-}$  TS cells.  $Arnt^{-/-}$  TS cells had significantly more large clusters of gold particles ( $p = 0.048$ ) and fewer small clusters of gold particles (Figure 5D) than wild-type cells. Therefore,  $\beta 3$  integrin seems to be accumulating in a post-Golgi, membrane-bound compartment in  $Arnt^{-/-}$  TS cells. Accumulation in this compartment either prevents transport of  $\beta 3$  integrin to the cell surface or  $\beta 3$  integrin accrues in this location because it is unable to be transported properly to the surface. These results provide further evidence that  $\beta 3$  integrin is intracellular but does not reach the cell surface in TS cells deficient for HIF.

#### *Hypoxia Regulates $\alpha v\beta 3$ Integrin-mediated Adhesion and Migration in Melanoma Cells*


Integrin expression and activation is important in both developmental processes and pathological conditions. It has been reported that hypoxia is associated with radiation resistance in tumors and predicts poor patient outcome (Vaupel *et al.*, 2001). Additionally,  $\alpha v\beta 3$  integrin expression correlates with metastasis or an increased invasive phenotype in a number of tumor cell types such as melanoma, ovarian carcinoma, hepatocellular carcinoma, and breast cancer (Carreiras *et al.*, 1996; Wong *et al.*, 1998; Nejari *et al.*, 2002; Jin and Varner, 2004). We examined the possible regulation of  $\alpha v\beta 3$  integrin by hypoxia in melanoma cells. We assayed both highly metastatic B16F10 murine melanoma cells and poorly metastatic B16F0 cells for cell surface expression of  $\alpha v\beta 3$  integrin. Based on FACS analysis, B16F10 cells exhibited 50% more  $\alpha v$  integrin ( $p < 0.001$ ) and 25% more  $\beta 3$  integrin ( $p = 0.01$ ) on the cell surface than B16F0 cells (Figure 6, A and B). Unlike the TS cells, culturing cells at 1.5%  $O_2$  for 24 h induced  $\alpha v\beta 3$  integrin protein expression in B16F0 and B16F10 cells approximately twofold (our unpublished data). Of note, the highly metastatic B16F10 cells were more adherent to vitronectin ( $p < 0.01$ ) and laminin ( $p = 0.025$ ) than the poorly metastatic B16F0 cells (Figure 6C). To further investigate the role of hypoxia in tumor progression, we examined changes in adhesion properties of melanoma cells in response to hypoxic culture conditions. We subjected B16F0 cells to 24 h of 1.5%  $O_2$  and measured adhesion to fibronectin, vitronectin, and laminin. Hypoxia increased B16F0 adhesion to fibronectin and vitronectin by 1.6-fold ( $p = 0.017$ ) and 1.7-fold ( $p = 0.001$ ), respectively, compared with culture in 20%  $O_2$  (Figure 7A). In contrast, adhesion to laminin was unaltered by hypoxic exposure (Figure 7A).



**Figure 5.** Subcellular localization of integrin  $\beta 3$  in TS cells. Confocal immunofluorescence microscopy was performed on  $Arnt^{+/+}$  and  $Arnt^{-/-}$  TS cells showing colocalization of  $\beta 3$  integrin with the Golgi marker 58K (A), calreticulin in the ER (B), and phosphorylated FAK Y397 (C). Integrin  $\beta 3$  is in red and the other markers are in green. The yellow image is a red/green merge to show colocalization. Intracellular small and large gold particle clusters for  $\beta 3$  integrin in wild-type and  $Arnt^{-/-}$  TS cells from immunoelectron micrographs was quantitated (D). Immunoelectron micrographs for  $\beta 3$  integrin in  $Arnt^{+/+}$  (E) and  $Arnt^{-/-}$  TS cells (F). Bar, 500 nm. Error bars represent  $\pm$  SEM. Student's *t* tests were performed to determine statistical significance.

We also measured B16F0 cell migration toward vitronectin. Treatment of B16F0 cells with 1.5%  $O_2$  resulted in increased vitronectin-associated migration ( $p = 0.045$ ) (Figure 7B). Assays for invasion of B16F0 and B16F10 cells into Matrigel were performed; unfortunately, both cell lines failed to perform in these experiments. When a blocking antibody for  $\beta 3$  integrin was added to B16F0 cells, adhesion to vitronectin under both normoxia and hypoxia was decreased ( $p < 0.01$ ) (Figure 7C), suggesting that adhesion to vitronectin is mediated by  $\beta 3$  integrin. In contrast to B16F0 cells, exposure of B16F10 cells to

1.5%  $O_2$  for 24 h did not significantly increase adhesion to vitronectin (our unpublished data). We then determined the effect of hypoxia on  $\alpha v\beta 3$  integrin cell surface expression. B16F0 cells were cultured at 20%  $O_2$  and 1.5%  $O_2$  for 24 h, and cells were harvested, stained with specific antibodies, and analyzed by FACS. B16F0 cells exhibited increased  $\alpha v$  integrin ( $p < 0.001$ ) and  $\beta 3$  integrin ( $p < 0.01$ ) cell surface expression under hypoxia (Figure 7D). However, cell surface expression of  $\alpha 6$  and  $\beta 1$  integrins did not increase after hypoxic exposure (Figure 7D).



**Figure 6.** Poorly metastatic B16F0 melanoma cells express less cell surface  $\alpha v\beta 3$  integrin than highly metastatic B16F10 cells. (A) Mean fluorescence intensity for  $\alpha v$  integrin on B16F0 and B16F10 cells based on FACS analysis. (B) Mean fluorescence intensity for  $\beta 3$  integrin on B16F0 and B16F10 cells. (C) B16F0 and B16F10 cells were plated on strips coated with FN, VN, or LN and stained with crystal violet. Solubilized crystal violet incorporated by adherent cells was measured on a plate reader at 560 nm. Error bars represent  $\pm$  SEM. Student's *t* tests were performed to determine statistical significance.

B16F0 cell surfaces also were labeled with biotin to detect membrane-bound proteins. Immunoprecipitation with antibodies to  $\alpha v$  integrin confirmed that B16F0 cells exhibited increased  $\alpha v\beta 3$  integrin cell surface expression under hypoxia, whereas immunoprecipitation with an antibody for  $\beta 1$  integrin showed no change in the abundance of integrin  $\beta 1$  on the cell surface after hypoxic treatment (Figure 7E). In conclusion, hypoxia specifically induces  $\alpha v\beta 3$  integrin expression in the poorly metastatic B16F0 melanoma cells, resulting in increased adhesion and migration. However, the mechanism of hypoxic induction is slightly different from that of TS cells because hypoxia increases  $\alpha v\beta 3$  total cellular protein in B16F0 melanoma cells.

## DISCUSSION

Data presented here demonstrate that HIF signaling is important for both adhesion of undifferentiated TS cells to vitronectin and migration properties of these cells, via control of subcellular localization of  $\alpha v\beta 3$  integrin. We also show that hypoxia promotes tumor cell migration through regulation of  $\alpha v\beta 3$  integrin-mediated adhesion. Therefore, the HIF signaling pathway and  $O_2$  availability are important in cell motility and adhesion in a number of biological contexts.

### HIF Regulates Adhesion and Migration


Placental development initially occurs in a natural  $O_2$  gradient, and  $O_2$  sensing via HIF activity in the placenta is critical for trophoblast cell fate adoption and trophoblast invasion (Adelman *et al.*, 2000; Cowden Dahl, Mack, Compennolle, Adelman, Carmeliet, and Simon, unpublished data). Trophoblast adhesion and invasion are critical for embryo implantation into the maternal tissue and subsequent placentation. Therefore, we examined the ability of HIF to regulate trophoblast adhesion and migration. Because embryos lacking ARNT or the HIF $\alpha$  subunits exhibit shallow placental invasion, we used TS cells derived from *Arnt*<sup>-/-</sup> blastocysts to assess whether the invasion defect in vivo is autonomous to trophoblasts. Normoxic undifferentiated TS cells were analyzed because they represent the most homogenous cell population. TS cells exhibit unusually high levels of HIF activity at 20%  $O_2$  conditions, thereby allowing us to study HIF-dependent processes under normal  $O_2$  tensions. TS cells deficient in the ARNT subunit of HIF mi-

grated and adhered poorly to vitronectin and fibronectin, suggesting HIF regulates these events. The specificity of the adhesion defect suggests that HIF specifically mediates adhesion through the  $\alpha v\beta 3$  integrin pair and not through other integrins.

Other groups have examined hypoxic regulation of  $\alpha v\beta 3$  integrin in endothelial cells. Hypoxia transcriptionally stimulates integrin  $\alpha v$  and  $\beta 3$  mRNA levels in human umbilical vein endothelial cells (HUVECs) (Walton *et al.*, 2000) and integrin  $\beta 2$  in human monocytes (Kong *et al.*, 2004). However, we did not observe changes in integrin  $\alpha v\beta 3$  protein levels in hypoxic TS cell or HUVECs (Figure 4C; out unpublished data). Treatment of bovine retinal endothelial cells with exogenous VEGF induces  $\alpha v$  and  $\beta 3$  integrin expression; however, VEGF treatment does not alter expression of either  $\alpha v$  integrin or  $\beta 3$  integrin in TS cells (Suzuma *et al.*, 1998; our unpublished data). In *Arnt*<sup>-/-</sup> TS cells,  $\beta 3$  integrin is transcribed, translated, and transits through the Golgi and ER, but it fails to cluster at focal adhesions on the cell surface. Therefore, the regulation of  $\alpha v\beta 3$  integrin in TS cells seems to be distinct from that in endothelial cells.

As the trophoblast of the blastocyst differentiates into trophoblasts, cell polarity, motility, and adhesion are altered (Sutherland, 2003). Integrins  $\alpha 5$ ,  $\alpha 6$ ,  $\alpha v$ ,  $\alpha 11b$ ,  $\beta 1$ , and  $\beta 3$  are expressed on the surface of blastocysts (Sutherland *et al.*, 1993; Schultz *et al.*, 1997). Antibodies against integrins  $\alpha 5$ ,  $\alpha v$ ,  $\beta 1$ , and  $\beta 3$  block adhesion of blastocysts to fibronectin, an early event in trophoblast differentiation; therefore,  $\alpha v\beta 3$  integrin-mediated adhesion may be important in trophoblast differentiation and function (Schultz and Armant, 1995). After the embryo implants into the uterus, integrin localization changes. Contact with uterine ECM induces protein trafficking of the laminin receptors  $\alpha 5\beta 1$  integrin and  $\alpha 7\beta 1$  integrin from the basal surface to the apical surface of the trophoblast (Schultz *et al.*, 1997; Klaffky *et al.*, 2001; Sutherland, 2003). Trafficking of integrins in trophoblasts seems to be an important regulatory step in trophoblast differentiation.

Placental development is coordinated with invasion into maternal decidua. In preeclampsia, cytotrophoblast invasion is greatly compromised. Preeclampsia is a pregnancy-associated disease that affects 4–5% of all pregnancies and can be fatal (VanWijk *et al.*, 2000). Hypoxia may affect placental invasion in vivo because preeclamptic placentas are hypoxic and exhibit reduced invasion into maternal deciduae (Van-



**Figure 7.** Adhesion to vitronectin and migration toward vitronectin is increased in B16F0 melanoma cells cultured in 1.5% O<sub>2</sub> versus 20% O<sub>2</sub>, via increased cell surface expression of αvβ3. (A) An increase in adhesion to FN and VN is observed in hypoxic B16F0 cells. Adhesion to LN is not changed by hypoxia. (B) Migration toward vitronectin is also enhanced by hypoxia in F0 cells. (C) Adhesion to vitronectin under normoxia and hypoxia is mediated by the β3 integrin because a blocking antibody for β3 integrin diminishes adhesion to vitronectin. (D) B16F0 cells exhibit increased αvβ3 integrin surface expression when treated with 1.5%O<sub>2</sub> for 24 h. Surface levels of α6 and β1 integrins are unaltered by hypoxia. FACS analysis for integrins αv, β3, α6, and β1 was performed on cells cultured in either 20% O<sub>2</sub> (solid line) or 1.5%O<sub>2</sub> (dotted line). (E) B16F0 cells were surface labeled with biotin and immunoprecipitated for αv integrin or β1 integrin. IPs were separated by nonreducing SDS-PAGE and blotted with streptavidin-HRP. Positions of αv integrin, β3 integrin, and β1 integrin are indicated. Error bars represent ± SEM. Student's *t* tests were performed to determine statistical significance.

Wijk *et al.*, 2000; Goldman-Wohl and Yagel, 2002). Pre-eclamptic placentas express low levels of β3 integrins throughout the placenta compared with normal placentas (Zhou *et al.*, 1997a), suggesting that this integrin may be important in proper placental progression. Furthermore, αvβ3 integrin and αIIbβ3 integrin regulate migration and trophoblast outgrowth on fibronectin (Rout *et al.*, 2004). Interestingly, β3 integrin is expressed throughout the murine placenta. However, the subcellular localization of β3 integrin is different for distinct trophoblast subtypes (Bowen and Hunt, 1999). β3 integrin is localized to the membrane in labyrinthine trophoblasts, whereas it is diffuse and cytoplasmic in spongiotrophoblasts (Bowen and Hunt, 1999), suggesting that β3 integrin localization and expression may be a developmentally regulated event. β3 integrin is not essen-

tial for placentation because integrin β3<sup>-/-</sup> mice are viable; however, β3<sup>-/-</sup> mice exhibit reduced fecundity. β3 integrin function is likely important for normal pregnancy, because the viability of integrin β3<sup>-/-</sup> embryos is reduced due to placental defects (Hodivala-Dilke *et al.*, 1999). Of note, 80% of αv integrin-null embryos die at midgestation with placental defects such as a reduced labyrinthine layer, fewer intermingled fetal and maternal blood vessels, and aberrant spongiotrophoblast and giant cell layers (Bader *et al.*, 1998). We have shown that HIF activity is critical to proper establishment of each placental layer (Cowden Dahl, Mack, Compennolle, Adelman, Carmeliet, and Simon, unpublished data). HIF may regulate trophoblast differentiation and invasion, in part, through regulation of αvβ3 integrin. Importantly, TS cells differentiated for 4 d also fail to express high

levels of cell surface  $\alpha v\beta 3$  integrin (our unpublished data), suggesting that  $\alpha v\beta 3$  integrin is not regulated properly during differentiation either. These results strongly suggest that misexpression of  $\alpha v\beta 3$  may occur in HIF-deficient placentas. Integrin-mediated defects are not as severe as those noted for HIF-deficient placentas, likely due to misexpression of additional target genes in HIF mutants. In summary, HIF regulation of genes critical to placentation in addition to the timing of  $\alpha v\beta 3$  integrin surface expression contributes to placental invasion of maternal tissue.

HIF is critical to trophoblast adhesion and migration and is also important in these processes in a variety of disease states. For example, hypoxia and oxidative stress are major factors in inflammation, cardiovascular disease, and cancer (Paul *et al.*, 2004; Peyssonau and Johnson, 2004). HIF is critical to inflammatory responses because myeloid cells lacking HIF1 $\alpha$  are defective in migration and invasion (Cramer *et al.*, 2003). Similar to our findings in TS cells,  $\alpha v\beta 3$  integrin may be important in the normal oxygen-regulated inflammatory responses of macrophages. For example, macrophage differentiation is marked by increased expression of  $\alpha v\beta 3$  integrin (Lafrenie *et al.*, 2002).

### Hypoxia Promotes Tumor Cell Motility

As tumors expand, they can outgrow their vascular supply, resulting in regions of the tumor deficient in O<sub>2</sub> and nutrients. Induction of genes in this hypoxic environment is important for further tumor progression. For example, VEGF induction in melanomas enhances both tumor angiogenesis and metastasis (Rofstad and Danielsen, 1999). In a variety of human cancer, tumor hypoxia is associated with increased metastatic potential. Although many mechanisms are involved in this process, changes in integrin activity may contribute to metastatic progression. We propose that one way tumor cells become more motile is by hypoxic exposure, because hypoxic exposure enhances  $\alpha v\beta 3$  integrin surface expression and adhesion to vitronectin in poorly metastatic B16F0 cells. Additionally, modulation of  $\alpha v\beta 3$  integrin expression could have consequences for tumor angiogenesis, which is linked to  $\alpha v\beta 3$  integrin. To this end, integrin inhibitors are currently in clinical trials as cancer therapeutics.

In summary, we demonstrate that HIF activity is a critical regulator of undifferentiated TS cell adhesion and migration. Functional inactivation of HIF results in decreased  $\alpha v\beta 3$  integrin cell surface expression.  $\alpha v\beta 3$  integrin is not appropriately transported to the cell surface of *Arnt*<sup>-/-</sup> TS cells, which adhere poorly to  $\alpha v\beta 3$  integrin ligands such as vitronectin and fibronectin. Furthermore, hypoxia regulates adhesion and migration of mouse melanoma cells through the regulation of  $\alpha v\beta 3$  integrin expression at the plasma membrane. These data illustrate that HIF activity controls adhesion, migration, and invasion through regulation of a key integrin in both development and cancer.

### ACKNOWLEDGMENTS

We thank Q. C. Yu and Neelima Shah for work on the immuno-EM experiments. This work is supported by National Research Service Award predoctoral fellowship award F31 HL10378 (to K.D.C.D.), National Institutes of Health grant R01 66331, and the Abramson Family Cancer Research Institute (to K.D.C.D. and M.C.S.). M.C.S. is an investigator for the Howard Hughes Medical Institute.

### REFERENCES

Abbott, B. D., and Buckalew, A. R. (2000). Placental defects in ARNT-knock-out conceptus correlate with localized decreases in VEGF-R2, Ang-1, and Tie-2. *Dev. Dyn.* 219, 526–538.

Adelman, D. M., Gertsenstein, M., Nagy, A., Simon, M. C., and Maltepe, E. (2000). Placental cell fates are regulated in vivo by HIF-mediated hypoxia responses. *Genes Dev.* 14, 3191–3203.

Adelman, D. M., Maltepe, E., and Simon, M. C. (1999). Multilineage embryonic hematopoiesis requires hypoxic ARNT activity. *Genes Dev.* 13, 2478–2483.

Bader, B. L., Rayburn, H., Crowley, D., and Hynes, R. O. (1998). Extensive vasculogenesis, angiogenesis, and organogenesis precede lethality in mice lacking all  $\alpha v$  integrins. *Cell* 95, 507–519.

Bashour, A. M., and Bloom, G. S. (1998). 58K, a microtubule-binding Golgi protein, is a formiminotransferase cyclodeaminase. *J. Biol. Chem.* 273, 19612–19617.

Bowen, J. A., and Hunt, J. S. (1999). Expression of cell adhesion molecules in murine placentas and a placental cell line. *Biol. Reprod.* 60, 428–434.

Carreiras, F., Denoux, Y., Staedel, C., Lehmann, M., Sichel, F., and Gauduchon, P. (1996). Expression and localization of  $\alpha v$  integrins and their ligand vitronectin in normal ovarian epithelium and in ovarian carcinoma. *Gynecol. Oncol.* 62, 260–267.

Ceradini, D. J., Kulkarni, A. R., Callaghan, M. J., Tepper, O. M., Bastidas, N., Kleinman, M. E., Capla, J. M., Galiano, R. D., Levine, J. P., and Gurtner, G. C. (2004). Progenitor cell trafficking is regulated by hypoxic gradients through HIF-1 induction of SDF-1. *Nat. Med.* 10, 858–864.

Compennolle, V., *et al.* (2002). Loss of HIF-2 $\alpha$  and inhibition of VEGF impair fetal lung maturation, whereas treatment with VEGF prevents fatal respiratory distress in premature mice. *Nat. Med.* 8, 702–710.

Compennolle, V., Brusselmans, K., Franco, D., Moorman, A., Dewerchin, M., Collen, D., and Carmeliet, P. (2003). Cardia bifida, defective heart development and abnormal neural crest migration in embryos lacking hypoxia-inducible factor-1 $\alpha$ . *Cardiovasc. Res.* 60, 569–579.

Cramer, T., Yamanishi, Y., Clausen, B. E., Forster, I., Pawlinski, R., Mackman, N., Haase, V. H., Jaenisch, R., Corr, M., Nizet, V., *et al.* (2003). HIF-1 $\alpha$  is essential for myeloid cell-mediated inflammation. *Cell* 112, 645–657.

Ding, Z. M., *et al.* (1999). Relative contribution of LFA-1 and Mac-1 to neutrophil adhesion and migration. *J. Immunol.* 163, 5029–5038.

Ema, M., Taya, S., Yokotani, N., Sogawa, K., Matsuda, Y., and Fujii-Kuriyama, Y. (1997). A novel bHLH-PAS factor with close sequence similarity to hypoxia-inducible factor 1 $\alpha$  regulates the VEGF expression and is potentially involved in lung and vascular development. *Proc. Natl. Acad. Sci. USA* 94, 4273–4278.

Felding-Habermann, B., Fransvea, E., O'Toole, T. E., Manzuk, L., Faha, B., and Hensler, M. (2002). Involvement of tumor cell integrin  $\alpha v\beta 3$  in hematogenous metastasis of human melanoma cells. *Clin. Exp. Metastasis.* 19, 427–436.

Firth, J. D., Ebert, B. L., Pugh, C. W., and Ratcliffe, P. J. (1994). Oxygen-regulated control elements in the phosphoglycerate kinase 1 and lactate dehydrogenase A genes: similarities with the erythropoietin 3' enhancer. *Proc. Natl. Acad. Sci. USA* 91, 6496–6500.

Forsythe, J. A., Jiang, B. H., Iyer, N. V., Agani, F., Leung, S. W., Koos, R. D., and Semenza, G. L. (1996). Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol. Cell. Biol.* 16, 4604–4613.

Gehlsen, K. R., Davis, G. E., and Sriramapo, P. (1992). Integrin expression in human melanoma cells with differing invasive and metastatic properties. *Clin. Exp. Metastasis* 10, 111–120.

Giancotti, F. G., and Ruoslahti, E. (1999). Integrin signaling. *Science* 285, 1028–1032.

Goldman-Wohl, D., and Yagel, S. (2002). Regulation of trophoblast invasion: from normal implantation to pre-eclampsia. *Mol. Cell. Endocrinol.* 187, 233–238.

Gu, Y. Z., Hogenesch, J. B., and Bradfield, C. A. (2000). The PAS superfamily: sensors of environmental and developmental signals. *Annu. Rev. Pharmacol. Toxicol.* 40, 519–561.

Hodivala-Dilke, K. M., McHugh, K. P., Tsakiris, D. A., Rayburn, H., Crowley, D., Ullman-Cullere, M., Ross, F. P., Coller, B. S., Teitelbaum, S., and Hynes, R. O. (1999).  $\beta 3$ -Integrin-deficient mice are a model for Glanzmann thrombasthenia showing placental defects and reduced survival. *J. Clin. Investig.* 103, 229–238.

Iwaki, T., Yamamoto, K., Matsuura, T., Sugimura, M., Kobayashi, T., and Kanayama, N. (2004). Alteration of integrins under hypoxic stress in early placenta and choriocarcinoma cell line BeWo. *Gynecol. Obstet. Investig.* 57, 196–203.

Iyer, N. V., *et al.* (1998). Cellular and developmental control of O<sub>2</sub> homeostasis by hypoxia-inducible factor 1 $\alpha$ . *Genes Dev.* 12, 149–162.

- Jin, H., and Varner, J. (2004). Integrins: roles in cancer development and as treatment targets. *Br. J. Cancer* 90, 561–565.
- Klaffky, E., Williams, R., Yao, C. C., Ziober, B., Kramer, R., and Sutherland, A. (2001). Trophoblast-specific expression and function of the integrin  $\alpha 7$  subunit in the peri-implantation mouse embryo. *Dev. Biol.* 239, 161–175.
- Koike, T., *et al.* (2004). Hypoxia induces adhesion molecules on cancer cells: a missing link between Warburg effect and induction of selectin-ligand carbohydrates. *Proc. Natl. Acad. Sci. USA* 101, 8132–8137.
- Kong, T., Eltzschig, H. K., Karhausen, J., Colgan, S. P., and Shelley, C. S. (2004). Leukocyte adhesion during hypoxia is mediated by HIF-1-dependent induction of  $\beta 2$  integrin gene expression. *Proc. Natl. Acad. Sci. USA* 101, 10440–10445.
- Kozak, K. R., Abbott, B., and Hankinson, O. (1997). ARNT-deficient mice and placental differentiation. *Dev. Biol.* 191, 297–305.
- Lartigau, E., Randrianarivelo, H., Avril, M. F., Margulis, A., Spatz, A., Eschwege, F., and Guichard, M. (1997). Intratumoral oxygen tension in metastatic melanoma. *Melanoma Res.* 7, 400–406.
- Liu, Y., Cox, S. R., Morita, T., and Kourembanas, S. (1995). Hypoxia regulates vascular endothelial growth factor gene expression in endothelial cells. Identification of a 5' enhancer. *Circ. Res.* 77, 638–643.
- Maltepe, E., Schmidt, J. V., Baunoch, D., Bradfield, C. A., and Simon, M. C. (1997). Abnormal angiogenesis and responses to glucose and oxygen deprivation in mice lacking the protein ARNT. *Nature* 386, 403–407.
- Nejjari, M., *et al.* (2002). Expression, regulation, and function of  $\alpha V$  integrins in hepatocellular carcinoma: an in vivo and in vitro study. *Hepatology* 36, 418–426.
- Nip, J., Shibata, H., Loskutoff, D. J., Cheresch, D. A., and Brodt, P. (1992). Human melanoma cells derived from lymphatic metastases use integrin  $\alpha v \beta 3$  to adhere to lymph node vitronectin. *J. Clin. Investig.* 90, 1406–1413.
- Okino, S. T., Chichester, C. H., and Whitlock, J. P., Jr. (1998). Hypoxia-inducible mammalian gene expression analyzed in vivo at a TATA-driven promoter and at an initiator-driven promoter. *J. Biol. Chem.* 273, 23837–23843.
- Paul, S. A., Simons, J. W., and Mabeesh, N. J. (2004). HIF at the crossroads between ischemia and carcinogenesis. *J. Cell. Physiol.* 200, 20–30.
- Peng, J., Zhang, L., Drysdale, L., and Fong, G. H. (2000). The transcription factor EPAS-1/hypoxia-inducible factor 2 $\alpha$  plays an important role in vascular remodeling. *Proc. Natl. Acad. Sci. USA* 97, 8386–8391.
- Pennacchietti, S., Michieli, P., Galluzzo, M., Mazzone, M., Giordano, S., and Comoglio, P. M. (2003). Hypoxia promotes invasive growth by transcriptional activation of the met protooncogene. *Cancer Cell* 3, 347–361.
- Peyssonaux, C., and Johnson, R. S. (2004). An unexpected role for hypoxic response: oxygenation and inflammation. *Cell Cycle* 3, 168–171.
- Rodriguez-Fernandez, J. L. (1999). Why do so many stimuli induce tyrosine phosphorylation of FAK? *Bioessays* 21, 1069–1075.
- Rofstad, E. K., and Danielsen, T. (1999). Hypoxia-induced metastasis of human melanoma cells: involvement of vascular endothelial growth factor-mediated angiogenesis. *Br. J. Cancer* 80, 1697–1707.
- Rout, U. K., Wang, J., Paria, B. C., and Armant, D. R. (2004).  $\alpha 5 \beta 1$ ,  $\alpha V \beta 3$  and the platelet-associated integrin  $\alpha IIb \beta 3$  coordinately regulate adhesion and migration of differentiating mouse trophoblast cells. *Dev. Biol.* 268, 135–151.
- Ryan, H. E., Lo, J., and Johnson, R. S. (1998). HIF-1 $\alpha$  is required for solid tumor formation and embryonic vascularization. *EMBO J.* 17, 3005–3015.
- Schultz, J. F., and Armant, D. R. (1995).  $\beta 1$ - and  $\beta 3$ -Class integrins mediate fibronectin binding activity at the surface of developing mouse peri-implantation blastocysts. Regulation by ligand-induced mobilization of stored receptor. *J. Biol. Chem.* 270, 11522–11531.
- Schultz, J. F., Mayernik, L., Rout, U. K., and Armant, D. R. (1997). Integrin trafficking regulates adhesion to fibronectin during differentiation of mouse peri-implantation blastocysts. *Dev. Genet.* 21, 31–43.
- Semenza, G. L., Jiang, B. H., Leung, S. W., Passantino, R., Concordet, J. P., Maire, P., and Giallongo, A. (1996). Hypoxia response elements in the aldolase A, enolase 1, and lactate dehydrogenase A gene promoters contain essential binding sites for hypoxia-inducible factor 1. *J. Biol. Chem.* 271, 32529–32537.
- Staller, P., Sulitkova, J., Lisztwan, J., Moch, H., Oakeley, E. J., and Krek, W. (2003). Chemokine receptor CXCR4 downregulated by von Hippel-Lindau tumour suppressor pVHL. *Nature* 425, 307–311.
- Sutherland, A. (2003). Mechanisms of implantation in the mouse: differentiation and functional importance of trophoblast giant cell behavior. *Dev. Biol.* 258, 241–251.
- Sutherland, A. E., Calarco, P. G., and Damsky, C. H. (1993). Developmental regulation of integrin expression at the time of implantation in the mouse embryo. *Development* 119, 1175–1186.
- Suzuma, K., Takagi, H., Otani, A., and Honda, Y. (1998). Hypoxia and vascular endothelial growth factor stimulate angiogenic integrin expression in bovine retinal microvascular endothelial cells. *Investig. Ophthalmol. Vis. Sci.* 39, 1028–1035.
- Tanaka, S., Kunath, T., Hadjantonakis, A. K., Nagy, A., and Rossant, J. (1998). Promotion of trophoblast stem cell proliferation by FGF4. *Science* 282, 2072–2075.
- Tian, H., Hammer, R. E., Matsumoto, A. M., Russell, D. W., and McKnight, S. L. (1998). The hypoxia-responsive transcription factor EPAS1 is essential for catecholamine homeostasis and protection against heart failure during embryonic development. *Genes Dev.* 12, 3320–3324.
- VanWijk, M. J., Kublickiene, K., Boer, K., and VanBavel, E. (2000). Vascular function in preeclampsia. *Cardiovasc. Res.* 47, 38–48.
- Vaupel, P., Kelleher, D. K., and Hockel, M. (2001). Oxygen status of malignant tumors: pathogenesis of hypoxia and significance for tumor therapy. *Semin. Oncol.* 28, 29–35.
- Walton, H. L., Corjay, M. H., Mohamed, S. N., Mousa, S. A., Santomenna, L. D., and Reilly, T. M. (2000). Hypoxia induces differential expression of the integrin receptors  $\alpha(v\beta 3)$  and  $\alpha(v\beta 5)$  in cultured human endothelial cells. *J. Cell. Biochem.* 78, 674–680.
- Wang, G. L., and Semenza, G. L. (1993). Characterization of hypoxia-inducible factor 1 and regulation of DNA binding activity by hypoxia. *J. Biol. Chem.* 268, 21513–21518.
- Wong, N. C., Mueller, B. M., Barbas, C. F., Ruminski, P., Quaranta, V., Lin, E. C., and Smith, J. W. (1998).  $\alpha v$  integrins mediate adhesion and migration of breast carcinoma cell lines. *Clin. Exp. Metastasis* 16, 50–61.
- Zhou, Y., Damsky, C. H., and Fisher, S. J. (1997a). Preeclampsia is associated with failure of human cytotrophoblasts to mimic a vascular adhesion phenotype. One cause of defective endovascular invasion in this syndrome? *J. Clin. Investig.* 99, 2152–2164.
- Zhou, Y., Fisher, S. J., Janatpour, M., Genbacev, O., Dejana, E., Wheelock, M., and Damsky, C. H. (1997b). Human cytotrophoblasts adopt a vascular phenotype as they differentiate. A strategy for successful endovascular invasion? *J. Clin. Investig.* 99, 2139–2151.