October 2002

Effect of Shear Stress on Platelet Activation via the Glycoprotein VI Receptor

Kendra L. Sarratt

University of Pennsylvania, ksarratt@seas.upenn.edu

Hong M. Chen

University of Pennsylvania, hong2@mail.med.upenn.edu

Mark L. Kahn

University of Pennsylvania, markkahn@mail.med.upenn.edu

Daniel A. Hammer

University of Pennsylvania, hammer@seas.upenn.edu

Follow this and additional works at: https://repository.upenn.edu/be_papers

Recommended Citation

Sarratt, K. L., Chen, H. M., Kahn, M. L., & Hammer, D. A. (2002). Effect of Shear Stress on Platelet Activation via the Glycoprotein VI Receptor. Retrieved from https://repository.upenn.edu/be_papers/38

Copyright 2002 IEEE. Reprinted from *Proceedings of the Second Joint Engineering in Medicine and Biology Society/ Biomedical Engineering Society Conference, (EMBS/BMES 2002)*, Volume 1, pages 674-675.

Publisher URL: <http://ieeexplore.ieee.org/xpl/tocresult.jsp?isNumber=25190&page=22>

This material is posted here with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the University of Pennsylvania's products or services. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org. By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

This paper is posted at ScholarlyCommons. https://repository.upenn.edu/be_papers/38
For more information, please contact repository@pobox.upenn.edu.

Effect of Shear Stress on Platelet Activation via the Glycoprotein VI Receptor

Abstract

Cardiovascular diseases are the nation's leading cause of death. Such diseases are caused by platelet response to collagen especially in the event of vascular injury leading to thrombosis. One of the platelet receptors known to bind to the collagen ligand is glycoprotein VI (GPVI) with co-receptor Fc receptor γ chain (FcR γ). By stably expressing the GPVI receptor in rat basophilic leukemia cells (RBL-2H3), which abundantly express FcR γ , but endogenously lack GPVI, studies have shown that GPVI-FcR γ is sufficient to confer adhesion as well as signaling responses to collagen as long as the receptor density is equivalent to that found on human platelets. While those investigations confirm that the GPVI receptor mediate binding to collagen under static conditions, they do not provide information on how the GPVI receptor interacts with collagen under dynamic conditions. In the present study we have used the GPVI-expressing RBL-2H3 cells to observe the kinetics of adhesion to collagen under hydrodynamic flow conditions *in vitro* using a parallel plate flow chamber coupled with video microscopy. We demonstrate that these cells do adhere to the surface at a low shear rate and do so at a greater adherent cell density than wild-type RBL-2H3 (WT-RBL) cells.

Keywords

GPVI-expressing RBL-2H3 cells, GPVI-173, collagen, cell adhesion, flow chamber

Comments

Copyright 2002 IEEE. Reprinted from *Proceedings of the Second Joint Engineering in Medicine and Biology Society/Biomedical Engineering Society Conference, (EMBS/BMES 2002)*, Volume 1, pages 674-675.

Publisher URL: <http://ieeexplore.ieee.org/xpl/tocresult.jsp?isNumber=25190&page=22>

This material is posted here with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the University of Pennsylvania's products or services. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org. By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

EFFECT OF SHEAR STRESS ON PLATELET ACTIVATION VIA THE GLYCOPROTEIN VI RECEPTOR

K. L. Sarratt¹, H. M. Chen², M. L. Kahn³, D. A. Hammer⁴

^{1,4}Department of Bioengineering, University of Pennsylvania, Philadelphia, PA, USA

^{2,3}Department of Cardiovascular Medicine, University of Pennsylvania, Philadelphia, PA, USA

Abstract-Cardiovascular diseases are the nation's leading cause of death. Such diseases are caused by platelet response to collagen especially in the event of vascular injury leading to thrombosis. One of the platelet receptors known to bind to the collagen ligand is glycoprotein VI (GPVI) with co-receptor Fc receptor gamma chain (FcRγ). By stably expressing the GPVI receptor in rat basophilic leukemia cells (RBL-2H3), which abundantly express FcRγ, but endogenously lack GPVI, studies have shown that GPVI-FcRγ is sufficient to confer adhesion as well as signaling responses to collagen as long as the receptor density is equivalent to that found on human platelets. While those investigations confirm that the GPVI receptor mediate binding to collagen under static conditions, they do not provide information on how the GPVI receptor interacts with collagen under dynamic conditions. In the present study we have used the GPVI-expressing RBL-2H3 cells to observe the kinetics of adhesion to collagen under hydrodynamic flow conditions *in vitro* using a parallel plate flow chamber coupled with video microscopy. We demonstrate that these cells do adhere to the surface at a low shear rate and do so at a greater adherent cell density than wild-type RBL-2H3 (WT-RBL) cells.

Keywords - GPVI-expressing RBL-2H3 cells, GPVI-173, collagen, cell adhesion, flow chamber

I. INTRODUCTION

One of the primary and critical reactions that can lead to cardiovascular disorders is the adherence and aggregation of platelets at sites of vascular injury in flowing blood [1,2]. Platelet response is generally brought about through interactions between specific receptors on the platelet surface and exposed ligands of the subendothelium, with collagen being a major agonist. The initial interaction between circulating platelets and the subendothelium is believed to occur indirectly through the von Willebrand factor (vWF) bound to collagen and the GPIb platelet receptor [3]. Other platelet surface proteins implicated as being direct platelet/collagen receptors include integrin $\alpha_2\beta_1$, CD36, glycoprotein VI (GPVI), and P65 [4-8].

Although the indirect interaction has been clearly identified, the direct interaction between platelet receptors and collagen remain poorly understood, especially in the case of the GPVI receptor [9-11]. This receptor has been found to be important, since human platelets deficient in it show impaired levels of adhesion and aggregation compared to whole blood [1]. To address the functional role of GPVI in isolation of other receptors, GPVI was expressed in rat basophilic leukemia (RBL-2H3) cells, a cell line that expresses the Fc receptor gamma chain (FcRγ), but does not express endogenous GPVI or integrin $\alpha_2\beta_1$ [12]. Studies with these cells under static conditions demonstrate that "GPVI expression is sufficient to confer both adhesive and

signaling responses to collagen independent of other collagen receptors but that GPVI-collagen responses are strictly dependent on receptor density" [11]. Of the three clones studied, GPVI-expressing RBL clone 173 (GPVI-173), expressing the highest level of GPVI, adhered to collagen at a higher optical density than the other two clones [11]. Using the GPVI-173 cells, this study quantifies the adhesion of the cells to collagen type I under flow conditions. The preliminary results show that the GPVI receptor adheres to the collagen surface at a low shear rate.

II. METHODOLOGY

Slide preparation. Microscope slides (3"x1"; Sigma Diagnostics, St. Louis, MO) were coated with fibrillar collagen (100 ug/mL) derived from equine tendons (Chronolog; Havertown, PA). The slide was incubated overnight at 4°C. On the day of experimentation the slide was rinsed with phosphate buffered saline with 0.05% polyoxyethylene-sorbitan monolaurate to remove unbound collagen. The slide was then incubated with 2% bovine serum albumin (BSA) for 1 hour at room temperature on a rotator.

Cell preparation. GPVI-173 cells were generated as previously described. [11,12]. The cells were cultured in Dulbecco Modified Eagle Medium with 10% fetal bovine serum and 10μg/mL gentamicin. On the day of experimentation, the cells were suspended in PBS with 1% BSA (PBS⁺) to achieve a cell concentration of 1×10^6 cells/mL.

Parallel plate flow chamber assembly and video microscopy. The parallel plate flow chamber was assembled in water with a straight channel template and collagen coated glass slide. The flow chamber assembly was mounted onto the stage of a phase contrast inverted microscope (Diaphot-TMD; Nikon, Tokyo, Japan). A CCD camera (Cohu Inc.; San Diego, CA) and Sony VHS recorder (Sony Medical Systems; Montvale, NJ) were used to obtain and record images. After the chamber was flushed with PBS⁺, wild-type RBL-2H3 (WT-RBL) and GPVI-173 cells were passed through the flow chamber at a controlled flow rate for 20 minutes using a syringe pump (Model 55-1143, Harvard Apparatus; South Natic, MA). The actual shear stress was calculated by the following equation

$$\tau_w = 6\mu Q/wh^2 \quad (1)$$

where τ_w is the wall shear stress, μ is the fluid viscosity, Q is the volumetric flow rate, w is the chamber width, and h is the gap thickness [13,14].

Adherent cell densities were determined by counting the number of adherent cells and dividing them by the field of view, 0.32 mm². Based on three experiments, one-way analysis of variance with a probability value of 0.05 (95% confidence) was performed to test for significance.

III. RESULTS

Visually analyzing the video, the adhesion of GPVI-173 to collagen was studied at a wall shear rate of 63s⁻¹. The number of cells per 0.32 mm² was counted and averaged over 3 experiments. We have presented the average adherent cell densities of GPVI-173 and WT-RBL cells at 5, 10, 15, and 20 minutes (see Table I). It was found that there was significant adhesion of GPVI-173 cells to collagen, with the cells remaining adherent. Over the twenty-minute segment, the adherent cell density gradually increased for both cell types. Control experiments performed with WT-RBL cells showed significantly lower adherent cell densities compared to that of GPVI-173 cells.

IV. DISCUSSION

Studies using static assays show that GPVI-173 is sufficient to mediate adhesion to collagen. While this may be true, the results do not account for the fact that adhesion of platelets, *in vivo*, occur under physiologically shear conditions. We demonstrate that at a low shear rate of 63s⁻¹, GPVI-173 cells adhere to collagen with a significantly higher adherent cell density than WT-RBL cells. This result suggests that GPVI alone is sufficient to confer adhesion to collagen under hydrodynamic conditions. However the shear rate range over which the receptor is reactive has yet to be determined. Thus, more experiments must be done to test for nonspecific adhesion.

V. CONCLUSION

This study provides further evidence that GPVI, a collagen receptor, mediates collagen adhesion. In doing the adhesion assay under dynamic conditions, we hope to gain a better understanding of the mechanistic details of the GPVI receptor's involvement in collagen activation of platelets. By determining its functionality, we plan to elucidate the pathomechanism GPVI plays in cardiovascular disorders such as myocardial infarction, angina, stroke, and arteriosclerosis. From there future investigations can provide novel targets for anti-platelet strategies to treat cardiovascular diseases.

ACKNOWLEDGMENTS

We gratefully acknowledge the support of NIH H162250 and HL64388 to Dr. Daniel Hammer.

REFERENCES

- [1] Moroi, M, Jung, SM, Shimmyozu, K, Tomiyama, Y, Ordians, A, Diaz-Ricart, M. Analysis of platelet adhesion to a collagen-coated surface under flow conditions: The involvement of glycoprotein VI in the platelet adhesion. *Blood* 1996; 88:2081-92.
- [2] Savage, B, Almus-Jacobs, F, Ruggeri, ZM. Specific synergy of multiple substrate-receptor interactions in platelet thrombus formation under flow. *Cell* 1998; 94:657-666.
- [3] Kehrel, B, "Collagen Receptors: Biology and Functions" In: Rao, GHR, ed., *Handbook of Platelet Physiology and Pharmacology*, Kluwer Academic Publishers, pp. 80-101. (1999).
- [4] Nieuwenhuis, HK, Akkermann, JWN, Houdijk, WPM, Sixma, JJ. Human blood platelets showing no response to collagen fail to express surface glycoprotein Ia. *Nature* 1985; 318:470-472.
- [5] Staatz, WD, Walsh, JJ, Pexton, T, Santoro, SA. The $\alpha_2\beta_1$ integrin cell surface collagen receptor binds to the $\alpha 1$ (I)-CB3 peptide and collagen. *J Biol Chem* 1990; 265:4778-4781.
- [6] Tandon, NN, Kralisz, U, Jamieson, GA. Identification of glycoprotein IV (CD36) as a primary receptor for platelet-collagen adhesion. *J Biol Chem* 1989; 264:7576-7583.
- [7] Moroi, M, Jung, SM, Okuma, M, Shimmyozu, K. A patient with platelets deficient in glycoprotein VI that lack both collagen-induced aggregation and adhesion. *J Clin Invest* 1989; 84:1440-1445.
- [8] Chaing, TM, Kang, AH. Isolation and purification of collagen $\alpha 1$ (I) receptor from human platelet membrane. *J Biol Chem* 1982; 257:7581-7586.
- [9] Sixma, JJ, van Zanten, GH, Huizinga, EG, van der Plas, RM, Verkley, M, Wu, YP, Gros, P, de Groot, PG. *Thromb Haemostasis* 1997; 78:434-438.
- [10] Barnes, MJ, Knight, CG, Farndale, RW. *Curr Opin Hematol* 1998; 5:314-320.
- [11] Chen, H, Locke, D, Liu, C, Kahn, ML. The platelet receptor GPVI mediates both adhesion and signaling responses to collagen in a receptor density-dependent fashion. *J Biol Chem* 2002; 277:3011-3019.
- [12] Zheng, Y, Liu, C, Chen, H, Locke, D, Ryan, J, Kahn, M. Expression of the platelet receptor GPVI confers signaling via the Fc receptor γ chain in response to the snake venom convulxin but not to collagen. *J Biol Chem* 2001; 276:12999-13006.
- [13] Mohandas, N, Hochmuth, RM, Spaeth, EE. Adhesion of red cells to foreign surfaces in the presence of flow. *J Biomed Mat Res* 1974; 8:119-136.
- [14] Hochmuth, RM, Mohanda, N, Blackshear, PL. Measurement of the elastic modulus for red cell membrane, using a fluid mechanical technique. *Biophys J* 1973; 13:747-762.

Table I. Average adherent cell densities (cells/mm²) $\pm$ SE

	Cell Type	
	GPVI-173	WT-RBL
5 min	87.47 $\pm$ 20.56	45.32 $\pm$ 15.44
10 min	160.72 $\pm$ 33.10	69.03 $\pm$ 15.70
15 min	182.33 $\pm$ 33.94	72.98 $\pm$ 15.33
20 min	192.08 $\pm$ 32.85	75.09 $\pm$ 17.73