



5-2014

## Using Growth Mixture Modeling to Identify Classes of Sodium Adherence in Adults with Heart Failure

Ruth Masterson-Creber

Christopher S. Lee


Terry A. Lennie

Maxim Topaz

Barbara Riegel

*University of Pennsylvania*, [briegel@nursing.upenn.edu](mailto:briegel@nursing.upenn.edu)

Follow this and additional works at: <https://repository.upenn.edu/nrs>

 Part of the [Cardiology Commons](#), [Cardiovascular Diseases Commons](#), [Circulatory and Respiratory Physiology Commons](#), [Health Services Research Commons](#), [Medical Humanities Commons](#), [Nursing Commons](#), and the [Preventive Medicine Commons](#)

---

### Recommended Citation

Masterson-Creber, R., Lee, C. S., Lennie, T. A., Topaz, M., & Riegel, B. (2014). Using Growth Mixture Modeling to Identify Classes of Sodium Adherence in Adults with Heart Failure. *Journal of Cardiovascular Nursing*, 29 (3), 209-217. <http://dx.doi.org/10.1097/JCN.0b013e3182834191>

This paper is posted at ScholarlyCommons. <https://repository.upenn.edu/nrs/108>  
For more information, please contact [repository@pobox.upenn.edu](mailto:repository@pobox.upenn.edu).

---

# Using Growth Mixture Modeling to Identify Classes of Sodium Adherence in Adults with Heart Failure

## Abstract

**BACKGROUND:** The prevention of fluid retention is important to reduce hospitalizations in patients with heart failure (HF). Following a low-sodium diet helps to reduce fluid retention.

**OBJECTIVE:** The primary objective of this study was to use growth mixture modeling to identify distinct classes of sodium adherence-characterized by shared growth trajectories of objectively measured dietary sodium. The secondary objective was to identify patient-level determinants of the nonadherent trajectory.

**METHODS:** This was a secondary analysis of data collected from a prospective longitudinal study of 279 community-dwelling adults with previously or currently symptomatic HF. Growth mixture modeling was used to identify distinct trajectories of change in 24-hour urinary sodium excretion measured at 3 time points over 6 months. Logistic modeling was used to predict membership in observed trajectories.

**RESULTS:** The sample was predominantly male (64%), had a mean age of 62 years, was functionally compromised (59% New York Heart Association class III), and had nonischemic HF etiology. Two distinct trajectories of sodium intake were identified and labeled adherent (66%) and nonadherent (34%) to low-sodium diet recommendations. Three predictors of the nonadherent trajectory were identified, confirming our previous mixed-effect analysis. Compared with being normal weight (body mass index/m<sup>2</sup>), being overweight and obese was associated with a 4-fold incremental increase in the likelihood of being in the nonadherent trajectory (odds ratio [OR], 4.63; 95% confidence interval [CI], 1.66-12.91;  $P < .002$ ). Being younger than 65 years (OR, 4.66; 95% CI, 1.04-20.81;  $P = .044$ ) or having diabetes (OR, 4.15; 95% CI, 1.29-13.40;  $P = .016$ ) were both associated with more than 4 times the odds of being in the nonadherent urine sodium trajectory compared with being older than 65 years or not having diabetes, respectively.

**CONCLUSIONS:** Two distinct trajectories of sodium intake were identified in patients with HF. The nonadherent trajectory was characterized by an elevated pattern of dietary sodium intake shown by others to be associated with adverse outcomes in HF. Predictors of the nonadherent trajectory included higher body mass index, younger age, and diabetes.

## Keywords

Adult, Comorbidity, Diabetes Mellitus, Diet, Sodium-Restricted, Female, Heart Failure, Humans, Male, Middle Aged, Obesity, Patient Compliance, Prospective Studies, Sodium

## Disciplines

Cardiology | Cardiovascular Diseases | Circulatory and Respiratory Physiology | Health Services Research | Medical Humanities | Medicine and Health Sciences | Nursing | Preventive Medicine



Published in final edited form as:

*J Cardiovasc Nurs.* 2014 ; 29(3): 209–217. doi:10.1097/JCN.0b013e3182834191.

## Using Growth Mixture Modeling to Identify Classes of Sodium Adherence in Adults With Heart Failure

**Ruth Masterson Creber, MSc, RN [PhD Student],**  
School of Nursing, University of Pennsylvania, Philadelphia

**Christopher S. Lee, PhD, RN [Assistant Professor],**  
Oregon Health & Science University, Portland

**Terry A. Lennie, PhD, RN, FAHA, FAAN [Professor],**  
College of Nursing, University of Kentucky, Lexington

**Maxim Topaz, MA, RN [PhD Student], and**  
School of Nursing, University of Pennsylvania, Philadelphia

**Barbara Riegel, DNSc, RN, FAHA, FAAN [Professor]**  
School of Nursing, University of Pennsylvania, Philadelphia

### Introduction

Adherence to a low-sodium diet is an important self-care behavior for patients with heart failure (HF). Excess sodium intake promotes higher ventricular filling pressures, pulmonary congestion, clinical symptoms of fluid retention, and puts patients with HF at high risk for acute decompensation and hospitalizations.<sup>1</sup> As such, adherence to a low-sodium diet is a key factor in reducing hospitalizations<sup>2</sup> and one of the most frequently recommended self-care behaviors.<sup>3-6</sup> Both the 2012 European Society of Cardiology and 2009 American College of Cardiology/American Heart Association guidelines recommend less than 2 g/day of dietary sodium.<sup>7,8</sup> Likewise, the Heart Failure Society of America recommends that all patients with clinical symptoms of HF restrict sodium to 2-3 g/day.<sup>6</sup> It is estimated, however, that only about 22-55% of HF patients are adherent to a low-sodium diet.<sup>9-12</sup>

Adherence, as defined by the World Health Organization (WHO), is a multidimensional phenomenon, determined by multiple dimensions related to social and economic conditions, the health care system in addition to condition, therapy or patient-related factors.<sup>13</sup> Like medication adherence in this patient population, adherence to a low sodium diet is influenced by a number of factors.<sup>14</sup> Although several factors have been associated with non-adherence to a low-sodium diet in the adult population with HF, there are limitations to these research findings. Most prior studies on 24-hour urine sodium excretion are cross-sectional in design and conducted with small samples. Deterministic methods, including mixed effects modeling have previously been used to identify patient-level factors associated with high sodium intake using an *a priori* determined cutoff of adherence.<sup>15</sup> This study builds on that analysis by applying a growth mixture modeling methodology to

Correspondence: Ruth Masterson Creber, MSc, RN, University of Pennsylvania, Philadelphia, PA (rumaster@nursing.upenn.edu; ruth.masterson@gmail.com).

The authors have no conflicts of interest to disclose.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

identify the “most likely” trajectory membership and quantify uncertainty in trajectory membership over time.

The specific aim of this study was to use growth mixture modeling to identify distinct patterns of change in 24-hour urine sodium excretion, as a proxy for dietary sodium intake, over six months of observation. The secondary aim was to build on a previous mixed effects analysis<sup>15</sup> and examine patient-level determinants of observed trajectories of urine sodium.

## Methods

### Design and Study population

The methodology of this study has been previously reported in detail.<sup>14</sup> In brief, this was a prospective, observational study of the association between excessive daytime sleepiness and HF self-care among 279 community-dwelling adults with chronic HF. The participants were enrolled from three outpatient settings in Philadelphia, Pennsylvania and Newark, Delaware between 2007 and 2010. Most of the participants were followed in specialty HF clinics and most were on optimal medical therapy. Given the observational nature of this study, the usual care provided by HF specialty clinics was not augmented. Participants did not receive supplemental HF self-care education outside of what was provided as part of routine care.

Inclusion criteria for the study specified adults with previously or currently symptomatic HF confirmed by echocardiographic and clinical evidence.<sup>14</sup> The participants needed to be able to actively participate in the study, so inclusion criteria specified that patients must be able to read and speak English and have sufficient cognition and health literacy to complete the questionnaires. Otherwise eligible individuals were excluded if they lived in a long-term care setting and were not responsible for their own self-care, including preparing and making choices about food. People were also excluded if they had renal failure requiring dialysis, imminent plans to move out of the area, a history of drug or alcohol abuse within the past year or an imminently terminal illness. People with major depressive illness were also excluded because self-care behaviors are influenced by major depression.<sup>16</sup> This study was in compliance with the Declaration of Helsinki.<sup>17</sup> Institutional Review Board approval was obtained from all three sites and all participants gave written informed consent. Data were collected at baseline, 3-and 6-months by research assistants during home visits. Institutional Review Board approval was obtained for this secondary analysis.

### Measurement of dietary sodium

A 24-hour urine sodium collection is considered a reliable and valid estimate of sodium intake<sup>18,19</sup> and has been used as an objective measure of sodium intake in other studies with patients with HF.<sup>1,20</sup> Sodium is under tight homeostatic control. If individuals do not perspire heavily, urine sodium accounts for approximately 95% to 98% of daily dietary sodium intake.<sup>21</sup> Variations in 24-hour urine sodium are primarily attributed to changes in sodium intake over the previous 24 hours.<sup>22</sup> Daily urine sodium levels are estimated to fluctuate only about 11%.<sup>23</sup> Validation for 24-hour urine collection has been performed by urine recovery of oral doses of para-amino benzoic acid.<sup>19,23,24</sup>

The methodology for collecting and measuring dietary sodium intake has been reported elsewhere<sup>15</sup> and is summarized here. Dietary sodium intake was estimated using three 24-hour urine sodium levels measured at enrollment, and at three and six months after enrollment. The procedure for the collection and quantification of the 24-hour urine samples is summarized here. At the time of enrollment, participants and their family members were given training for the specimen collection. Participants were provided with all necessary supplies including: urine containers, collection devices, verbal and written instructions (with

pictures) specifying when they should start and finish collecting the specimen. Prior to each urine collection, patients were given a reminder call and training for the procedure was reinforced by the research assistant. Participants self-reported the start and stop time of the urine collection. If the recorded time was less than 22 hours or greater than 26 hours it was considered inaccurate and excluded from the analysis. Specimens were collected from participant's homes by a courier service shortly after the 24-hour sample was complete.<sup>15</sup>

Urine sodium analyses were completed at the central laboratory at the Hospital of the University of Pennsylvania using the ion-selective electrode method (Beckman LX<sup>®</sup> 20 Chemistry Analyzer, Beckman Coulter, Inc., Fullerton, CA) to determine the amount of sodium in the urine specimens. A normal urinary sodium level is 40 to 120 mEq/L/day.<sup>11</sup> In order to compare urine excretion with the recommended diet of 2-3 g/day sodium intake,<sup>6</sup> urine sodium excretion in millimoles was converted to g ( $g = \text{mmol} * .02299$ ) to estimate dietary sodium intake. As a quality assurance measure, urinary creatinine was measured concurrently. Normal urinary creatinine is  $1 \pm 10\%$  g<sup>25</sup> so any samples with creatinine levels  $<0.9$  g in 24 hours were considered incomplete.<sup>26</sup>

### Statistical Analysis

Standard descriptive (proportions, means and standard deviations), unadjusted odds ratios and *t* tests were used to describe the sample and compare trajectories. Latent growth mixture modeling was used to identify distinct trajectories of urinary sodium using data from three time points.

Latent class growth modeling is used to examine patterns in longitudinal data with repeated measures to identify classes or subgroups within a population. Unique trajectories or pathways differentiate classes from one another. Using repeated measures to identify subgroups within a population has been used across multiple disciplines. Latent class growth mixture analysis also has been used across multiple disciplines to identify trajectories of development;<sup>27</sup> nighttime continence;<sup>28</sup> physical aggression in young boys;<sup>29</sup> the natural history of prostate disease;<sup>30</sup> depression recovery profiles<sup>31</sup> and preventative interventions for reducing classroom violence.<sup>32</sup> It has also been used extensively in the substance abuse literature to identify trajectories and predictors of high-risk behavior.<sup>33-44</sup> Within the heart failure literature, it has been used to identify patterns of cognitive change,<sup>45</sup> as well as patterns of medication nonadherence.<sup>46</sup>

The purpose of growth mixture modeling is to identify homogenous subgroups within larger heterogeneous populations and represent unobserved heterogeneity among subjects in a sample. Finite mixture random effects are used to represent the departure of individuals' latent growth parameters from the population mean growth parameter.<sup>32,47</sup> Class membership of an individual cannot be measured directly; however, it can be inferred based on membership in a specific trajectory.<sup>48</sup> According to Wang and colleagues growth mixture modeling can be described as having three steps. The first step is analogous to random effects modeling which specifies individual-level observed data as a sum of fixed effects, random effects and measurement errors at each observation point.<sup>48</sup> The second step represents the distribution of class-specific random effects and covariate effects on class-specific mean growth trajectories. The third step includes the covariate effects on class membership using regression models.<sup>48</sup>

Growth mixture modeling has many benefits over alternative methods, as reviewed by Preacher.<sup>49</sup> Growth mixture modeling can model changes in factors over time as random effects, allow for variations between individuals and subgroups (instead of individuals and the population average as in mixed-effect modeling) and judge comparative fit between iterative models with various statistics; the focus of growth mixture modeling is on

capturing inter-and-intra-individual differences over time.<sup>50</sup> Conventional deterministic approaches minimize within group and maximize between-group variance. In contrast, growth mixture modeling employs a model-based approach to calculate the probability of membership in each trajectory, identify “most likely” trajectory membership, and quantifies uncertainty in trajectory membership. In this article the terms subgroups and classes are used synonymously to describe people who share a specific growth trajectory.

Growth mixture modeling was performed using MPlus v.6.12 (Los Angeles, CA). Model fit between 2 to 4 trajectories were compared ( $k$  vs  $k-1$  trajectories) using the Lo-Mendell-Rubin adjusted likelihood ratio test ( $p < 0.05$ ), parametric bootstrapped likelihood ratio test ( $p < 0.05$ ), Bayesian Information Criteria (BIC), convergence (entropy closest to 1.0), the proportion of the sample in each trajectory (not less than 5%), and average posterior probabilities (closest 1.0).<sup>50,51</sup> Predictors of observed trajectories of dietary sodium were quantified using logistic regression modeling in MPlus. Unadjusted and multivariate odds ratios (OR) and 95% confidence intervals (CI) were calculated for each model factor, taking into account our method of multiple imputation, described below.

The following ten determinants of non-adherence were tested based on the five dimensions of adherence posed by the WHO.<sup>52</sup> Four of the five WHO adherence dimensions were included in this analysis. The fifth factor was not included because data on the health care system was not available for this secondary data analysis. Patient related factors included gender, age and BMI. Social and economic factors included race, income and highest level of education. Condition related factors included New York Heart Association (NYHA) functional class, HF type (diastolic or systolic), etiology of HF and diabetes as a comorbidity. Therapy related included diuretic use.

### Robust Handling of Missing Data

In an effort to reduce bias, a principled method of multiple imputation<sup>53</sup> was performed to account for missing urine sodium excretion values (23% of the sample had at least one missing value). Data were imputed using the method of chained equations (Stata v. 11.2).<sup>54-56</sup> The method of chained equations is a contemporary technique that imputes missing data based on multiple covariates in the dataset (i.e. a full information method of imputation). The latent class variable was fit on imputed data in Mplus v.6.12. Multiple sensitivity analyses were performed to examine potential differences in the significance and effect size of predictors of urine sodium trajectories comparing complete case with imputed data samples. There were no differences in the predictors identified in the models except for the size of the confidence intervals; thus, for economy of presentation, only the final analysis using imputed data is shown here.

## Results

The overall demographic and clinical characteristics of the sample population overall at baseline are presented in Table 1. This sample of adults ( $n=279$ ) with HF was predominantly male (64%), had a mean age of 62, non-ischemic HF etiology and was functionally compromised (59% NYHA class III). The unadjusted odds ratios for the analysis are presented in Table 2. The covariates that had a statistically significant unadjusted association with the non-adherent trajectory were diabetes ( $p=0.008$ ), income ( $p=0.031$ ) and BMI ( $p < 0.001$ ).

Modeling of three and four trajectories resulted in poor model fit and small trajectories (*data not shown*). Entropy for the two-class model was 0.752 and there were high average posterior probabilities of membership in both trajectories (94% and 91%), indicating very limited uncertainty in trajectory membership. Based on the observed characteristics, we

labeled the first and largest trajectory “adherent” (n=178, 66%) and second trajectory “non-adherent” (n=91, 34%).

As displayed in the Figure there was heterogeneity within each trajectory, as demonstrated by the error bars, which indicate two standard deviations around each mean urine sodium level at each point in time. Paired t-tests demonstrate statistically significant differences between the two trajectories at each time point ( $p < 0.001$ ). Those in the adherent group started off at baseline with a mean sodium consumption of 2.42 g/day (standard deviation (SD)  $\pm 0.19$ , 95% CI: 2.39-2.45) versus 4.41 g/day (SD:  $\pm 0.34$ , 95% CI: 4.33-4.48) in the non-adherent trajectory ( $t = 61.68$ ,  $p < 0.001$ ). At three months, the mean difference between the two trajectories was 2.16 g/day (SE: 0.36,  $t = 59.99$ ,  $p < 0.001$ ) and by six months was 1.98 g/day (SE:  $\pm 0.03$ ,  $t = 62.56$ ,  $p < 0.001$ ). Within the adherent and non-adherent trajectories there were no differences from baseline to 6 months (SE:  $\pm 0.02$ ,  $t = -1.39$ ,  $p = 0.166$  and SE:  $\pm 0.05$ ,  $t = -0.47$ ,  $p = 0.640$ ), indicating that there were limited changes in dietary sodium intake over time.

Based on the WHO model, predictors of membership in the non-adherence trajectory were identified. The conceptual model was also supported using data driven statistical approaches including univariate associations. Compared with normal weight patients (BMI  $< 25$ ), being overweight and obese was associated with a 4-fold incremental increase in the likelihood of being in the non-adherent trajectory (OR: 4.63, 95% CI: 1.66-12.91,  $p < 0.002$ ). Being less than 65 years of age (OR: 4.66, 95% CI: 1.04-20.81,  $p = 0.044$ ) or having diabetes (OR: 4.15, 95% CI: 1.29-13.40,  $p = 0.016$ ) were both associated with over four times the odds of being in the non-adherent urine sodium trajectory compared with people who are over 65 or do not have diabetes, respectively. There were no significant differences in the odds of being in the non-adherent trajectory based on gender, cognition, diuretic use, income level, education, race, HF etiology, HF type or NYHA functional status.

## Discussion

The primary aim of this study was to identify trajectories of sodium adherence in a sample of community dwelling HF patients and two groups were identified. Based on the more liberal Heart Failure Society of America sodium recommendations of 3g/day sodium intake,<sup>6</sup> one trajectory was labeled “adherent” and the other “non-adherent” to a low sodium diet. Both groups were consistently adherent or non-adherent over six months of follow-up. That is, there was no change in the level of adherence or non-adherence over time.

To the best of our knowledge, this is the first study that describes adherence to a low-sodium diet using a growth mixture approach; however, this method has been applied to medication adherence in this study sample.<sup>46</sup> The predictors of the non-adherence to a low sodium diet were being overweight or obese, age less than 65 years and having diabetes. These results that obesity and diabetes predict higher sodium consumption are consistent with an analysis that employed a mixed-effect analytic approach;<sup>15</sup> however, in this study we also identified younger age as a risk factor for being non-adherent to a sodium restricted diet.

HF patients who were overweight or obese, age less than 65 years or have diabetes were identified as being more likely to have trouble following a restricted sodium diet and may need more focused, tailored self-care interventions.<sup>57</sup> One approach to tailoring self-care interventions that we are testing in a pilot randomized control study is a tailored motivational interviewing and skill-building approach to improving self-care behaviors including adherence to a low-sodium diet.

The finding that being obese is strongly associated with greater odds of being non-adherent to a restricted sodium diet implies that these patients may be consuming more food and by default more sodium, or that they are specifically consuming more foods that are higher in sodium. This finding is consistent with the Intersalt international study of electrolyte excretion and blood pressure, in which investigators reported a positive correlation between BMI and sodium excretion.<sup>19</sup> They proposed that overweight individuals consuming higher sodium intake most likely explains this finding.<sup>58</sup> In contrast, Chung et al<sup>20</sup> found no differences in BMI between people who self-reported being adherent or non-adherent to low-sodium guidelines. The approach in our study is different than that used by Chung and colleagues in that we did not apply a 2,000 mg cutoff for sodium adherence. Instead, we identified distinct and naturally occurring trajectories of urinary sodium excretion that happened to track with current dietary sodium recommendations. Moreover, the current study was longitudinal in design. Thus, methodological differences may explain our incongruent findings.

Other authors studying sodium adherence in the HF population have not reported our finding of an association between younger age and non-adherence to a low sodium diet. In studies of adherence to self-maintenance behaviors, an association between age and adherence to medication has been reported. Several studies report that younger patients are more adherent to taking medication<sup>59,60</sup> while others found no consistent difference in medication adherence by age.<sup>61-63</sup> In this specific sample, we found no difference in overall medication adherence by age.<sup>64</sup> The association with younger age and the non-adherent trajectory could be that younger patients with HF may also be more newly diagnosed and not yet adept in the skills of identifying and avoiding hidden sodium in foods. Patients with HF who are over 65 years may be living that long because they have been more stringent with the self-maintenance behavior of eating a low salt diet.

The finding that diabetes is a strong predictor of the non-adherence to a low sodium diet is most likely explained by overconsumption of sodium, rather than a physiologic mechanism. In fact, in the diabetic state of hyperinsulinemia, sodium reabsorption is enhanced in the nephron segments of the kidney.<sup>65-68</sup> Clinically, patients with both HF and diabetes may struggle to closely monitor both their sugar and their sodium intake. There may receive conflicting messages from multiple providers about nutritional priorities. The implications of this finding may be that interventions are needed to specifically address the dietary complexities of managing both HF and diabetes in this patient population.

Several factors that we anticipated would predict sodium adherence did not reach statistical significance, including higher income and education. We expected that people who reported having inadequate income would potentially be more likely to be non-adherent to a low sodium diet because low-sodium food is more expensive and time consuming to prepare than processed, higher-sodium foods; however, there were no differences between people who reported more income than needed, enough to meet needs or less than needed. Several investigators have suggested that people with more years of education would be more adherent with medication;<sup>62,69</sup> however, education has not been found to be associated with sodium adherence. In this study, we found no differences in sodium adherence based on education. The lack of association between years of education and adherence to low-sodium diet is consistent with the propositions of the WHO adherence model,<sup>13</sup> which suggests that higher education in and of itself does not necessarily lead to higher health literacy or behavior change. In this specific study, poor health literacy was an exclusion criterion. As an observational study, there was also no intervention tested, limiting our ability to comment on behavior change.



There are probably many other barriers to following a low-sodium diet that are more important than knowledge. For instance, an identified barrier for patients is the perception that foods will taste bland and unappealing without added salt.<sup>57</sup> This barrier has been addressed by clinicians who promote flavoring with salt substitutes, using low-sodium cook books and allowing taste buds to adapt to lower-sodium foods.<sup>70</sup> Other contextual barriers identified in a qualitative descriptive study of older women with hypertension and HF include eating alone with no motivation to cook healthy meals. These women also identified that healthcare providers need to provide more education on a low-sodium diet, as well as large-print informational materials.<sup>71</sup> A study by De Brito-Ashurt (2011) identified barriers to dietary sodium restriction in people of Bangladeshi origin with chronic renal disease living the UK.<sup>72</sup> Though the disease and patient population are different from this study population, the barriers, including deeply rooted dietary beliefs, attitudes and culturally established taste for salt,<sup>72-74</sup> are likely to be universal across patient populations.

Other authors found low-sodium diet adherence rates to be about 28% or 29%.<sup>11,12</sup> The findings from this study offer a different perspective from that of population-average adherence. What we can say is that 66% of the sample fit the adherent profile to sodium recommendations and 34% fit the non-adherent profile. These numbers cannot be compared with other population averages or direct calculations of sodium adherence. This non-deterministic approach offers advantages to understanding how non-adherence can be examined and provides a list of useful predictors that are relevant to clinicians for tailoring self-care interventions and education.

## Strengths and Limitations

The primary strength of this study is the use of growth mixture modeling, which enables moving away from quantifying the overall population average to examining and quantifying heterogeneity and identifying distinct subgroups. Another strength of this research is that it includes longitudinal data collected at three time points, on a relatively large group of community dwelling stable patients with HF. Many other studies that have examined sodium adherence included only one, 24-hour urine collection in small sample sizes. With an even larger sample size, we may have been able to identify more latent categories of adherence and more predictors of trajectory membership. Further research is needed to determine if the two classes identified in this study will hold across multiple populations of patients with HF. In other words, are these predictors consistent and generalizable to different populations of patients with HF, or localized findings? The primary limitation of this dataset was that originally 23% of the total sample had missing outcome data; however, this was accounted for, at least in part, by contemporary and robust imputation techniques.

## Conclusion

Two distinct trajectories of sodium intake were identified. The non-adherent trajectory identifies a distinct pattern of dietary sodium intake that is not consistent with current sodium guidelines. Identified predictors (overweight or obese, less than 65, or having diabetes) can be used by clinicians to target interventions to patients who are most likely to struggle with eating a lower sodium diet. Results can also be used to tailor interventions and patient education programs to the highest risk groups. They can also be incorporated into HF management programs. Future studies identifying the specific barriers within these patient populations are also warranted so that the specific barriers can be targeted with interventions.

## Acknowledgments

This study was funded by grant received from the National Institutes of Health, National Heart, Lung, and Blood Institute (RO1 HL084394-01A1) and by the Philadelphia Veterans Affairs Medical Center, VISN 4 Mental Illness Research, Education, and Clinical Center.

The National Hartford Centers of Geriatric Nursing Excellence Patricia G. Archbold Scholarship program and the New Courtland Center for Transitions and Health and the Center for Integrative Science in Aging, NIH/NINR (T32-NR009356), provided predoctoral funding for Ruth Masterson Creber.


## References

1. Lennie TA, Song EK, Wu JR, et al. Three gram sodium intake is associated with longer event-free survival only in patients with advanced heart failure. *J Card Fail.* Apr; 2011 17(4):325–330. [PubMed: 21440871]
2. Hummel SL, DeFranco AC, Skorcz S, Montoye CK, Koelling TM. Recommendation of low-salt diet and short-term outcomes in heart failure with preserved systolic function. *Am J Med.* Nov; 2009 122(11):1029–1036. [PubMed: 19854331]
3. Riegel B, Moser DK, Powell M, Rector TS, Havranek EP. Nonpharmacologic care by heart failure experts. *J Card Fail.* Mar; 2006 12(2):149–153. [PubMed: 16520265]
4. Hunt SA, Abraham WT, Chin MH, et al. 2009 focused update incorporated into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the International Society for Heart and Lung Transplantation. *Circulation.* Apr 14; 2009 119(14):e391–479. [PubMed: 19324966]
5. Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure): developed in collaboration with the American College of Chest Physicians and the International Society for Heart and Lung Transplantation: endorsed by the Heart Rhythm Society. *Circulation.* Sep 20; 2005 112(12):e154–235. [PubMed: 16160202]
6. Lindenfeld J, Albert NM, Boehmer JP, et al. HFSA 2010 Comprehensive Heart Failure Practice Guideline. *J Card Fail.* Jun; 2010 16(6):e1–194. [PubMed: 20610207]
7. Hunt SA, Abraham WT, Chin MH, et al. 2009 Focused Update Incorporated Into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults. *Journal of the American College of Cardiology.* 2009; 53(15):e1–e90. [PubMed: 19358937]
8. McMurray JJV, Adamopoulos S, Anker SD, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *European Heart Journal.* 2012
9. Tsuyuki RT, McKelvie RS, Arnold JM, et al. Acute precipitants of congestive heart failure exacerbations. *Arch Intern Med.* Oct 22; 2001 161(19):2337–2342. [PubMed: 11606149]
10. Lennie TA, Worrall-Carter L, Hammash M, et al. Relationship of heart failure patients' knowledge, perceived barriers, and attitudes regarding low-sodium diet recommendations to adherence. *Prog Cardiovasc Nurs.* Winter;2008 23(1):6–11. [PubMed: 18326994]
11. Chung ML, Moser DK, Lennie TA, et al. Gender differences in adherence to the sodium-restricted diet in patients with heart failure. *J Card Fail.* Oct; 2006 12(8):628–634. [PubMed: 17045182]
12. Evangelista LS, Berg J, Dracup K. Relationship between psychosocial variables and compliance in patients with heart failure. *Heart Lung.* Jul-Aug;2001 30(4):294–301. [PubMed: 11449216]
13. World Health Organization. Adherence to long-term therapies: Evidence for action. Geneva, Switzerland: WHO; 2003.
14. Riegel B, Moelter ST, Ratcliffe SJ, et al. Excessive Daytime Sleepiness is Associated With Poor Medication Adherence in Adults With Heart Failure. *J Card Fail.* Apr; 2011 17(4):340–348. [PubMed: 21440873]

15. Masterson Creber R, Lennie T, Topaz M, Riegel B. Identifying predictors of higher than recommended sodium intake in HF patients: a mixed effect analysis of longitudinal data. *Heart and Lung: The Journal of Acute and Critical Care*. in press.
16. Gonzalez JS, Safren SA, Delahanty LM, et al. Symptoms of depression prospectively predict poorer self-care in patients with Type 2 diabetes. *Diabet Med*. Sep; 2008 25(9):1102–1107. [PubMed: 19183315]
17. Rickham PP. Human experimentation. Code of ethics of the world medical association. Declaration of Helsinki. *Br Med J*. Jul 18.1964 2(5402):177. [PubMed: 14150898]
18. Day NE, McKeown N, Wong MY, Welch A, Bingham S. Epidemiological assessment of diet: A comparison of a 7-day diary with a food frequency questionnaire using urinary markers of nitrogen, potassium and sodium. *International Journal of Epidemiology*. 2001; 30(2):309–317. [PubMed: 11369735]
19. Intersalt cooperative research group. Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 hour urinary sodium and potassium excretion. Intersalt Cooperative Research Group. *Bmj*. Jul 30; 1988 297(6644):319–328. [PubMed: 3416162]
20. Chung ML, Lennie TA, de Jong M, Wu JR, Riegel B, Moser DK. Patients differ in their ability to self-monitor adherence to a low-sodium diet versus medication. *J Card Fail*. Mar; 2008 14(2):114–120. [PubMed: 18325457]
21. Bates CJ, TD.; Bingham, SA. Bio- chemical markers of nutrient intake. Oxford, England: Oxford University Press; 1997.
22. Espeland MA, Kumanyika S, Wilson AC, et al. Statistical issues in analyzing 24-hour dietary recall and 24-hour urine collection data for sodium and potassium intakes. *Am J Epidemiol*. May 15; 2001 153(10):996–1006. [PubMed: 11384956]
23. Bingham SA, Williams R, Cole TJ, Price CP, Cummings JH. Reference values for analytes of 24-h urine collections known to be complete. *Ann Clin Biochem*. Nov; 1988 25(Pt 6):610–619. [PubMed: 3267076]
24. Day N, McKeown N, Wong M, Welch A, Bingham S. Epidemiological assessment of diet: a comparison of a 7-day diary with a food frequency questionnaire using urinary markers of nitrogen, potassium and sodium. *Int J Epidemiol*. Apr; 2001 30(2):309–317. [PubMed: 11369735]
25. Bates CJ, TD.; Bingham, SA.; Margetts, BM.; Nelson, M. Biochemical markers of nutrient intake. In: Margetts BM, NM., editor. *Design in nutritional epidemiology*. Oxford, England: Oxford University Press; 1997. p. 170-240.
26. Korhonen MH, Litmanen H, Rauramaa R, Vaisanen SB, Niskanen L, Uusitupa M. Adherence to the salt restriction diet among people with mildly elevated blood pressure. *Eur J Clin Nutr*. Nov; 1999 53(11):880–885. [PubMed: 10557001]
27. Brendgen M, Vitaro F, Bukowski WM, Doyle AB, Markiewicz D. Developmental profiles of peer social preference over the course of elementary school: associations with trajectories of externalizing and internalizing behavior. *Dev Psychol*. May; 2001 37(3):308–320. [PubMed: 11370908]
28. Croudace TJ, Jarvelin MR, Wadsworth MEJ, Jones PB. Developmental typology of trajectories to nighttime bladder control: Epidemiologic application of longitudinal latent class analysis. *American Journal of Epidemiology*. 2003; 157(9):834–842. [PubMed: 12727677]
29. Nagin DS, Tremblay RE. Parental and early childhood predictors of persistent physical aggression in boys from kindergarten to high school. *Archives of General Psychiatry*. 2001; 58(4):389–394. [PubMed: 11296100]
30. Pearson JD, Morrell CH, Landis PK, Carter HB, Brant LJ. Mixed-effects regression models for studying the natural history of prostate disease. *Statistics in Medicine*. 1994; 13(5-7):587–601. [PubMed: 7517570]
31. Dew MA, Reynolds CF 3rd, Houck PR, et al. Temporal profiles of the course of depression during treatment. Predictors of pathways toward recovery in the elderly. *Arch Gen Psychiatry*. Nov; 1997 54(11):1016–1024. [PubMed: 9366658]
32. Muthen B, Brown CH, Masyn K, et al. General growth mixture modeling for randomized preventive interventions. *Biostatistics*. Dec; 2002 3(4):459–475. [PubMed: 12933592]

33. Wiesner M, Weichold K, Silbereisen RK. Trajectories of alcohol use among adolescent boys and girls: identification, validation, and sociodemographic characteristics. *Psychol Addict Behav*. Mar; 2007 21(1):62–75. [PubMed: 17385956]
34. Warner LA, White HR, Johnson V. Alcohol initiation experiences and family history of alcoholism as predictors of problem-drinking trajectories. *J Stud Alcohol Drugs*. Jan; 2007 68(1):56–65. [PubMed: 17149518]
35. Orlando M, Tucker JS, Ellickson PL, Klein DJ. Concurrent use of alcohol and cigarettes from adolescence to young adulthood: an examination of developmental trajectories and outcomes. *Subst Use Misuse*. 2005; 40(8):1051–1069. [PubMed: 16040368]
36. Li F, Barrera M Jr, Hops H, Fisher KJ. The longitudinal influence of peers on the development of alcohol use in late adolescence: a growth mixture analysis. *J Behav Med*. Jun; 2002 25(3):293–315. [PubMed: 12055779]
37. Capaldi DM, Feingold A, Kim HK, Yoerger K, Washburn IJ. Heterogeneity in Growth and Desistance of Alcohol Use for Men in Their 20s: Prediction from Early Risk Factors and Association with Treatment. *Alcohol Clin Exp Res*. Oct 18.2012
38. Brook JS, Ning Y, Brook DW. Personality risk factors associated with trajectories of tobacco use. *Am J Addict*. Nov-Dec;2006 15(6):426–433. [PubMed: 17182444]
39. Witkiewitz K, Masyn KE. Drinking trajectories following an initial lapse. *Psychol Addict Behav*. Jun; 2008 22(2):157–167. [PubMed: 18540713]
40. Tobler AL, Komro KA. Trajectories of parental monitoring and communication and effects on drug use among urban young adolescents. *J Adolesc Health*. Jun; 2010 46(6):560–568. [PubMed: 20472213]
41. Jacob T, Koenig LB, Howell DN, Wood PK, Haber JR. Drinking trajectories from adolescence to the fifties among alcohol-dependent men. *J Stud Alcohol Drugs*. Nov; 2009 70(6):859–869. [PubMed: 19895762]
42. Feldman BJ, Masyn KE, Conger RD. New approaches to studying problem behaviors: a comparison of methods for modeling longitudinal, categorical adolescent drinking data. *Dev Psychol*. May; 2009 45(3):652–676. [PubMed: 19413423]
43. Muthen B, Muthen LK. Integrating person-centered and variable-centered analyses: growth mixture modeling with latent trajectory classes. *Alcohol Clin Exp Res*. Jun; 2000 24(6):882–891. [PubMed: 10888079]
44. Li F, Duncan TE, Hops H. Examining developmental trajectories in adolescent alcohol use using piecewise growth mixture modeling analysis. *J Stud Alcohol*. Mar; 2001 62(2):199–210. [PubMed: 11327186]
45. Riegel B, Lee CS, Glaser D, Moelter ST. Patterns of Change in Cognitive Function over Six Months in Adults with Chronic Heart Failure. *Cardiol Res Pract*. 2012; 2012:631075. [PubMed: 22928145]
46. Riegel B, Lee CS, Ratcliffe SJ, et al. Predictors of objectively measured medication nonadherence in adults with heart failure. *Circ Heart Fail*. Jul 1; 2012 5(4):430–436. [PubMed: 22647773]
47. Muthen B, Shedden K. Finite mixture modeling with mixture outcomes using the EM algorithm. *Biometrics*. Jun; 1999 55(2):463–469. [PubMed: 11318201]
48. Wang CP, Hendricks Brown C, Bandeen-Roche K. Residual Diagnostics for Growth Mixture Models. *Journal of the American Statistical Association*. 2005; 100(471):1054–1076.
49. Preacher KJ, WA.; MacCallum, RC.; Briggs, NE. *Latent growth curve modeling*. Thousand Oaks, California: Sage; 2008.
50. Jung T, Wickrama K. An Introduction to Latent Class Growth Analysis and Growth Mixture Modeling. *Social and Personality Psychology Compass*. 2008; 2(1):302–317.
51. Nylund KL AT, Muthen B. Deciding on the number of classes in latent class analysis and growth mixture modeling: A monte carlo simulation study. *Struct Equ Modeling*. 2007; 14:535–569.
52. World Health Organization. *Overview: Medication Adherence--Where Are We Today?*. Geneva, Switzerland: WHO; 2006.
53. Kenward M, Carpenter J. Multiple imputation: current perspectives. *Statistical Methods in Medical Research*. 2007; 16:199–218. [PubMed: 17621468]
54. Royston P. Multiple imputation of missing values. *The Stata Journal*. 2004; 4(3):227–241.

55. Royston P. Multiple imputation of missing values: Update of ice. *The Stata Journal*. 2005; 5(4): 527–536.
56. Stata Statistical Software: Release 11.2 [computer program]. College Station TX; StataCorp LP; 2009.
57. Bennett SJ, Lane KA, Welch J, Perkins SM, Brater DC, Murray MD. Medication and dietary compliance beliefs in heart failure. *West J Nurs Res*. Dec; 2005 27(8):977–993. discussion 994-979. [PubMed: 16275694]
58. Dyer AR, Elliott P, Shipley M, Stamler R, Stamler J. Body mass index and associations of sodium and potassium with blood pressure in INTERSALT. *Hypertension*. Jun; 1994 23(6 Pt 1):729–736. [PubMed: 8206570]
59. Gonzalez B, Lupon J, Parajon T, et al. Nurse evaluation of patients in a new multidisciplinary Heart Failure Unit in Spain. *Eur J Cardiovasc Nurs*. Apr; 2004 3(1):61–69. [PubMed: 15053889]
60. Granger BB, Ekman I, Granger CB, et al. Adherence to medication according to sex and age in the CHARM programme. *Eur J Heart Fail*. Nov; 2009 11(11):1092–1098. [PubMed: 19875409]
61. Graveley EA, Oseasohn CS. Multiple drug regimens: medication compliance among veterans 65 years and older. *Res Nurs Health*. Feb; 1991 14(1):51–58. [PubMed: 2017581]
62. Wu JR, Moser DK, Lennie TA, Burkhart PV. Medication adherence in patients who have heart failure: a review of the literature. *Nurs Clin North Am*. Mar; 2008 43(1):133–153. vii–viii. [PubMed: 18249229]
63. Evangelista L, Doering LV, Dracup K, Westlake C, Hamilton M, Fonarow GC. Compliance behaviors of elderly patients with advanced heart failure. *J Cardiovasc Nurs*. Jul-Aug;2003 18(3): 197–206. quiz 207-198. [PubMed: 12837010]
64. Riegel B, Lee C, Ratcliffe S, De Geest S, Potashnik S, Patey M, Sayers S, Goldberg L, Weintraub W. Predictors of Objectively Measured Medication Nonadherence in Adults with Heart Failure. *Circulation: Heart Failure*. 2012
65. Finch D, Davis G, Bower J, Kirchner K. Effect of insulin on renal sodium handling in hypertensive rats. *Hypertension*. May; 1990 15(5):514–518. [PubMed: 2185152]
66. Rocchini AP, Katch V, Kveselis D, et al. Insulin and renal sodium retention in obese adolescents. *Hypertension*. 1989; 14(4):367–374. [PubMed: 2676858]
67. Sechi LA. Mechanisms of insulin resistance in rat models of hypertension and their relationships with salt sensitivity. *J Hypertens*. Sep; 1999 17(9):1229–1237. [PubMed: 10489099]
68. DeFronzo RA, Cooke CR, Andres R, Faloona GR, Davis PJ. The effect of insulin on renal handling of sodium, potassium, calcium, and phosphate in man. *J Clin Invest*. Apr; 1975 55(4): 845–855. [PubMed: 1120786]
69. Chui MA, Deer M, Bennett SJ, et al. Association between adherence to diuretic therapy and health care utilization in patients with heart failure. *Pharmacotherapy*. Mar; 2003 23(3):326–332. [PubMed: 12627931]
70. Sethares KA, Elliott K. The effect of a tailored message intervention on heart failure readmission rates, quality of life, and benefit and barrier beliefs in persons with heart failure. *Heart Lung*. Jul-Aug;2004 33(4):249–260. [PubMed: 15252415]
71. Sheahan SL, Fields B. Sodium dietary restriction, knowledge, beliefs, and decision-making behavior of older females. *J Am Acad Nurse Pract*. Apr; 2008 20(4):217–224. [PubMed: 18387019]
72. de Brito-Ashurst I, Perry L, Sanders TA, Thomas JE, Yaqoob MM, Dobbie H. Barriers and facilitators of dietary sodium restriction amongst Bangladeshi chronic kidney disease patients. *J Hum Nutr Diet*. Feb; 2011 24(1):86–95. [PubMed: 21114553]
73. Little, RJ.; Rubin, DB. *Statistical analysis with missing data*. 2nd. Vol. Vol. New York: John Wiley & Sons; 2002.
74. Rubin, DB. *Multiple imputation for nonresponse in surveys*. New York, NY: John Wiley & Sons, Inc; 1987.



**Figure.**  
Trajectories of adherence to low sodium diet recommendations for patients with heart failure

**Table 1**

Characteristics of the overall sample at baseline (n=279)

	Overall n (%)		Overall n (%)
Age		Diuretic	
< 65 years	165 (59)	Yes	225 (81)
>=65 years	114 (41)	No	54 (19)
Gender		Cognitive Impairment* (n=271)	
Male	179 (64)	Yes	107 (40)
Female	100 (36)	No	164 (61)
Race		Heart failure type	
Black/Other	104 (38)	Diastolic/Mixed	86 (31)
White	175 (63)	Systolic	193 (69)
Income		Body mass index* (n=278)	
More than needed	98 (35)	Normal	74 (26)
Enough to meet needs	137 (49)	Overweight	73 (26)
Less than needed	44 (16)	Obese	131 (47)
NYHA functional class		Highest level of education	
Class I+II	65 (23)	Less than high school	27 (10)
Class III	164 (59)	High school graduate	102 (37)
Class IV	50 (18)	More than high school	150 (54)
Heart failure etiology		Diabetes	
Ischemic HF	102 (37)	Yes	171 (61)
Non-ischemic HF	177 (63)	No	108 (39)
Hypertension		COPD	
Yes	180 (65)	Yes	58 (21)
No	99 (35)	No	221 (79)

NYHA = New York Heart Association, COPD = Chronic obstructive pulmonary disease

\* Variables containing missing data

**Table 2**

Unadjusted odds of membership in the non-adherent urine sodium trajectory (n=279)

	Unadjusted Odds Ratio	95% CI	p-value
Gender			
Female		(reference)	
Male	1.51	0.57-4.03	>0.2
Body Mass Index			
Normal		(reference)	
Overweight	5.27	2.23-13.06	<0.001
Age			
>=65 years		(reference)	
< 65 years	4.80	0.45-51.80	0.20
Diabetes			
No		(reference)	
Yes	3.71	1.41-9.80	0.008
Race			
White		(reference)	
Black/Other	2.60	0.97-7.00	0.059
Income	2.07	1.07-4.02	0.031
NYHA Functional Class	1.27	0.69-2.37	>0.20
Heart failure etiology	1.57	0.37-6.68	>0.20
Heart failure type	2.68	0.89-8.03	0.078
Diuretic	3.58	0.35-36.62	>0.20
Cognitive Impairment * n=271	1.00	0.44-2.27	>0.20
LVEF * n=278	0.99	0.97- 1.00	0.089

CI = confidence intervals, NYHA = New York Heart Association, LVEF = Left Ventricular Ejection Fraction

\* Variables containing missing data



**Table 3**

Identifying predictors of non-adherence to the low-sodium dietary recommendation using multivariate logistic regression modeling (n=270)

	Odds Ratio	95% CI	p-value
Gender			
Female		(reference)	
Male	3.52	0.93-13.26	0.063
Body Mass Index			
Normal		(reference)	
Overweight	4.63	1.66-12.91	0.002**
Age			
>=65 years		(reference)	
< 65 years	4.66	1.04-20.81	0.044**
Diabetes			
No		(reference)	
Yes	4.15	1.29-13.40	0.016**
Race			
White		(reference)	
Black/Other	1.20	0.28-5.15	>0.20
Income	0.95	0.33-2.75	>0.20
NYHA Functional Class	0.71	0.43-1.17	>0.20
Heart failure etiology	1.51	0.39-5.86	>0.20
Heart failure type	1.27	0.28-5.80	>0.20
Diuretic	1.18	0.15-9.02	>0.20
Cognitive Impairment	0.65	0.21-2.04	>0.20
LVEF	0.99	0.96-1.03	>0.20

CI = confidence intervals, NYHA = New York Heart Association, LVEF = Left Ventricular Ejection Fraction

\*\* Statistically significant p<0.05