5-17-2007

Evaluating Dark Energy Probes Using Multidimensional Dark Energy Parameters

Andreas Albrecht
University of California - Davis

Gary Bernstein
University of Pennsylvania, garyb@physics.upenn.edu

Follow this and additional works at: https://repository.upenn.edu/physics_papers

Recommended Citation

Albrecht, A., & Bernstein, G. (2007). Evaluating Dark Energy Probes Using Multidimensional Dark Energy Parameters. Retrieved from https://repository.upenn.edu/physics_papers/168

Suggested Citation:

A. Albrecht and G. Bernstein. (2007). Evaluating dark energy probes using multidimensional dark energy parameters. *Physical Review D* 75, 103003.

© 2007 The American Physical Society
<http://dx.doi.org/10.1103/PhysRevD.75.103003>

This paper is posted at ScholarlyCommons. https://repository.upenn.edu/physics_papers/168
For more information, please contact repository@pobox.upenn.edu.

Evaluating Dark Energy Probes Using Multidimensional Dark Energy Parameters

Abstract

We investigate the value of future dark-energy experiments by modeling their ability to constrain the dark-energy equation of state. Similar work was recently reported by the Dark Energy Task Force (DETF) using a two dimensional parameterization of the equation-of-state evolution. We examine constraints in a nine-dimensional dark-energy parameterization, and find that the best experiments constrain significantly more than two dimensions in our 9D space. Consequently the impact of these experiments is substantially beyond that revealed in the DETF analysis, and the estimated cost per “impact” drops by about a factor of 10 as one moves to the very best experiments. The DETF conclusions about the relative value of different techniques and of the importance of combining techniques are unchanged by our analysis.

Disciplines

Physical Sciences and Mathematics | Physics

Comments

Suggested Citation:

A. Albrecht and G. Bernstein. (2007). Evaluating dark energy probes using multidimensional dark energy parameters. *Physical Review D* 75, 103003.

© 2007 The American Physical Society

<http://dx.doi.org/10.1103/PhysRevD.75.103003>

Evaluating dark energy probes using multidimensional dark energy parameters

Andreas Albrecht

Department of Physics, University of California at Davis, One Shields Avenue, Davis, California 95616, USA

Gary Bernstein

Department of Physics and Astronomy, University of Pennsylvania, 209 South 33rd Street, Philadelphia, Pennsylvania 19104, USA

(Received 14 August 2006; published 17 May 2007)

We investigate the value of future dark-energy experiments by modeling their ability to constrain the dark-energy equation of state. Similar work was recently reported by the Dark Energy Task Force (DETF) using a two dimensional parameterization of the equation-of-state evolution. We examine constraints in a nine-dimensional dark-energy parameterization, and find that the best experiments constrain significantly more than two dimensions in our 9D space. Consequently the impact of these experiments is substantially beyond that revealed in the DETF analysis, and the estimated cost per “impact” drops by about a factor of 10 as one moves to the very best experiments. The DETF conclusions about the relative value of different techniques and of the importance of combining techniques are unchanged by our analysis.

DOI: [10.1103/PhysRevD.75.103003](https://doi.org/10.1103/PhysRevD.75.103003)

PACS numbers: 95.36.+x, 98.80.Es

I. INTRODUCTION

The observed cosmic acceleration requires “dark energy” to achieve consistency with the current cosmological paradigm. The dark energy must be the dominant component of the universe today (about 70% of the energy density), yet an understanding of its fundamental nature has proved elusive. Many believe that resolving the mystery of the dark energy will force a radical change in our understand of fundamental physics. This expectation has generated great interest in the dark energy and widespread enthusiasm for an aggressive observational program to help resolve this mystery.

Recently, the Dark Energy Task Force (DETF) [1] released a report to guide the planning of future dark-energy observations. The DETF used a dark-energy “figure of merit” (FoM) based on a two-parameter description of the dark-energy evolution in order to produce quantitative findings. An interesting question is whether using the DETF FoM might lead to poor choices in shaping an observational program because of its simplicity. In particular, could the relative value of two possible experiments be distorted by the DETF FoM?

We consider this question by examining an alternative FoM. We model the dark-energy evolution with a multi-parameter model and formulate a FoM (the “D9 FoM”) which gives an experiment credit for *any* constraint it places on the dark-energy evolution. Because theory currently offers little guidance on the functional form of this evolution, we should seek from experiment any available constraint on its behavior. We use this alternative FoM to assess many of the same simulated data sets (or “data models”) considered by the DETF.

The DETF dark-energy parameters are almost completely unconstrained by current data which is typically analyzed in smaller parameters spaces in order to manifest

a noticeable impact (see for example [2]). In contrast, the DETF approach of evaluating experiments making no prior assumptions on the cosmic curvature and using a two-parameter model for the equation of state looks very ambitious (*too* ambitious to some [3]). Our work shows that the best data models constrain significantly more than two equation-of-state parameters and thus their impact was underestimated even using the ambitious DETF parameterization. As a consequence, we find that the highest quality large-scale projects also have a much lower estimated cost per FoM improvement than is achieved by the medium scale projects. Our work represents only one of several interesting advances that can reveal impact of dark-energy experiments that is greater than seen by the DETF (see, for example, Refs. [4,5]). In order to give a more focused discussion our new FoM is the only significant technical difference between our methods and those of the DETF.

II. DARK-ENERGY PARAMETERS

Following the DETF (and many others) we model the dark energy as a homogeneous and isotropic fluid. The complete dark-energy history can then be given by the dark-energy density today and the “equation-of-state parameter” (the ratio of the density and pressure of the fluid) $w(a)$ as a function of time or cosmic scale factor a .

This paper is about the choice of parametrization of $w(a)$. The DETF used a standard linear form

$$w(a) = w_0 + w_a(1 - a). \quad (1)$$

For this work we used a piecewise constant model of $w(a)$

$$w(a) = -1 + \sum_{i=1}^{N_G} w_i T(a_i, a_{i+1}), \quad (2)$$

where the “top-hat function” $T(a_i, a_{i+1})$ is unity for $a_i > a \geq a_{i+1}$ and zero otherwise. Any nonzero value of a w_i gives a deviation from a cosmological constant ($w = -1$). The DETF linear model is well approximated by a subspace of our larger space. Here we use $N_G = 9$.

The DETF considered the degree to which w_0 and w_a would be constrained by a variety of data models. The DETF FoM (first used in Ref. [6]) is given by the reciprocal area in $w_0 - w_a$ space enclosed by the 95% confidence contour for a given data model. Correspondingly our 9D FoM is the reciprocal hypervolume enclosed by the 95% contour in our larger space.

III. METHODS, $w(a)$ EIGENMODES

Like the DETF we use the Fisher matrix formalism and assume a Gaussian probability distribution to evaluate the 9D FoM. A general (unnormalized) Gaussian probability distribution for parameters $\vec{x}$ around a central value $\vec{x}_0$ in N dimensions is given by

$$\exp\{(-\Delta\vec{x}\mathbf{F}\Delta\vec{x})/2\}, \quad (3)$$

where $\Delta\vec{x} = \vec{x} - \vec{x}_0$. The N eigenvectors $\vec{f}_i$ of $\mathbf{F}$ give the directions of the principle axes (or “principle components”) of the error ellipsoid, and the width of the error ellipsoid along axis $\vec{f}_i$ is given by σ_i , the inverse square root of the i -th eigenvalue of $\mathbf{F}$. The $\vec{f}_i$ describe the independently measured “modes” of $w(a)$ in the 9D space.

We only calculate ratios of FoM’s so one can define the FoM as $\prod_i \sigma_i^{-1}$ (additional constant factors will drop out). Depending on whether $\mathbf{F}$ is defined in the DETF or 9D space of w parameters, this formula gives the DETF or the 9D FoM. We use many DETF data models, but we exclude models of galaxy-cluster data, since the extension of the DETF calculations to our scheme is not straightforward, especially as regards systematic error estimates.

Each data model corresponds to its own Fisher matrix $\mathbf{F}$ from which the $\vec{f}_i$ ’s, σ_i ’s and the FoM can be calculated. Figure 1 gives the errors σ_i and modes $\vec{f}_i$ corresponding to the DETF “Stage 2” data minus clusters. Stage 2 is the DETF projection of the data upon completion of existing projects. The DETF defines major longer term projects as “Stage 4” and smaller faster future programs as “Stage 3.” Fig. 2 gives the same information for a particular Stage 4 data model. None of the DETF data models are powerful enough to constrain all nine directions in our parameter space. This is reflected in large values of σ_i for larger i . We do not trust our formalism to give meaningful answers in directions that place bounds weaker than unity on the w_i ’s, and such weak constraints are not likely to enlighten us about dark energy. We handle this issue by resetting all $\sigma_i > 1$ to unity when we calculate the FoM. That way changes in σ_i in directions where the method is not trusted do not contribute to changes in the FoM.

FIG. 1 (color online). Projected impact of experiments currently underway (DETF Stage 2) in the 9D space of w parameters: The upper panel gives the errors σ_i in increasing order. The lower panels give the corresponding independently measured modes of $w(a)$ (the $\vec{f}_i$). The nine markers on each curve give the nine components of each $\vec{f}_i$ positioned at $\bar{a}_j \equiv \sqrt{a_j a_{j+1}}$. The dotted vertical lines show the values of a_j .

The 9D FoM’s reported here use grids $\{a_i\}$ with minimum redshift $z_{\min} \equiv 1/(a_{\min} - 1) = 0.01$, $z_{\max} = 4$ with the grid uniform in a . We considered a variety of different grids (varying the spacing pattern, redshift range and total grid points N_G) [7]. The final grid was chosen to give the highest FoM’s which means we use the parameters which are best measured by the model data sets. Increasing N_G

FIG. 2 (color online). The impact of the DETF “Stage 4 space weak lensing optimistic” data model: This is a much higher quality data set vs Stage 2. More modes are well measured and the well measured modes reach to higher redshift (lower a). This plot has the same format as Fig. 1.

beyond the point where the well measured modes are well resolved does not significantly change the FoM's or the shapes and σ_i 's of the well measured modes. We find this convergence under increasing N_G an attractive feature of our parameterization and we found $N_G = 9$ large enough to achieve convergence.

Independently measured modes of $w(a)$ (the $\vec{f}_i$'s) reveal interesting details about each data model. The fact that the best measured Stage 2 $\vec{f}_i$'s (shown in Fig. 1) approach zero for $a < 0.5$ (larger redshifts) but the corresponding Stage 4 modes (in Fig. 2) do not illustrates the deeper redshift reach of the Stage 4 data. Other specific differences among data models can be understood by inspecting the plots in [7].

Our form for $w(a)$ has been used previously by others applied to different data models [6,8–10] (see also Ref. [11]). To the extent that comparison is possible our work is consistent with these earlier papers. In particular, the claims in Ref. [10] that there are only two measurable w parameters stems from a different formal definition of what it means to measure a parameter usefully. The choice we make here is best suited to our purpose, which is to make a direct comparison with the DETF methods.

There are some small technical differences between our calculations and those in the DETF report. When we use

two supernova data models in combination they share the same nuisance parameters (except for the photometric redshifts). The DETF calculations keep the nuisance parameters separate. The PLANCK prior is the same one used by the DETF, but the Fisher matrix is expressed in the variables $\{n_s, \delta_\zeta, \omega_m, \omega_b, \theta_s\}$ where we use $\ln(\theta_s) = -0.252\ln(\omega_m) - 0.83\ln(\omega_b) - \ln(D_A^{co}(a^*)) + 8.2094$ (from Ref. [12]). These parameters are defined in the DETF report. Some other modest differences stem from the shortcomings of the transfer function formalism discussed in section 9.2 of the DETF Appendix. These small differences from the DETF calculations are included in all the FoM's presented here (DETF and 9D) so our comparisons of FoM's will reflect the different w parameter choices only.

IV. RESULTS AND INTERPRETATION

The DETF present their main results in 4 bar charts showing the FoM ranges for particular data models. The values are given as ratios to the Stage 2 FoM so that progress beyond Stage 2 can be read directly, with increasing progress corresponding to larger values along the y axis. The four panels in Fig. 3 show equivalent plots (using the same DETF data models minus clusters). The dark bars

FIG. 3 (color online). Figure of merit improvements vs Stage 2 for DETF data models: Dark bars show the DETF FoM. To the right of each dark bar is a light bar that reflects the 9D FoM for the identical data model. The 9D FoM registers a much greater impact from each data model, and also shows a much greater improvement at Stage 4 vs Stage 3. Each panel corresponds to one of the four main bar charts in the DETF report (B = “baryon oscillations,” S = “supernovae,” W = “weak lensing,” A = “All” see pp. 16–20 of Ref. [1]). The x-axis labels are similar to those on the DETF plots, but abbreviated.

show the DETF FoM and the light bars show the 9D FoM. The 9D FoM shows much larger values for all the strong data models.

One can see that good combinations of Stage 3 data give 9D FoM improvements of an order of magnitude and strong Stage 4 data combinations give 9D FoM improvements of 3 orders of magnitude or more. Compared with the DETF results (about half an order of magnitude to Stage 3 and 1 order of magnitude to Stage 4) this is not only a strong showing overall, but specifically the 9D FoM exhibits a greater improvement factor going from Stage 3 to Stage 4 as compared with the DETF FoM. Given the ballpark costs quoted (but not independently verified) by the DETF of a few $\times 10^7$ for Stage 3 and $0.3 - 1 \times 10^9$ for Stage 4, our work indicates that good Stage 4 projects tend to be much more cost effective (about 10 times better in \$ per FoM increment) than Stage 3 projects. The opposite seems to be the case when using the DETF FoM, but our work shows that this is only because the 2D DETF parametrization prevents the better experiments from showing their full capabilities.

The DETF FoM is constructed in a 2-D parameter space. To build intuition, consider the following effective “reduction to 2D” of the 9D FoM:

$$\mathcal{F}_2 \equiv \mathcal{F}_9^{2/D_e}. \quad (4)$$

Here $\mathcal{F}_2$ and $\mathcal{F}_9$ are the reduced and regular 9D FoM's, respectively. One can think of $\mathcal{F}_2$ as the product of only two $1/\sigma_{\text{eff}}$'s where σ_{eff} is a suitably defined geometric mean of the σ_i 's. The “effective dimension” D_e can be thought of as the number of directions constrained by the data model in 9D parameter space.

Figure 4 is the same as Fig. 3 except Eq. (4) has been applied to all the 9D FoM's. Values of D_e were assigned to the data models (details in the caption) to create an approximate match (by eye) between rescaled 9D and DETF FoM's. Inspection of Fig. 4 suggests that the rescaling accounts for many of the differences between the FoM's. A refined assignment of D_e 's gradually increasing with improving data models would give an even better account of the differences (but still would not match up every detail). Successful matching of the DETF and 9D FoM's after rescaling implies that D_e directions in the 9D space are measured as well as the two DETF parameters. This applies only in an average sense: The best 9D modes are measured much better than the best DETF parameter, others are less well measured. We present this rescaling to give intuition about the similarities and differences between the DETF and 9D FoM's. The similarity of the DETF and 9D bars in Fig. 4 help one see how the detailed DETF conclusions about the relative value of different techniques and combinations of techniques are unchanged

FIG. 4 (color online). Rescaled FoM's: This is identical to Fig. 3 except all the 9D FoM's have been scaled according to Eq. (4). We use $D_e = 2.5$ for Stage 2, $D_e = 3$ for Stage 3, $D_e = 4$ for Stage 4 pessimistic and $D_e = 4.5$ for Stage 4 optimistic. These plots suggest that the scaling gives a reasonable account of the impact of measuring more parameters in 9D vs 2D spaces demonstrated in Fig. 3.

FIG. 5 (color online). Impact of adding additional constraints on curvature (left side) and the Hubble constant (right side): Dark bars show the impact on the DETF FoM and light bars show the impact on the 9D FoM. The y axis is the factor by which the FoM's change (for Stage 4 space, without Stage 2) when the additional constraints are imposed. Changing from the DETF to the 9D parameters does not undermine the insensitivity to additional constraints reported by the DETF. This plot corresponds to the plot on p. 14 of Ref. [1] (in a different format).

by our analysis. However, we do not expect Eq. (4) to offer a universal relationship between the two FoM's that extrapolates to all possible data models.

Imposing additional constraints or “priors” on specific parameters generally will improve the figures of merit. The DETF report has a plot similar to Fig. 5 (dark bars) which illustrates the “impact factor” (factor by which the FoM improves) from imposing stronger priors on the curvature and the Hubble constant. The DETF found the impact to be modest, and noted that the best data models actually determine the parameters so well themselves that the impact of additional priors was small.

Our work shows that DETF data models can constrain more $w(a)$ parameters than the two considered by the DETF. Does this improvement come at the expense of poorer constraints on other parameters, due to the greater flexibility of the dark-energy model? The lighter bars in Fig. 5 show the impact factor on the 9D FoM. In no case is the impact substantially greater for the 9D case, indicating that going to the 9D $w(a)$ model does not significantly undermine the constraints on the curvature and Hubble constant from the modeled data. In fact, the 9D impact factors are much smaller with the Hubble prior than for the DETF FoM, suggesting that the directions in the 9D space constrained by these data models have even less degeneracy with the Hubble parameter than w_0 and w_a .

V. CONCLUSIONS

We have analyzed a FoM for dark-energy probes which is defined in a nine-dimensional parameter space, up from the (already ambitious) two-parameter space used by the DETF. Our 9D FoM gives a more complete account of the impact of a given data model. We find the DETF data models constrain significantly more parameters than the

two used by the DETF, leading to vastly improved FoM's in the 9D space. Our rescaling law gives an intuitive account of the impact of measuring more parameters.

The 9D FoM follows the same logic as the DETF FoM, namely, it attaches equal weight to any constraint on $w(a)$. The measurement of a nonzero value for any of the independently measured combinations $\vec{f}_i$ would be equally significant in that it would rule out a pure cosmological constant. Thus the higher values of the 9D FoM (vs the DETF) reflect genuinely greater discovery power.

The effective number of parameters measured (D_e) increases with the quality of the project, and consequently so does the gap in the impact registered by the 9D vs DETF FoM's. Good Stage 4 projects measure more than four parameters and produce a three or four order of magnitude improvement in the 9D FoM vs a 1 order of magnitude improvement in the DETF FoM. As a result, our 9D work indicates that good Stage 4 projects achieve a 10 times lower estimated cost per FoM increase than Stage 3 projects, the reverse of the DETF conclusion. Aside from this major difference, the detailed DETF conclusions about the relative value of different techniques and the importance of combining techniques are unchanged by the 9D analysis.

ACKNOWLEDGMENTS

We thank the DETF for a great collaboration on which this work was based, and Lloyd Knox for his simulated PLANCK Fisher matrices and helpful discussions. This work was supported in part by DOE Grants No. DE-FG03-91ER40674 (A. A.) and No. DE-FG02-95ER40893 (G. B.) and by NASA Grant No. BEFS-04-0014-0018 (G. B.).

- [1] A. Albrecht *et al.*, astro-ph/0609591.
- [2] D. N. Spergel *et al.*, astro-ph/0603449.
- [3] E. V. Linder, *Astropart. Phys.* **24**, 391 (2005).
- [4] H. Zhan, *J. Cosmol. Astropart. Phys.* 08 (2006) 008.
- [5] M. Schneider, L. Knox, H. Zhan, and A. Connolly, *Astrophys. J.* **651**, 14 (2006).
- [6] D. Huterer and M. S. Turner, *Phys. Rev. D* **64**, 123527 (2001).
- [7] See EPAPS Document No. E-PRVDAQ-75-053710 for full details about this exploration (including $\hat{f}_i$ plots for all data models); this information can be found at www.physics.ucdavis.edu/Cosmology/albrecht/MI0608269 or at EPAPS. For more information on EPAPS, see <http://www.aip.org/pubservs/epaps.html>.
- [8] D. Huterer and G. Starkman, *Phys. Rev. Lett.* **90**, 031301 (2003).
- [9] L. Knox, A. Albrecht, and Y. S. Song, astro-ph/0408141.
- [10] E. V. Linder and D. Huterer, *Phys. Rev. D* **72**, 043509 (2005).
- [11] A. G. Riess *et al.*, astro-ph/0611572.
- [12] W. Hu, *ASP Conf. Ser.* **399**, 215 (2005).