



11-4-2013

Effects of Self-Care on Quality of Life in Adults with Heart Failure and Their Spousal Caregivers: Testing Dyadic Dynamics Using the Actor-Partner Interdependence Model

Ercole Vellone

Misook L. Chung

Antonello Cocchieri

Gennaro Rocco

Rosaria Alvaro

See next page for additional authors

Follow this and additional works at: <https://repository.upenn.edu/nrs>

 Part of the [Behavioral Medicine Commons](#), [Cardiology Commons](#), [Cardiovascular Diseases Commons](#), [Circulatory and Respiratory Physiology Commons](#), [Medical Humanities Commons](#), [Nursing Commons](#), and the [Preventive Medicine Commons](#)

Recommended Citation

Vellone, E., Chung, M. L., Cocchieri, A., Rocco, G., Alvaro, R., & Riegel, B. (2013). Effects of Self-Care on Quality of Life in Adults with Heart Failure and Their Spousal Caregivers: Testing Dyadic Dynamics Using the Actor-Partner Interdependence Model. *Journal of Family Nursing*, 20 (1), 120-141. <http://dx.doi.org/10.1177/1074840713510205>

This paper is posted at ScholarlyCommons. <https://repository.upenn.edu/nrs/180>
For more information, please contact repository@pobox.upenn.edu.

Effects of Self-Care on Quality of Life in Adults with Heart Failure and Their Spousal Caregivers: Testing Dyadic Dynamics Using the Actor-Partner Interdependence Model

Abstract

Emotions are contagious in couples. The purpose of this study was to analyze the manner in which adults with chronic heart failure (HF) and their informal caregivers influence each other's self-care behavior and quality of life (QOL). A sample of 138 HF patients and spouses was enrolled from ambulatory centers across Italy. The Actor-Partner Interdependence Model (APIM) was used to analyze dyadic data obtained with the Self-Care of Heart Failure Index (SCHFI), the Caregivers Contribution to the SCHFI, and the Short Form 12. Both actor and partner effects were found. Higher self-care was related to lower physical QOL in patients and caregivers. Higher self-care maintenance in patients was associated with better mental QOL in caregivers. In caregivers, confidence in the ability to support patients in self-care was associated with improved caregivers' mental QOL, but worsened physical QOL in patients. Interventions that build the caregivers' confidence are needed.

Keywords

family caregivers, heart failure, dyads, couples, Actor-Partner Interdependence Model, self-care, quality of life

Disciplines

Behavioral Medicine | Cardiology | Cardiovascular Diseases | Circulatory and Respiratory Physiology | Medical Humanities | Medicine and Health Sciences | Nursing | Preventive Medicine

Author(s)

Ercole Vellone, Misook L. Chung, Antonello Cocchieri, Gennaro Rocco, Rosaria Alvaro, and Barbara Riegel

Effects of Self-Care on Quality of Life in Adults with Heart Failure and Their Spousal Caregivers:**Testing Dyadic Dynamics using the Actor-Partner Interdependence Model**

Ercole Vellone, MSN, RN
Research fellow, School of Nursing
Tor Vergata University, Via Montpellier, 1
00133 Rome, Italy
Phone: +390672596871
Fax: +390672596961
e-mail: ercole.vellone@uniroma2.it

Misook L. Chung, PhD, RN
Associate Professor
Co-Director, RICH Heart Program
College of Nursing, University of Kentucky
529 CON Building, 760 Rose Street
Lexington, KY 40536-0232
Tel 859-323-8024
Fax 859-323-1057
misook.chung@uky.edu

Antonello Cocchieri, MSN, RN
PhD student, School of Nursing
Tor Vergata University, Via Montpellier, 1
00133 Rome, Italy
Phone: +390672596871
Fax: +390672596961
e-mail: antonello.cocchieri@uniroma2.it

Gennaro Rocco, MSN, RN
Director, Center of Excellence for Nursing Scholarship,
Viale Giulio Cesare, 78
00192 Rome, Italy
Phone +39 0637511597; Fax +39 0645437034
e-mail: genna.rocco@gmail.com

Rosaria Alvaro, MSN, RN
Associate Professor, School of Nursing
Tor Vergata University, Via Montpellier, 1
00133 Rome, Italy
Phone: +390672596871; Fax: +390672596961
e-mail: rosaria.alvaro@uniroma2.it

Barbara Riegel, RN, DNSc, FAHA, FAAN
Professor and Edith Clemmer Steinbright Chair of Gerontology

School of Nursing, University of Pennsylvania
 418 Curie Boulevard
 Philadelphia, PA USA 19104-4217
 215-898-9927 (W)
 240-282-7707 (eFAX)
briegel@nursing.upenn.edu

Keywords: caregivers, heart failure, dyads, couples, actor-partner interdependence model, self-care, quality of life

Acknowledgements: The authors grateful acknowledge Patricia D'Antonio, PhD, RN, FAAN and George J. Knaf, PhD for their review of a prior version of this manuscript.

Declaration of Conflicting Interests: None

Funding: This study was funded by the Center of Excellence for Nursing Scholarship, Rome, Italy

Barbara Riegel, DNSc, RN is the Edith Clemmer Steinbright Professor of Gerontology at the University of Pennsylvania, School of Nursing. She is known internationally as an expert in heart failure self-care.

Publications:

1. Riegel, B., Moser, D. K., Anker, S. D., Appel, L. J., Dunbar, S. B., Grady, K. L., . . . Whellan, D. J. (2009). State of the science: Promoting self-care in persons with heart failure: A scientific statement from the American Heart Association. *Circulation, 120*(12), 1141-1163. doi:10.1161/CIRCULATIONAHA.109.192628
2. Riegel, B., Lee, C. S., & Dickson, V. V. (2011). Self care in patients with chronic heart failure. *Nature Reviews. Cardiology, 8*(11), 644-654. doi:10.1038/nrcardio.2011.95

Misook Lee Chung, PhD, RN, is Associate Professor, at the University of Kentucky, College of Nursing. She studies family caregivers and patients with heart failure. Her particular interest is in understanding the dynamic interdependent relationships between patients and family caregivers in terms of the behavioral, physiological, and psychological outcomes of both dyad members and developing effective patient-caregiver dyad-based interventions to improve outcomes. Publications:

1. Chung, M. L., Lennie, T. A., Dekker, R. L., Wu, J. R., & Moser, D. K. (2011). Depressive symptoms and poor social support have a synergistic effect on event-free survival in patients with heart failure. *Heart and Lung, 40*(6), 492-501. doi:10.1016/j.hrtlng.2010.08.001
2. Chung, M. L., Moser, D. K., Lennie, T. A., & Frazier, S. K. (2012). Perceived social support predicted quality of life in patients with heart failure, but the effect is mediated by depressive symptoms. *Quality of Life Research*. Advanced online publication. doi:10.1007/s11136-012-0294-4
3. Chung, M. L., Moser, D. K., Lennie, T. A., & Rayens, M. K. (2009). The effects of depressive symptoms and anxiety on quality of life in patients with heart failure and their spouses: Testing dyadic dynamics using Actor-Partner Interdependence Model. *Journal of Psychosomatic Research, 67*(1), 29-35. doi:10.1016/j.jpsychores.2009.01.009

Antonello Cocchieri, RN, MSN is a PhD student at Tor Vergata University in Rome, Italy. His interest is in heart failure self-care. He is currently conducting a survival analysis of the effects of heart failure self-care on hospitalization and mortality. Publications:

1. Vellone, E., Fida, R., Cocchieri, A., Sili, A., Piras, G., & Alvaro, R. (2001). Positive and negative impact of caregiving to older adults: a structural equation model. *Professioni Infermieristiche* 64(4), 237-248.
2. Vellone, E., Riegel, B., Cocchieri, A., Barbaranelli, C., D'Agostino, F., Glaser, D., . . . Alvaro, R. (2012). Validity and reliability of the caregiver contribution to Self-care of Heart Failure Index. *Journal of Cardiovascular Nursing*. Advanced online publication. doi:10.1097/JCN.0b013e318256385e.

Gennaro Rocco, RN, MSN is the Director of the Center of Excellence for Nursing Scholarship in Rome, Italy. His main interest is ethics and in establishing an infrastructure in Italy for nursing research.

Publications:

1. Stievano, A., De Marinis M.G., Russo, M.T., Rocco, G. & Alvaro, R. (2012). Professional dignity in nursing in clinical and community workplaces. *Nursing Ethics* 19(3), 341-356. doi:10.1177/0969733011414966
2. Stievano, A., Bonfigli, A., Fanfera, E., Finocchi, G., Montevicchi, A., Nappini, P., Tallarita, F., Turci, C. & Rocco, G. (2011). ILISI® digital index of the Italian scientific literature of nursing. *Nurse Education Today*, 31(3), 294-298. doi:10.1016/j.nedt.2010.11.019

Ercole Vellone, RN, MSN is a research fellow at Tor Vergata University in Rome, Italy. His primary interest is in heart failure self-care and caregiver contributions to patients' self-care. Publication:

1. Vellone, E., Riegel, B., Cocchieri, A., Barbaranelli, C., D'Agostino, F., Glaser, D., . . . Alvaro, R. (2012). Validity and reliability of the caregiver contribution to Self-care of Heart Failure Index. *Journal of Cardiovascular Nursing*. Advanced online publication. doi:10.1097/JCN.0b013e318256385e.
2. Vellone, E., Riegel, B., D'Agostino, F., Fida, R., Rocco, G., Cocchieri, A., & Alvaro, R. (in press) Structural equation model testing the situation-specific theory of heart failure self-care. *Journal of Advanced Nursing*.

Rosaria Alvaro, RN, MSN is Associate Professor in nursing science at Tor Vergata University. Her main interest is in quality of life in people with cardiovascular disease. Publications:

1. Vellone, E., Savini, S., Simeone, S., Barbato, N., Carovillano, G., Caramia, M.D. & Alvaro, R. (2011). [Reliability of caregivers in assessing the quality of life of stroke survivors: an explorative study]. *Assistenza Infermieristica e Ricerca*, 30(4), 180-188.
2. Vellone, E., Savini, S., Barbato, N., Carovillano, G., Caramia, M. & Alvaro, R. (2010). Quality of life in stroke survivors: first results from the reliability and validity of the Italian version of the Stroke Impact Scale 3.0. *Annali d'Igiene*, 22(5), 469-479.

Vellone E, Chung M, Cocchieri A, Gennaro R, Alvaro R, Riegel B, Effects of Self-Care on Quality of Life in Adults with Heart Failure and Their Spousal Caregivers: Testing Dyadic Dynamics using the Actor-Partner Interdependence Model, *Journal of Family Nursing*, 2014,

20(1), 120-41. doi: 10.1177/1074840713510205*

Abstract

Emotions are contagious in couples. The purpose of this study was to analyze the manner in which adults with chronic heart failure (HF) and their informal caregivers influence each other's self-care behavior and quality of life (QOL). A sample of 138 HF patients and spouses was enrolled from ambulatory centers across Italy. The actor-partner interdependence model (APIM) was used to analyze dyadic data obtained with the Self-Care of Heart Failure Index, the Caregivers Contribution to the Self-Care of Heart Failure Index, and the Short Form 12. Both actor and partner effects were found. Higher self-care was related to lower physical QOL in patients and caregivers. Higher self-care maintenance in patients was associated with better mental QOL in caregivers. In caregivers, confidence in the ability to support patients in self-care was associated with improved caregivers' mental QOL and improved patients' physical QOL. Interventions the build caregiver confidence are needed.

Funding: This study was funded by the Center of Excellence for Nursing Scholarship, Rome, Italy

Acknowledgements: The authors grateful acknowledge Patricia D'Antonio, PhD, RN, FAAN and George J. Knafl, PhD for their review of a prior version of this manuscript.

Introduction

A growing body of literature addresses the unique bond that forms between individuals in a committed personal relationship (M. A. Lewis et al., 2006). Observing the emotional expressions of another person causes an automatic tendency to share the person's experiences (Frith & Frith, 2006) and emotions are particularly contagious in couples (Ridker, 2009; Rohrbaugh, Shoham, Cleary, Berman, & Ewy, 2009; Satpathy et al., 2008). Yet, the study of emotional contagion within couples is in its infancy in couples in which one individual is the caregiver for a spouse with a chronic illness. Therefore, the purpose of this study was to analyze the manner in which adult patients with chronic heart failure (HF) and their informal spousal caregivers influence each other's self-care behavior and quality of life (QOL).

Background

Heart failure affects more than 5 million individuals in the United States (US) and many more million worldwide, especially in countries with large numbers of older adults (Go et al., 2012). In 2002, it was reported that HF had an overall population prevalence of approximately 1-3%, rising to approximately 10% in the very elderly (McMurray & Stewart, 2002). In the US, currently about 2.8% of the adult population has HF (Go, et al., 2012). In Italy, where these data were collected, 1.1% of the overall population has HF with a prevalence that increases to 12.1% in people aged 70 and over (Valle et al., 2006). Self-care is an integral element of the treatment of HF that is necessary at all stages of HF (Ueland et al., 2005).

The trajectory of HF is unpredictable, with a gradual decline in functional capacity that is accompanied by significant symptom burden, impaired mental acuity, and poor QOL (Bennett & Sauve, 2003; Gott et al., 2006; Johnson, 1997). This decline in functioning, accompanied by an increase in caregiving responsibilities, significantly burdens informal caregivers (Calvert,

Freemantle, & Cleland, 2005; Friedman, 1997; Goldstein & Lynn, 2006; D. T. Lee, Yu, Woo, & Thompson, 2005; E. F. Lewis et al., 2001; Pressler et al., 2012). In couples, the burden of caregiving has been shown to impair the couple's relationship by causing difficulties in communication and social roles that accentuate the emotional distress experienced by both patients and caregivers (Luttik, Blaauwbroek, Dijker, & Jaarsma, 2007; Molloy, Johnston, & Witham, 2005).

Although the interdependence between individuals in a committed relationship has been discussed in the health care literature for decades (Geden & Taylor, 1999; Vellone, Piras, Venturini, Alvaro, & Cohen, 2012), relatively few investigators have studied HF patient and caregiver dyads. Seven such studies were located (Agren, Evangelista, Davidson, & Stromberg, 2010; Agren, Evangelista, Hjelm, & Stromberg, 2012; Allison & Waterman, 2002; Kurylo, Elliott, DeVivo, & Dreer, 2004; Retrum, Nowels, & Bekelman, 2012; Ridker, 2009; Sebern & Riegel, 2009) but only one of these used an analytic approach that allowed assessment of the effect of one individual on the other person in the dyad (Ridker, 2009). Three of the studies compared outcomes in separate patient and partner groups (Agren, et al., 2010; Agren, et al., 2012; Allison & Waterman, 2002), an approach referred to as family-related rather than family research (Feetham, 1991). Four of the studies examined the influence of one person's emotions on behavior of the partner but none of these studies fully accounted for the dynamics of the dyad. For example, in one study of 27 patients with HF and their family caregivers, a negative orientation toward solving problems in caregivers was a significant predictor of higher depression and decreased QOL in patients (Kurylo, et al., 2004). In another study exploring the contribution of supportive relationships to HF self-care in 75 dyads, strong, positive relationships were found between patient and spousal caregiver shared care decision making and HF patients'

self-care maintenance and self-care confidence (Sebern & Riegel, 2009). Another study examined congruence or consistency in perspective in 17 patient and caregiver dyads using a qualitative approach (Retrum, et al., 2012). These authors found that much of the tension and distress among couples was due to conflicting views about how emotions should be expressed.

Only one prior study of HF and caregiver dyads used the actor-partner interdependence model (APIM), an analytic approach that fully accounts for dyadic dynamics (Ridker, 2009). Chung and colleagues enrolled 58 HF patient and spousal caregiver dyads from a single site in the Southeastern US. Depressive symptoms and anxiety were assessed using the Brief Symptom Inventory. Quality of life was assessed using the Minnesota Living with Heart Failure Questionnaire. Using a cross-sectional design, the investigators found both actor and partner effects. Depression and anxiety had significant actor effects on both patients and spouses; those with more depression and higher anxiety had poorer QOL. Partner effects were evident on patients' QOL; patients whose spouses had more depressive symptoms and higher anxiety were more likely to have poorer QOL (Ridker, 2009).

No other studies of dyadic dynamics were located in the existing HF literature. Because so little is known about the manner in which HF patients and their informal caregivers influence each other and because self-care and QOL are such fundamental issues in HF, we explored the manner in which self-care contributes to QOL in patients and spousal caregivers.

Methods

The research question that stimulated this study was: How does chronic illness affect the manner in which individuals in a couple influence each other? The specific aim of this study was to describe the manner in which QOL and self-care vary within a couple as a function of chronic HF. We hypothesized that, in dyads of HF patients and spousal caregivers, the QOL and self-

care of one member of the dyad would influence the QOL and self-care of the other member of the dyad. All analyses were guided by the model shown in Figure 1.

Design and Sample

A cross-sectional design was used to conduct the study. A convenience sample of 1192 HF patients was enrolled from ambulatory cardiovascular centers in 19 Italian provinces located in the north, center and south of Italy. Inclusion criteria specified enrollment of only adults (older than 18 years) with a confirmed diagnosis of HF regardless of etiology or type. Heart failure was confirmed by echocardiography and clinical evidence of HF. Patient participants had to be stable, without an acute coronary event in the prior three months. Otherwise eligible patients were excluded if they had obvious evidence of dementia. No other inclusion and exclusion criteria were specified so that we could include as many patients as possible. This analysis was conducted using data from enrolled patient participants who were accompanied by a spousal caregiver at the time of enrollment. The rationale for focusing on spouses was that generally spousal caregivers are considered the primary caregivers and their relationship with patients is more interdependent than other family members or caregivers (M. A. Lewis, et al., 2006). Any spouse accompanying an enrolling HF patient was invited to enroll; no further inclusion or exclusion criteria were specified for the spousal caregivers.

Procedures

The Institutional Review Board of each center where patients and caregivers were recruited approved the study. Data collection was done by nurses who were trained in the standardized collection of research data. Data collectors screened patients during routine ambulatory visits. When patients met the inclusion criteria, the study was explained and patients were invited to participate. If an informal primary caregiver was attending the patient's visit, the

research assistant invited him/her to participate as well. Both patients and caregivers gave written informed consent before data collection was done. All data were obtained in a private, quiet room. Anecdotally, most caregivers completed the instruments on their own, while most patients asked research assistants to read the instrument items to them.

Instruments.

Both patients and caregivers completed a socio-demographic survey, a measure of self-care, and a measure of QOL, as detailed below. An investigator-designed survey was used to collect information regarding the dyad's socio-demographic characteristics (e.g. age, gender, education, relationship between patient and caregiver). Information on the patient's clinical status (e.g. illness duration, New York Heart Association [NYHA] functional class, ejection fraction) was abstracted by nurses from the medical record.

The Self-Care of Heart Failure Index version 6.2 (SCHFI v.6.2) (Riegel et al., 2004; Riegel, Lee, Dickson, & Carlson, 2009) was used to measure self-care in HF patients. The SCHFI is a widely used instrument measuring self-care behaviors in people with HF that consists of three scales: the self-care maintenance, the self-care management, and the self-care confidence scales. The self-care maintenance scale has 10 items measuring self-monitoring (e.g. checking weight daily) and treatment adherence (e.g. following a low salt diet, taking medications). The self-care management scale has 6 items that measure how quickly patients recognize and evaluate symptoms of a HF exacerbation (shortness of breath or ankle swelling), treatment implemented after symptoms are identified (e.g. reducing salt and fluid intake) and treatment evaluation. The self-care confidence scale, with 6 items, evaluates how much confidence patients perceive in each phase of the self-care process (e.g. confidence in symptom management). All SCHFI v.6.2 items use a 4-point self-report scale. Each SCHFI v.6.2 scale yields a standardized

score from 0 to 100 where higher scores mean better self-care. The SCHFI v.6.2 has adequate psychometric characteristic in the US population (Riegel, et al., 2009) and has been found to have adequate validity and reliability in the Italian population as well (Vellone et al., 2013).

The Caregiver Contribution to Self-care of Heart Failure Index [CC-SCHFI] (Vellone et al., 2012) was used to measure self-care contributions by caregivers. The CC-SCHFI is a modified version of the SCHFI v.6.2 for use with caregivers of HF patients that measures the extent to which a caregiver contributes to a patient's self-care maintenance and self-care management as well as the caregiver's confidence in his/her ability to contribute to the patient's self-care. Items in the CC-SCHFI correspond perfectly to those of the SCHFI v.6.2. The caregiver contribution to self-care maintenance scale, with 10 items, measures how often a caregiver recommends to the patient that she/he monitor HF symptoms (e.g. checking ankles for swelling) and adhere to HF treatment (e.g. reminding the patient not to forget his/her medicine). The scale also assesses how often the caregiver substitutes for the patient because she/he is not able to do specific behaviors (e.g. checking the patient's ankles or administering the medicines). The caregiver contribution to self-care management scale has 6 items that measure the caregiver's ability to recognize HF symptoms in the patient (e.g. trouble breathing or ankle swelling), implement treatments for symptoms (e.g. reducing salt in the patient's diet), and evaluating the treatment implemented. The caregiver's confidence in his/her contribution to self-care has 6 items that assess the extent to which the caregiver feels confident contributing to the patient's self-care (e.g. confidence in evaluating the importance of the patient's symptoms). Each CC-SCHFI item uses a 4-point self-report scale. A standardized score from 0 to 100 is computed for each scale; higher scores indicate more caregiver contribution to patient's self-care. Construct and discriminant validity, internal consistency and test-retest reliability of the

CC-SCHFI have been shown in the Italian population (Vellone, Riegel, et al., 2012).

The Short Form 12 [SF-12] (Ware, Kosinski, & Keller, 1996), a generic measure of QOL appropriate for adult populations, was used in both patients and caregivers. The SF-12 is composed of 12 items that are aggregated into two dimensions: the Physical Component Summary (PCS) and the Mental Component Summary (MCS). The PCS items address physical QOL as influenced by physical health, physical functioning, role limitations caused by physical health and bodily pain. The MCS items address mental QOL as influenced by mental health, vitality, role limitations caused by mental health and social functioning. A standardized score ranging from 0 to 100 is obtained from each SF-12 dimension, with higher scores indicating better QOL. The SF-12 has been demonstrated to be a valid and reliable instrument in the Italian population (Gandek et al., 1998) and has been used in several studies to measure QOL in HF patients (Ad, Henry, & Hunt, 2011; Hopman et al., 2009; Prior, Jordan, & Kadam, 2012) and in caregivers of older adults with chronic conditions (Akosile, Okoye, Nwankwo, & Mbada, 2011; Roth, Perkins, Wadley, Temple, & Haley, 2009; Vellone et al., 2011). In this study the SF-12 was administered to both members of the dyad.

Statistical analysis.

The actor-partner interdependence model (APIM) was used to analyze the dyadic data, as this approach allows investigators to evaluate the influence of partners on each other (Ridker, Hennekens, Buring, & Rifai, 2000). In APIM, the actor effect is the influence of a person's emotions on him/herself (e.g. effect of depression on QOL). The partner effect is the impact of one person's emotions on his/her partner (e.g. effect of depression in one person on the partner's QOL). This analytic approach has been applied to adaptation in families of young children with asthma (Ridker, et al., 2000), fear in cancer survivors and their family caregivers (Mellon,

Kershaw, Northouse, & Freeman-Gibb, 2007), distress in couples in therapy (Anderson & Johnson, 2010; Cook & Snyder, 2005), physical and mental health of survivors of advanced prostate cancer and their spousal caregivers (Zhou et al., 2011), QOL of children with cystic fibrosis and their caregivers (Driscoll, Schatschneider, McGinnity, & Modi, 2012), mental and physical health of patients awaiting coronary artery bypass grafting surgery and their partners or family caregivers (Thomson, Molloy, & Chung, 2012), distress in stroke patients and caregivers (Twiddy, House, & Jones, 2012), and psychological well-being in Korean family caregiving dyads (Kim, Reed, Hayward, Kang, & Koenig, 2011).

All analyses were conducted using SPSS for windows, version 20. An α level of .05 was used throughout. Paired sample t-tests were used to provide characteristics of distinguishable dyad members, patients and spousal caregivers. To determine how an individual's QOL was influenced by his/her own self-care and the partner's self-care, we used a multilevel modeling dyadic analysis, APIM with distinguishable dyads (Kenny, Kashy, & Cook, 2006). In this APIM, the residual structure was treated as heterogeneous compound symmetry, including an inter-spousal correlation and separate variances for spouses' QOL. Prior to analysis, data were restructured to a pairwise dyadic data set, and the actor and partner scores on predictors were grand-mean centered and standardized using z-score to obtain both unstandardized and standardized regression coefficients for the actor and partner effects. We also used Pearson product-moment correlation coefficients to identify bivariate association among predictors.

Results

Characteristics of Patient-spouse Dyads

Within the sample of 1192 HF patients, only 339 (28.4%) attended the medical visit with a caregiver. Of these 339 patients, 138 (40.7%) of the caregivers were spouses. These 138 dyads

form the unit of analysis. The mean age of patients was 73.6 years (SD = 9.6) and most were male (64.4%). More than 85% of patients had less than high school education and were retired or unemployed. The median illness duration was almost 4 years. Patients were distributed throughout the four categories of NYHA functional class, although class IV (i.e. symptomatic at rest) was less represented than the other classes. The average ejection fraction was 43% (Table 1). Spousal caregivers were 70.4 years old (SD = 9.4) on average and most were female (67.4%). Education was low for most caregivers as only 22% of them had a high school or university education. Few caregivers (15.9%) were working outside the home (Table 1). Comparing socio-demographic variables within the dyads, patients were 3 years older than caregivers (paired $t = 5.6$, $P < .001$). Patients were more likely to be male while caregivers were more likely to be female ($P < .001$ Fisher exact test). Patients and caregivers were not different in years of education ($P = .45$ Mann Whitney U test) or in employment ($P = .74$ Fisher exact test).

Comparing patients from the dyads studied in our sample to those in the entire sample, the only difference was in comorbidity; patients cared for spousal caregivers had higher comorbidity than the rest of the sample (2.8 vs. 3.4 illnesses, $p = .02$). The two groups were statistically equivalent in the patients' age ($p = .14$), education ($p = .59$), NYHA functional class ($p = .96$), ventricular ejection fraction ($p = .10$), and illness duration ($p = .11$).

Self-care and Quality of Life in Patient-spousal Caregiver Dyads

As shown in Table 2, patients and spousal caregivers reported similar levels of self-care maintenance, management and confidence but, as expected, patients' QOL (i.e., physical QOL and mental QOL) was worse than that of their spousal caregivers. Patients' self-care maintenance, management, and confidence were significantly correlated with those of caregivers, with the strongest correlation found for self-care management (Table 3). Patients'

self-care scores were negatively correlated with their own physical QOL; higher self-care maintenance was associated with worse physical QOL. Self-care management was not significantly correlated with physical QOL of patients or caregivers. However, spousal emotional QOL was significantly correlated with patients' self-care maintenance and confidence. There were no high correlations among predictors; all correlation coefficients were less than .6. Although patients' physical QOL was not significantly correlated with spouses' physical QOL ($r = .013, p > 0.5$), patients' mental QOL was significantly correlated with spouses' mental QOL ($r = .345, p < .001$).

Influence of Self-care on Physical QOL

The results of the APIM analysis are presented in Figures 2 and 3. Details are shown in Table 4. There is evidence for an actor effect of self-care maintenance on physical QOL for both patients and caregivers; individuals higher in self-care maintenance were lower in physical QOL. Specifically, a one unit increase in patient self-care maintenance was associated with a 0.141 unit decrease in physical QOL. For caregivers, a one unit increase in self-care maintenance was associated with 0.098 unit decrease in their physical QOL. For self-care management, the only actor effect was on physical QOL of patients ($b = -.146, p = .034$); higher self-care management was associated with lower physical QOL. The only partner effect of self-care confidence was on physical QOL of caregivers to patients; every one unit increase in the caregiver's confidence was associated with a 0.162 unit decrease in the patient's physical QOL.

Influence of Self-care on Mental QOL

As shown in Figure 3, only one partner effect and one actor effect were found in mental QOL. Self-care maintenance of patients had a significant partner effect on the mental QOL of caregivers; a one unit increase in the patient's self-care maintenance was associated with a .117

unit increase in the caregiver's mental QOL. Actor effects of self-care confidence on mental QOL were significant for caregivers ($b = .124, p = .020$) but the actor effect of self-care confidence on emotional well-being did not reach statistical significance for patients ($b = .103, p = .051$).

Discussion

In this study we used the actor-partner interdependence model to analyze the manner in which adults with HF and their spousal caregivers interact as a unit to influence self-care and QOL. We found both actor and partner effects in this sample of older couples. Surprisingly, higher self-care was related to lower physical QOL in both patients and caregivers. But, the relationship between self-care on mental QOL differed, with higher self-care maintenance in patients associated with better mental QOL in caregivers.

Only two partner effects were seen in this sample. First, we found that better self-care maintenance in patients was associated with better mental QOL in caregivers. This finding that mental QOL of caregivers improved when patients were adhered to therapy captures statistically the energy and emotions that caregivers put into their efforts to support self-care in their loved ones. For example, Rosland and colleagues described how most of the chronically ill patients in their study reported criticism or nagging about treatment adherence from their caregivers (Rosland, Heisler, Choi, Silveira, & Piette, 2010). This result is also consistent with our earlier results from a study of the relationship between shared care in couples and HF self-care (Sebern & Riegel, 2009). In that study we found that caregiver decision making contributed in a positive fashion to patients' self-care maintenance. Finding that patients who follow the treatment regimen and engage in healthy behaviors improves the mental QOL of their spouses is not a surprise to anyone engaged in a long-term relationship.

The second partner effect is not so easily explained; higher caregiver self-care confidence was associated with poorer physical QOL in patients. Some insight into this finding can be gained from a prior study (C. S. Lee, Suwanno, & Riegel, 2009) demonstrating that confidence is an important moderator of the relationship between self-care and health status. That is, self-care confidence can change the strength and direction of the relationship between these variables. Clinically, this interesting relationship between caregiver confidence and patient QOL may indicate that caregivers are assuming more responsibility as the patients they are caring for experience a physical decline. If this is true, the gain in confidence bodes well for the ability of caregivers to continue in the caregiving role (Gallagher et al., 2011; Gonyea, O'Connor, Carruth, & Boyle, 2005; Merluzzi, Philip, Vachon, & Heitzmann, 2011).

Several actor effects were observed in both patients and caregivers. A negative relationship was found between patients' self-care and their physical QOL, which differs from prior results demonstrating that self-care has a positive effect on health status (Jaarsma et al., 1999; Suwanno, Petpichetchian, Riegel, & Issaramalai, 2009). Perhaps this result can be explained by the cross-sectional nature of these data, not causality of the relationship. It may be that patients with poor physical QOL are motivated to engage in better self-care rather than self-care causing a decline in physical QOL. We and others have shown previously that HF patients defer self-care until their functional status deteriorates (Ayas et al., 2003; Riegel, Lee, & Dickson, 2011).

In caregivers, higher contributions to self-care maintenance were associated with poorer physical QOL. This finding can be explained by the physical and emotional efforts required of caregivers to oversee and to provide direct care in order to support and maintain self-care of patients. Numerous prior studies have demonstrated that caregivers are burdened with their

contributions to keeping HF patients stable (Ayas, et al., 2003; Calvert, et al., 2005; Gott, et al., 2006) and high caregiving stress has been associated with poor physical health in caregivers. It is not surprising that caregivers contributing to self-care of patients experience exhaustion that impairs their physical QOL.

In spousal caregivers, higher confidence in their contributions to patients' self-care was associated with better mental QOL. A positive relationship between self-care confidence and QOL has been found in HF patients (Britz & Dunn, 2010) but we are not aware of studies demonstrating this relationship in caregivers. This finding is important because the studies conducted to date describe poor physical and emotional QOL in HF caregivers (Allison & Waterman, 2002; Johnson, 1997; Luttik, Jaarsma, Veeger, & van Veldhuisen, 2005; Pressler, et al., 2012). Identifying a factor associated with better emotional QOL provides a direction for intervention.

We were surprised to not see more of a dyadic effect on self-care. In a prior mixed-methods study we found that one of the primary factors distinguishing those patients who developed expertise in self-care was family engagement; expert patients had more support from engaged loved ones who fostered skill development in self-care (Riegel, Vaughan Dickson, Goldberg, & Deatrack, 2007). It may be that self-care is fairly new in Italy, so the caregivers did not know how to help their loved ones improve their self-care skills. Indeed, their scores on the CC-SCHF were low. This result also might be explained by cultural differences in the interpersonal relationship between couples. Italian spouses may have a different perception about their role as caregivers of an ill family member than US caregivers. Further investigation is needed to explore this hypothesis.

One major clinical implication of these results is a renewed appreciation of the importance of confidence in the performance of self-care. In caregivers, confidence in the ability to support patients in the performance of self-care improved caregivers' mental QOL and was associated with improved physical QOL in patients. Interventions that build caregivers' confidence are needed. In addition, these results add to the growing body of literature describing the effort required of HF patient caregivers. Innovative approaches to support these caregivers are needed.

The major strength of this study is the analytic method, which can yield significant insights into the dynamics that occur in a couple. The major limitation of this study is the cross-sectional nature of the data. Thus, although the analysis was conducted specifying the direction of the relationships, we acknowledge that the dynamics of the relationship are difficult to discern without longitudinal data. Investigators in this area are encouraged to study these relationships over time. Another limitation is that no power analysis was conducted prior to data collection. The sample of spousal caregivers was small, which may have biased our abilities to detect significant relationships. Further, spousal caregivers were available in only a small proportion of the full sample of HF patients. Although this subset of patients appears to be similar to those from the larger sample, there could be a systematic reason explaining the lack of availability of a spousal caregiver. Further, the regression coefficients in the analysis were quite small so readers should not over-interpret these results. For these reasons, the results of this study should be used with caution.

Table 1. Patients' (n = 138) and Caregivers (n = 138) Characteristics on Enrollment

	Patients N (%)	Spousal Caregivers N (%)
Age (mean, SD)	73.6 (9.6)	70.4 (9.4)
Gender		
Male	93 (67.4)	45 (32.6)
Female	45 (32.6)	93 (67.4)
Education		
Elementary	48 (35.0)	47 (34.3)
Middle School	45 (32.8)	41 (29.9)
Professional School	24 (17.5)	18 (13.1)
High School	16 (11.7)	24 (17.1)
University Degree	4 (2.9)	7 (5.1)
Profession		
Employed	19 (13.8)	22 (15.9)
Unemployed	119 (86.2)	116 (84.1)
Illness duration (years, median, interquartile ranges)	3.7 (2.1 - 6)	
NYHA		
I	30 (21.7)	
II	46 (33.3)	
III	49 (35.5)	
IV	13 (9.4)	
Ejection fraction (mean, SD)	43 (10.8)	

NYHA = New York Heart Association
SD = standard deviation

Table 2. Comparisons of self-care and quality of life between patients and spousal caregivers.

Characteristics	Patients	Spousal Caregivers	Paired t	p-value
	Mean (SD)	Mean (SD)		
Self-Care				
Maintenance	53.7 (15.5)	55.2 (16.6)	-.894	.373
Management	51.0 (18.5)	52.4 (16.6)	-.620	.576
Confidence	49.8 (18.5)	52.2 (17.8)	-1.548	.124
Quality of life				
Physical	36.1 (9.7)	43.1 (8.3)	- 6.9	<.001
Mental	43.1 (9.8)	46.6 (9.7)	-3.5	.001

Note: SD= standard deviation

Table 3.

Correlation among predictors and outcome variable.

	1	2	3	4	5	6	7	8	9
1. Patient self-maintenance									
2. Caregiver self-maintenance	.228 **								
3. Patient self-management	.488**	.336**							
4. Caregiver self-management	.444**	.418**	.478**						
5. Patient self-confidence	.423**	.257*	.545**	.412**					
6. Caregiver self-confidence	.230	.357**	.338**	.492**	.508**				
7. Patient physical QOL	-.225**	-.039	-.178	.118	-.050	-.248**			
8. Caregiver physical QOL	-.070	-.203*	-.149	-.077	-.154	-.143	.103		
9. Patient mental QOL	.120	-.075	-.023	.173	.169*	.050	-.082	.107	
10. Caregiver mental QOL	.220**	.193*	-.025	-.010	.185*	.262**	-.182	.092	.304**

** P<.01; * p<.05

Table 5.

The actor and partner effects of self-care on quality of life

Self-care subscales	Actor and partner effects	Physical QOL			Mental QOL		
		<i>b</i>	β	p-value	<i>b</i>	β	p-value
Self-care maintenance	Actor: patient	-.141	-.236	.009	.092	.149	.099
	Actor: caregiver	-.098	-.163	.024	.088	.143	.079
	Partner: patient	.007	.012	.884	-.064	-.104	.216
	Partner: caregiver	-.013	-.022	.773	.117	.190	.031
Self-care management	Actor: patient	-.146	-.253	.034	-.055	-.102	.431
	Actor: caregiver	-.025	-.043	.735	-.005	-.010	.947
	Partner: patient	.147	.253	.055	.140	.259	.076
	Partner: caregiver	-.040	-.069	.547	.002	.003	.980
Self-care confidence	Actor: patient	.053	.099	.295	.103	.187	.051
	Actor: caregiver	-.041	-.076	.377	.124	.227	.020
	Partner: patient	-.162	-.306	.002	-.026	-.048	.627
	Partner: caregiver	-.048	-.092	.266	.037	-.067	.469

Figure 1. The actor and partner effects of self-care on quality of life in patient-caregiver dyads




Figure 2. The actor and partner effects of self-care maintenance, management, confidence on physical quality of life in patient-caregiver dyads (N=138)





Figure 3. The actor and partner effects of self-care maintenance, confidence, management on mental quality of life in patient-caregiver dyads (N=138)



References

- Ad, N., Henry, L., & Hunt, S. (2011). The impact of surgical ablation in patients with low ejection fraction, heart failure, and atrial fibrillation. *European Journal of Cardio-Thoracic Surgery*, 40(1), 70-76. doi:10.1016/j.ejcts.2010.11.016
- Agren, S., Evangelista, L., Davidson, T., & Stromberg, A. (2010). The influence of chronic heart failure in patient-partner dyads-A comparative study addressing issues of health-related quality of life. *Journal of Cardiovascular Nursing*. doi:10.1097/JCN.0b013e3181ec0281
- Agren, S., Evangelista, L. S., Hjelm, C., & Stromberg, A. (2012). Dyads affected by chronic heart failure: a randomized study evaluating effects of education and psychosocial support to patients with heart failure and their partners. *Journal of Cardiac Failure*, 18(5), 359-366. doi:10.1016/j.cardfail.2012.01.014
- Akosile, C. O., Okoye, E. C., Nwankwo, M. J., & Mbada, C. E. (2011). Quality of life and its correlates in caregivers of stroke survivors from a Nigerian population. *Quality of Life Research*, 20(9), 1379-1384. doi:10.1007/s11136-011-9876-9
- Allison, P. D., & Waterman, R. P. (2002). Fixed-effects negative binomial regression models. *Sociological Methodology*, 32, 247-265. doi:10.1111/1467-9531.0017
- Anderson, S. R., & Johnson, L. N. (2010). A dyadic analysis of the between- and within-system alliances on distress. [Multicenter Study]. *Family Process*, 49(2), 220-235. doi:10.1111/j.1545-5300.2010.01319.x
- Ayas, N. T., White, D. P., Manson, J. E., Stampfer, M. J., Speizer, F. E., Malhotra, A., & Hu, F. B. (2003). A prospective study of sleep duration and coronary heart disease in women. *Archives of Internal Medicine*, 163(2), 205-209. doi:10.1001/archinte.163.2.205

- Bennett, S., & Sauve, M. (2003). Cognitive deficits in patients with heart failure: A review of the literature. *Journal of Cardiovascular Nursing, 18*(3), 219-242.
- Britz, J. A., & Dunn, K. S. (2010). Self-care and quality of life among patients with heart failure. *Journal of the American Academy of Nurse Practitioners, 22*(9), 480-487.
doi:10.1111/j.1745-7599.2010.00538.x
- Calvert, M. J., Freemantle, N., & Cleland, J. G. (2005). The impact of chronic heart failure on health-related quality of life data acquired in the baseline phase of the CARE-HF study. *European Journal of Heart Failure, 7*(2), 243-251. doi:10.1016/j.ejheart.2005.01.012
- Cook, W. L., & Snyder, D. K. (2005). Analyzing nonindependent outcomes in couple therapy using the actor-partner interdependence model. *Journal of Family Psychology, 19*(1), 133-141. doi:10.1037/0893-3200.19.1.133
- Driscoll, K. A., Schatschneider, C., McGinnity, K., & Modi, A. C. (2012). Application of dyadic data analysis in pediatric psychology: cystic fibrosis health-related quality of life and anxiety in child-caregiver dyads. *Journal of Pediatric Psychology, 37*(6), 605-611.
doi:10.1093/jpepsy/jss063
- Feetham, S. L. (1991). Conceptual and methodological issues in research of families. In A. L. Whall & J. Fawcett (Eds.), *Family theory development in nursing: State of the science and art*. (pp. 55-68). Philadelphia: F.A. Davis.
- Friedman, M. M. (1997). Older adults' symptoms and their duration before hospitalization for heart failure. *Heart and Lung, 26*, 169-176. doi:10.1016/s0147-9563(97)90053-4
- Frith, C. D., & Frith, U. (2006). How we predict what other people are going to do. *Brain Research, 1079*(1), 36-46. doi:10.1016/j.brainres.2005.12.126

- Gallagher, D., Ni Mhaolain, A., Crosby, L., Ryan, D., Lacey, L., Coen, R. F., . . . Lawlor, B. A. (2011). Self-efficacy for managing dementia may protect against burden and depression in Alzheimer's caregivers. [Research Support, Non-U.S. Gov't]. *Aging & Mental Health, 15*(6), 663-670. doi:10.1080/13607863.2011.562179
- Gandek, B., Ware, J. E., Aaronson, N. K., Apolone, G., Bjorner, J. B., Brazier, J. E., . . . Sullivan, M. (1998). Cross-validation of item selection and scoring for the SF-12 Health Survey in nine countries: Results from the IQOLA Project. International Quality of Life Assessment. *Journal of Clinical Epidemiology, 51*(11), 1171-1178. doi:10.1016/S0895-4356(98)00109-7
- Geden, E. A., & Taylor, S. G. (1999). Theoretical and empirical description of adult couples' collaborative self-care systems. *Nursing Science Quarterly, 12*(4), 329-334. doi:10.1177/08943189922107061
- Go, A. S., Mozaffarian, D., Roger, V. L., Benjamin, E. J., Berry, J. D., Borden, W. B., . . . Stroke Statistics, S. (2012). Heart Disease and Stroke Statistics--2013 Update A Report from the American Heart Association. *Circulation, 127*(1), e6-e245. doi:10.1161/CIR.0b013e31828124ad
- Goldstein, N. E., & Lynn, J. (2006). Trajectory of end-stage heart failure: the influence of technology and implications for policy change. *Perspectives in Biology and Medicine, 49*(1), 10-18. doi:10.1353/pbm.2006.0008
- Gonyea, J. G., O'Connor, M., Carruth, A., & Boyle, P. A. (2005). Subjective appraisal of Alzheimer's disease caregiving: the role of self-efficacy and depressive symptoms in the experience of burden. [Research Support, Non-U.S. Gov't]. *American Journal of*

Alzheimer's Disease & Other Dementias, 20(5), 273-280.

doi:10.1177/153331750502000505

Gott, M., Barnes, S., Parker, C., Payne, S., Seamark, D., Gariballa, S., & Small, N. (2006).

Predictors of the quality of life of older people with heart failure recruited from primary care. *Age and Ageing*, 35(2), 172-177. doi:10.1093/ageing/afj040

Hopman, W. M., Harrison, M. B., Coo, H., Friedberg, E., Buchanan, M., & VanDenKerkhof, E.

G. (2009). Associations between chronic disease, age and physical and mental health status. *Chronic Diseases in Canada*, 29(3), 108-116.

Jaarsma, T., Halfens, R., Abu-Saad, H. H., Dracup, K., Stappers, J., & van Ree, J. (1999).

Quality of life in older patients with systolic and diastolic heart failure. *European Journal of Heart Failure*, 1(2), 151-160. doi:10.1016/s1388-9842(99)00007-0

Johnson, J. E. (1997). Insomnia, alcohol, and over-the-counter drug use in old-old urban women.

Journal of Community Health Nursing, 14(3), 181-188.

doi:10.1207/s15327655jchn1403_5

Kenny, D. A., Kashy, D. A., & Cook, W. L. (2006). *Dyadic data analysis*. New York, NY: The Guilford Press.

Kim, S. S., Reed, P. G., Hayward, R. D., Kang, Y., & Koenig, H. G. (2011). Spirituality and psychological well-being: testing a theory of family interdependence among family caregivers and their elders. *Research in Nursing and Health*, 34(2), 103-115.

doi:10.1002/nur.20425

Kurylo, M., Elliott, T. R., DeVivo, L., & Dreer, L. E. (2004). Caregiver social problem solving abilities and family member adjustment following congestive heart failure. *Journal of*

Clinical Psychology in Medical Settings, 11(3), 151-157.

doi:10.1023/B:JOCS.0000037609.23007.24

- Lee, C. S., Suwanno, J., & Riegel, B. (2009). The relationship between self-care and health status domains in Thai patients with heart failure. *European Journal of Cardiovascular Nursing*, 8(4), 259-266. doi:10.1016/j.ejcnurse.2009.04.002
- Lee, D. T., Yu, D. S., Woo, J., & Thompson, D. R. (2005). Health-related quality of life in patients with congestive heart failure. *European Journal of Heart Failure*, 7(3), 419-422. doi:10.1016/j.ejhert.2004.08.004
- Lewis, E. F., Johnson, P. A., Johnson, W., Collins, C., Griffin, L., & Stevenson, L. W. (2001). Preferences for quality of life or survival expressed by patients with heart failure. *Journal of Heart and Lung Transplantation*, 20(9), 1016-1024. doi:10.1016/s1053-2498(1)00298-4
- Lewis, M. A., McBride, C. M., Pollak, K. I., Puleo, E., Butterfield, R. M., & Emmons, K. M. (2006). Understanding health behavior change among couples: an interdependence and communal coping approach. *Social Science and Medicine*, 62(6), 1369-1380. doi:10.1016/j.socscimed.2005.08.006
- Luttik, M. L., Blaauwbroek, A., Dijker, A., & Jaarsma, T. (2007). Living with heart failure: partner perspectives. *Journal of Cardiovascular Nursing*, 22(2), 131-137.
- Luttik, M. L., Jaarsma, T., Veeger, N. J. G. M., & van Veldhuisen, D. J. (2005). For better and for worse: Quality of life impaired in HF patients as well as in their partners. *European Journal of Cardiovascular Nursing*, 4(1), 11-14. doi:10.1016/j.ejcnurse.2004.12.002
- McMurray, J. J. V., & Stewart, S. (2002). The burden of heart failure (supplement). *European Heart Journal*, 4, D50-D58. doi:10.1016/s1520-760(02)90164-x

- Mellon, S., Kershaw, T. S., Northouse, L. L., & Freeman-Gibb, L. (2007). A family-based model to predict fear of recurrence for cancer survivors and their caregivers. *Psychooncology*, *16*(3), 214-223. doi:10.1002/pon.1074
- Merluzzi, T. V., Philip, E. J., Vachon, D. O., & Heitzmann, C. A. (2011). Assessment of self-efficacy for caregiving: the critical role of self-care in caregiver stress and burden. *Palliative & Supportive Care*, *9*(1), 15-24. doi:10.1017/s1478951510000507
- Molloy, G. J., Johnston, D. W., & Witham, M. D. (2005). Family caregiving and congestive heart failure. Review and analysis. *European Journal of Heart Failure*, *7*(4), 592-603. doi:10.1016/j.ejheart.2004.07.008
- Pressler, S. J., Gradus-Pizlo, I., Chubinski, S. D., Smith, G., Wheeler, S., Sloan, R., & Jung, M. (2012). Family caregivers of patients with heart failure: A longitudinal study. *Journal of Cardiovascular Nursing*. Advanced online publication. doi:10.1097/JCN.0b013e3182563877
- Prior, J. A., Jordan, K. P., & Kadam, U. T. (2012). Influence of chronic diseases on long-term change in physical health: a consultation-survey linkage cohort study in general practice. *Quality of Life Research*, *21*(4), 581-591. doi:10.1007/s11136-011-9974-8
- Retrum, J. H., Nowels, C. T., & Bekelman, D. B. (2012). Patient and caregiver congruence: The importance of dyads in heart failure care. *Journal of Cardiovascular Nursing*, *28*(2), 129-136. doi:10.1097/JCN.0b013e3182435f27
- Ridker, P. M. (2009). C-reactive protein: Eighty years from discovery to emergence as a major risk marker for cardiovascular disease. *Clinical Chemistry*, *55*(2), 209-215. doi:10.1373/clinchem.2008.119214

- Ridker, P. M., Hennekens, C. H., Buring, J. E., & Rifai, N. (2000). C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *New England Journal of Medicine*, 342(12), 836-843. doi:10.1056/NEJM200003233421202
- Riegel, B., Carlson, B., Moser, D. K., Sebern, M., Hicks, F. D., & Roland, V. (2004). Psychometric testing of the self-care of heart failure index. *Journal of Cardiac Failure*, 10(4), 350-360. doi:10.1016/j.cardfail.2003.12.001
- Riegel, B., Lee, C. S., & Dickson, V. V. (2011). Self care in patients with chronic heart failure. *Nature reviews. Cardiology*, 8(11), 644-654. doi:10.1038/nrcardio.2011.95
- Riegel, B., Lee, C. S., Dickson, V. V., & Carlson, B. (2009). An update on the self-care of heart failure index. *Journal of Cardiovascular Nursing*, 24(6), 485-497. doi:10.1097/JCN.0b013e3181b4baa0
- Riegel, B., Vaughan Dickson, V., Goldberg, L. R., & Deatricks, J. A. (2007). Factors associated with the development of expertise in heart failure self-care. *Nursing Research*, 56(4), 235-243. doi:10.1097/01.NNR.000280615.75447.f7
- Rohrbaugh, M. J., Shoham, V., Cleary, A. A., Berman, J. S., & Ewy, G. A. (2009). Health consequences of partner distress in couples coping with heart failure. *Heart & Lung: The Journal of Acute and Critical Care*, 38(4), 298-305. doi:10.1016/j.hrtlng.2008.10.008
- Rosland, A. M., Heisler, M., Choi, H. J., Silveira, M. J., & Piette, J. D. (2010). Family influences on self-management among functionally independent adults with diabetes or heart failure: do family members hinder as much as they help? *Chronic Illness*, 6(1), 22-33. doi:10.1177/1742395309354608
- Roth, D. L., Perkins, M., Wadley, V. G., Temple, E. M., & Haley, W. E. (2009). Family caregiving and emotional strain: associations with quality of life in a large national

- sample of middle-aged and older adults. *Quality of Life Research*, 18(6), 679-688.
doi:10.1007/s11136-009-9482-2
- Satpathy, H. K., Frey, D., Satpathy, R., Satpathy, C., Fleming, A., Mohiuddin, S. M., & Khandalavala, J. (2008). Peripartum cardiomyopathy. *Postgraduate Medicine*, 120(1), 28-32. doi:10.3810/pgm.2008.04.1757
- Sebern, M., & Riegel, B. (2009). Contributions of supportive relationships to heart failure self-care. *European Journal of Cardiovascular Nursing*, 8(2), 97-104.
doi:10.1016/j.ejcnurse.2008.07.004
- Suwanno, J., Petpichetchian, W., Riegel, B., & Issaramalai, S. A. (2009). A model predicting health status of patients with heart failure. *Journal of Cardiovascular Nursing*, 24(2), 118-126. doi:10.1097/JCN.0b013e318197a75c
- Thomson, P., Molloy, G. J., & Chung, M. L. (2012). The effects of perceived social support on quality of life in patients awaiting coronary artery bypass grafting and their partners: Testing dyadic dynamics using the Actor-Partner Interdependence Model. *Psychology Health & Medicine*, 17(1), 35-46. doi:10.1080/13548506.2011.579988
- Twiddy, M., House, A., & Jones, F. (2012). The association between discrepancy in illness representations on distress in stroke patients and carers. *Journal of Psychosomatic Research*, 72(3), 220-225. doi:10.1016/j.jpsychores.2011.12.004
- Ueland, T., Kjekshus, J., Froland, S. S., Omland, T., Squire, I. B., Gullestad, L., . . . Aukrust, P. (2005). Plasma levels of soluble tumor necrosis factor receptor type I during the acute phase following complicated myocardial infarction predicts survival in high-risk patients. *Journal of the American College of Cardiology*, 46(11), 2018-2021.
doi:10.1016/j.jacc.2005.08.039

- Valle, R., Baccichetto, R., Barro, S., Calderan, A., Carbonieri, E., Chinellato, M., . . . Milani, L. (2006). [Heart failure in Eastern Veneto: Prevalence, hospitalization rate, adherence to guidelines and social costs]. *Monaldi Archives for Chest Disease*, *66*(1), 63-74.
- Vellone, E., Fida, R., Cocchieri, A., Sili, A., Piras, G., & Alvaro, R. (2011). Positive and negative impact of caregiving to older adults: a structural equation model. *Professioni Infermieristiche*, *64*(4), 237-248.
- Vellone, E., Piras, G., Venturini, G., Alvaro, R., & Cohen, M. Z. (2012). The experience of quality of life for caregivers of people with Alzheimer's disease living in Sardinia, Italy. *Journal of Transcultural Nursing* *23*(1), 46-55. doi:10.1177/1043659611414199
- Vellone, E., Riegel, B., Cocchieri, A., Barbaranelli, C., D'Agostino, F., Antonetti, G., . . . Alvaro, R. (2013). Psychometric testing of the Self-care of Heart Failure Index version 6.2. Manuscript submitted for publication.
- Vellone, E., Riegel, B., Cocchieri, A., Barbaranelli, C., D'Agostino, F., Glaser, D., . . . Alvaro, R. (2012). Validity and reliability of the caregiver contribution to Self-care of Heart Failure Index. *Journal of Cardiovascular Nursing*. Advanced online publication. doi:10.1097/JCN.0b013e318256385e
- Ware, J., Jr., Kosinski, M., & Keller, S. D. (1996). A 12-Item Short-Form Health Survey: Construction of scales and preliminary tests of reliability and validity. *Medical Care*, *34*(3), 220-233. doi:10.1097/00005650-199603000-00003
- Zhou, E. S., Kim, Y., Rasheed, M., Benedict, C., Bustillo, N. E., Soloway, M., . . . Penedo, F. J. (2011). Marital satisfaction of advanced prostate cancer survivors and their spousal caregivers: The dyadic effects of physical and mental health. *Psychooncology*, *20*(12), 1353-1357. doi:10.1002/pon.1855