8-6-2012

Coarsening of a Two-dimensional Foam on a Dome

A. E. Roth
University of Pennsylvania

C. D. Jones
University of Pennsylvania

Douglas J. Durian
University of Pennsylvania, djdurian@physics.upenn.edu

Follow this and additional works at: https://repository.upenn.edu/physics_papers

---

Recommended Citation

Roth, A. E., Jones, C. D., & Durian, D. J. (2012). Coarsening of a Two-dimensional Foam on a Dome. Retrieved from https://repository.upenn.edu/physics_papers/259

Roth, A. E., Jones, C. D., & Durian, D. J. (2012). Coarsening of a two-dimensional foam on a dome. *Physical Review E*, 86(2), 021402. doi: [10.1103/PhysRevE.86.021402](https://doi.org/10.1103/PhysRevE.86.021402)
© 2012 American Physical Society

This paper is posted at ScholarlyCommons. https://repository.upenn.edu/physics_papers/259
For more information, please contact repository@pobox.upenn.edu.

Coarsening of a Two-dimensional Foam on a Dome

Abstract

In this paper we report on bubble growth rates and on the statistics of bubble topology for the coarsening of a dry foam contained in the narrow gap between two hemispheres. By contrast with coarsening in flat space, where six-sided bubbles neither grow nor shrink, we observe that six-sided bubbles grow with time at a rate that depends on their size. This result agrees with the modification to von Neumann's law predicted by J. E. Avron and D. Levine [[Phys. Rev. Lett. 69, 208 \(1992\)](#)]. For bubbles with a different number of sides, except possibly seven, there is too much noise in the growth rate data to demonstrate a difference with coarsening in flat space. In terms of the statistics of bubble topology, we find fewer three-, four-, and five-sided bubbles, and more bubbles with six or more sides, in comparison with the stationary distribution for coarsening in flat space. We also find good general agreement with the Aboav-Weaire law for the average number of sides of the neighbors of an n -sided bubble.

Disciplines

Physical Sciences and Mathematics | Physics

Comments

Roth, A. E., Jones, C. D., & Durian, D. J. (2012). Coarsening of a two-dimensional foam on a dome. *Physical Review E*, 86(2), 021402. doi: [10.1103/PhysRevE.86.021402](https://doi.org/10.1103/PhysRevE.86.021402)

© 2012 American Physical Society

Coarsening of a two-dimensional foam on a dome

A. E. Roth, C. D. Jones, and D. J. Durian*

Department of Physics & Astronomy, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6396, USA

(Received 11 June 2012; published 6 August 2012)

In this paper we report on bubble growth rates and on the statistics of bubble topology for the coarsening of a dry foam contained in the narrow gap between two hemispheres. By contrast with coarsening in flat space, where six-sided bubbles neither grow nor shrink, we observe that six-sided bubbles grow with time at a rate that depends on their size. This result agrees with the modification to von Neumann's law predicted by J. E. Avron and D. Levine [*Phys. Rev. Lett.* **69**, 208 (1992)]. For bubbles with a different number of sides, except possibly seven, there is too much noise in the growth rate data to demonstrate a difference with coarsening in flat space. In terms of the statistics of bubble topology, we find fewer three-, four-, and five-sided bubbles, and more bubbles with six or more sides, in comparison with the stationary distribution for coarsening in flat space. We also find good general agreement with the Aboav-Weaire law for the average number of sides of the neighbors of an n -sided bubble.

DOI: [10.1103/PhysRevE.86.021402](https://doi.org/10.1103/PhysRevE.86.021402)

PACS number(s): 82.70.Rr

I. INTRODUCTION

Coarsening is a process in foams by which there is diffusion of gas across films such that some bubbles grow and other bubbles shrink. This progresses in such a way that the average bubble area increases over time [1]. Coarsening is not limited to foams and is also relevant in other systems involving domain growth [2,3]. For an ideal dry two-dimensional foam, von Neumann showed that the rate of change of area of a bubble in a two-dimensional foam is [4]

$$\frac{dA}{dt} = K_o(n - 6), \quad (1)$$

where n is the number of sides of a bubble, and K_o is a constant of proportionality. Remarkably, the shape of the bubble, its edge lengths, and its set of neighbors all do not matter.

In 1992 Avron and Levine [5] generalized von Neumann's law to predict the rate of area change for bubbles coarsening on a curved surface. The essential ingredient is that the sum of turning angles around each bubble is no longer 2π , as in flat space but, rather, depends on the integral of Gaussian curvature, κ_G , over the bubble area. This modifies the von Neumann law to

$$\frac{dA}{dt} = K_o \left[(n - 6) + \frac{3}{\pi} \int \kappa_G dA \right]. \quad (2)$$

In the case of a surface of constant positive curvature, such as a dome of radius R , this reduces to

$$\frac{dA}{dt} = K_o \left[(n - 6) + \frac{3A}{\pi R^2} \right]. \quad (3)$$

The rate of change of the bubble area thus depends on the number of sides *and the area* of the bubble.

There have been numerous theoretical and simulation studies of coarsening for foams in two-dimensional flat space [6–14]. But to date we are aware of only one simulation that includes the effect of substrate curvature [15]. There, the authors used a modified Potts model for two-dimensional

foam coarsening on spheres, toroids, and pseudospheres. For spheres, they focus on how the area distribution and average area change over time and find that at late times the dynamics are dominated by the appearance of “singular bubbles” much larger than the average that quickly grow to cover the sphere. There is minimal discussion of the coarsening of individual bubbles and no discussion of side number or other distributions of the system.

While the coarsening of foams in two-dimensional flat space has been well measured [12,16–25], we are unaware of any experiments to test the modified law of Avron and Levine for foam in two-dimensional curved space. However, metallurgical grain growth on curved substrates has been reported. In Ref. [26], the results are said to be preliminary and no growth rate data are shown. In Ref. [27], the deviation from the coarsening rate for flat space is masked by noise, but statistical analysis is reported to demonstrate consistency with Eq. (2). In this paper we use a hemispheric cell to create a curved two-dimensional foam. We use image analysis to track individual bubbles and measure bubble dynamics such as coarsening rate. We also measure bubble statistics, such as the distribution of number of sides, and compare this to results from a flat cell. Our image quality and analysis methods are sufficient to demonstrate directly, for six-sided bubbles, that the growth rates are different from flat space and are consistent with Avron and Levine [5].

II. MATERIALS AND METHODS

To measure coarsening rates of two-dimensional foams on a curved surface, we constructed a cell from two hemispherical polycarbonate domes. The smaller dome has an outer diameter of 12.5 cm, and the larger dome has an inner diameter of 13.3 cm, creating a 4-mm gap. The smaller dome was glued to a flat acrylic plate. The larger dome was placed over the smaller dome and separated from the plate by an O-ring 0.25 in. in diameter. We were careful to ensure that the two domes were aligned concentrically. The upper dome was then screwed to the plate to create a sealed chamber of constant curvature.

The solution we used to create our foam was a liquid consisting of 75% deionized water, 20% glycerin, and 5%

*djdurian@physics.upenn.edu

FIG. 1. (Color online) Sample photograph of a two-dimensional foam, coarsening between nested polycarbonate hemispheres with a 4-mm gap. The scale bar is 1 cm.

Dawn Ultra Concentrated dishwashing liquid. This created a foam that was stable and generally lasted many days. The foam was prepared by putting 35 mL of solution into the chamber (this fills the dome to about 2 cm above the O-ring) and shaking it until a uniform opaque foam was created, with an average bubble size much less than the separation of the domes. The chamber was then left to coarsen until a single layer of bubbles remained between the two domes. This took about 24 h. Two-dimensional coarsening could then be observed for the next 2–4 days. Film ruptures were sometimes observed at the end of this period. Bubble statistics were not taken after any ruptures, though single-bubble dynamics were still considered.

To photograph the foam, the chamber was placed 65 cm above a Vista Point A lightbox, which provided a spatially and temporally uniform light source. A Nikon D80 camera with a Nikkor AF-S 300-mm 1:2.8 D lens was mounted 2.5 m above the chamber. The camera was controlled by a computer to take pictures every 2 min. The apparatus was left to collect pictures, for a period ranging from a few days up to a week. This process was repeated three times to build up statistics. A sample photograph is shown in Fig. 1. Note in this photograph that, especially towards the edge, it is possible to see both the plateau borders on the top dome, as well as the plateau borders on the bottom dome. This makes it difficult to identify the correct boundaries of the cells. We address this issue as part of our image analysis.

After we have taken a series of images of the dome, we perform analysis to get out the true areas of the bubbles on the dome. Note that the images constitute an orthographic projection of a sphere (or hemisphere) onto a plane, where the point of projection is infinity. This projection converts the positions according to the following equations [28]:

$$x = R \cos \varphi \sin \lambda, \quad y = R \sin \varphi, \quad (4)$$

where R is the radius of the sphere, λ is the longitude, φ is the latitude, and the center of the domes is defined as $\lambda = 0$, $\varphi = 0$. The first problem is that in a given image, the plateau borders on both the top and the bottom domes are visible. In order to isolate a single set of plateau borders so that the cells' edges are defined correctly, we recognize that the image is projected in two ways. The plateau borders on the top dome

FIG. 2. Transformation of the photograph in Fig. 1 by the Lambert equal area projection, to get the true areas of the cells. There are errors in this result, but for a region in the center, typically about $4.5 \times 4.5 \text{ cm}^2$, it works well and the areas of these cells can be tracked. This image is skeletonized before any actual areas are measured. The scale bar is 1 cm.

are an orthographic projection of the foam using the radius of the top dome, and the plateau borders on the bottom dome are an orthographic projection of the foam using the radius of the bottom dome; both are combined into the same image. To undo this transformation, we take the inverse of the transformation twice, once using the radius for the top dome and once using the radius for the bottom dome. The resulting two images are thresholded and dilated. The images are then multiplied. This kills the plateau borders that do not correspond to the transformation. That is, the plateau borders from the bottom dome that were transformed using the radius of the top dome are killed, and vice versa. The result is a binary image with the correct latitudes and longitudes of the plateau borders on the dome.

After we have accounted for the fact that the plateau borders on both the inner and the outer domes are visible, we can consider the areas of the individual bubbles. The binary image with the correct latitudes and longitudes of the plateau borders has errors but does well for a region of interest in the center. This resulting image, however, does not preserve the areas of the cells. A simple projection that will preserve areas is the Lambert cylindrical equal area projection. This projection is defined by [28]

$$x = R\lambda, \quad y = R \sin \varphi, \quad (5)$$

where, as before, λ is the longitude and φ is the latitude. The R used here is the average of the two domes used. This produces an image with cells that have distorted shapes but have the same areas as the actual cells on the curved surface. This allows us to track individual bubble areas over time. The result of this image analysis is shown in Fig. 2.

III. BUBBLE DYNAMICS

Using the method of finding the correct areas of individual bubbles described above, it is possible to track the areas of individual bubbles over time. The method of identifying the correct plateau borders and finding the correct areas sometimes has errors of failing to identify a plateau border or adding an

FIG. 3. (Color online) Area versus time for three six-sided bubbles, with initial area, A_0 , as labeled. Lines are a linear fit to the data, from which A_0 was found and subtracted from the data. Vertical error bars represent the perimeter times the pixel size divided by the square root of the number of points in the perimeter; this statistical uncertainty matches the scatter in the data points.

extra one, especially farther from the center, where distortion is greater. It was possible to find bubbles, especially near the center, that could be correctly identified for a significant length of time. The viewing region where bubbles can be tracked is typically about $4.5 \times 4.5 \text{ cm}^2$. It will not necessarily be possible to track all bubbles in this region, but it is very rare for a bubble outside this region to be trackable. The projections were compared to the original images, to ensure that the areas of the tracked bubbles were correct. Images were taken at 200-min intervals and analyzed to get the correct bubble areas. Correctly identified bubbles were labeled and tracked over this period. From this it was possible to get area-versus-time graphs for many bubbles.

We can now consider the form that the area-versus-time traces for the bubbles should take. Avron and Levine's prediction for the coarsening of foam on a spherical substrate of radius R , Eq. (3), is a linear differential equation that can be solved analytically for area versus time:

$$A = \left[A_0 + \frac{\pi R^2}{3} (n-6) \left(1 - e^{-\frac{3K_0 t}{\pi R^2}} \right) \right] e^{\frac{3K_0 t}{\pi R^2}} \quad (6)$$

$$= A_0 + K_0 t \left(n - 6 + \frac{3A_0}{\pi R^2} \right) \left(1 + \frac{3K_0 t}{2\pi R^2} + \dots \right). \quad (7)$$

Here A_0 is the area of a bubble at time 0, and Eq. (7) is the Taylor expansion in $(K_0 t / R^2)$. Note that the standard von Neumann result, $A = A_0 + K_0(n-6)t$, is obtained in the limit $R \rightarrow \infty$.

Example data for the area versus time of three six-sided bubbles with different initial areas are plotted in Fig. 3. There the initial area of each bubble was subtracted off so that the traces are easily comparable. The lines are a linear fit to the data, giving a constant growth rate dA/dt that is positive. It is possible to fit the data to the full exponential form of Eq. (6), but the additional terms in the expansion from Eq. (7) are much, much smaller than the linear term, so it is sufficient to fit to an ordinary line. The key feature in Fig. 3 is that the six-sided bubbles grow, and that the larger ones grow faster. This agrees with Avron and Levine and contrasts strongly with the case of a flat-sided cell, where six-sided bubbles neither grow nor shrink according to the usual von Neumann equation.

FIG. 4. (Color online) Growth rate of the bubble area versus the expected factor from Avron and Levine, Eq. (3), where A is the bubble area and R is the radius of the dome. Error bars represent the uncertainty in slopes of linear fits to bubble area versus time. The solid line is a proportionality fit to all data, with slope $K_0 = 0.00018 \pm 0.00008 \text{ cm}^2/\text{min}$. Inset: Blowup of the data for the $n = 6$ -sided bubbles; growth rates for the three representative bubbles in Fig. 3 are circled.

In particular, in our recent experiments on the coarsening of bubbles in a flat cell [25], where the liquid fraction was varied, the six-sided bubbles all had growth rates scattered around 0 to within statistical uncertainty.

We now measure the growth rate for *all* bubbles, as illustrated in Fig. 3, and we compare it to the expected relationship from Avron and Levine's modification to von Neumann's law. This is plotted in Fig. 4, where the y axis is the coarsening rate, and the x axis is the expected proportionality for a dome of constant curvature given by Eq. (3). Each point represents a single bubble. The line is a proportionality, with slope K_0 , which is the only fitting parameter. The inset is a blowup showing all the six-sided bubbles and highlighting the three bubbles featured in Fig. 3. For six-sided bubbles the growth rates are all positive, except for one or two outliers. There is notable scatter, but the evident trend is that dA/dt increases with bubble size.

Another way to compare growth rate data to Avron and Levine is to plot the coarsening rate against the area, as shown in Fig. 5. There each point represents a single bubble, color-coded by the number of sides. Horizontal lines are the expected relationship from the unmodified von Neumann's law, as seen in Eq. (1), using the same constant of proportionality K_0 as found in Fig. 4. Solid lines are the expected relationship from the modified von Neumann's law, as shown in Eq. (3), again using the same value of K_0 . We see that the data are generally consistent with the predicted modification. This is most evident in the six-sided bubbles, which are virtually all growing. The rate of area change also appears to increase with area for $n = 7$ but is masked by noise for other values of n .

IV. BUBBLE DISTRIBUTIONS

With our system it was also possible to measure distributions such as $p(n)$, the probability that a bubble has n sides, and $m(n)$, the average number of sides of the neighbors of an n -sided bubble. Unlike the case of the flat cell, we do not expect

FIG. 5. (Color online) Growth rate of bubbles versus area. Symbol types distinguish bubbles with different numbers n of sides, as labeled. Horizontal lines are the expected relationship from the uncorrected von Neumann law, shown in Eq. (1), using K_o from Fig. 4. Solid lines are the expected relationship using the corrected von Neumann law from Avron and Levine, shown in Eq. (3), using K_o from Fig. 4. Data and vertical error bars are the same as shown in Fig. 4. Horizontal error bars represent the range of area over which each bubble grew with the specified rate and number of sides.

to reach a scaling state where these statistical quantities remain constant over time. Because the growth rate of a bubble grows with its area, we expect, at long times, to have large bubbles grow rapidly to dominate the system, and this will cause bubble statistics and distributions to change with time. Our system is at much earlier times, where the modification to a bubble's growth rate due to its area is small. This modification still should have some impact, however, and we do indeed find that our statistics deviate from the ordinary scaling state observed in the flat cell.

To measure our statistics, an 8×8 -cm region of interest was defined in the center of the dome, and the number of sides of all bubbles completely or partially within this region was recorded by hand. This was done for photographs at 400-mine intervals from the earliest photograph of a two-dimensional interval to the first occurrence of a rupture, for a total period of typically 48 hs. This was repeated for three runs. These data were used to produce a distribution of the number of sides, which is shown in Fig. 6. Also shown, for comparison, is the distribution found for a flat two-dimensional foam in Ref. [25]. We see that, compared to the flat cell, the dome has a surplus of six-sided bubbles, and a deficit of three-, four-, and five-sided bubbles.

It is also possible to describe these distributions by their average, $\langle n \rangle = 6.06 \pm 0.1$, and by their variance, $\mu_2 = \sum p(n)(n - 6)^2$. We measure the variance to be $\mu_2 = 1.30 \pm 0.05$. This value is lower than measured for the flat cell, $\mu_2 = 1.56 \pm 0.02$ [25], indicating that the width of the distribution is narrower.

FIG. 6. (Color online) Side number distribution for a flat cell and for the dome. For the flat cell, the average number of sides is $\langle n \rangle = 5.92 \pm 0.01$ and the variance $\mu_2 = 1.56 \pm 0.02$. For the dome $\langle n \rangle = 6.06 \pm 0.1$ and $\mu_2 = 1.30 \pm 0.05$. Data for the flat cell are taken from Ref. [25]. Vertical error bars are from a fractional area of 1 over the square root of the number of bubbles with the specified side number.

From these same data we can measure $m(n)$, the average number of sides of the neighbors of an n -sided bubble. We expect $m(n)$ to be related to n by the Aboav-Weaire law, which predicts $m(n) = 6 - a + [(6a - \mu_2)/n]$. In this equation a is

FIG. 7. (Color online) Aboav-Weaire relationship for a flat cell and for a dome. The y axis is $m(n)$, the average number of sides of an n -sided bubble. Lines are fits to the Aboav-Weaire law, $m(n) = 6 - a + [(6a - \mu_2)/n]$, where a is the only fitting parameter. We find $a = 1.1 \pm 0.08$ for the flat cell and $a = 0.96 \pm 0.09$ for the dome [25]. The variances, μ_2 , for these systems are given in the caption to Fig. 6.

a fitting parameter which is usually found to be around 1. Our measurements for $m(n)$ are shown in Fig. 7, along with the measurements of $m(n)$ for a flat cell. Fits to the Aboav-Weaire law are also shown, using the relevant value for μ_2 in each case. We find $a = 1.1 \pm 0.08$ for the flat cell [25] and $a = 0.96 \pm 0.09$ for the dome. We see that in both cases there appears to be more curvature in the data than predicted. The data for the flat cell also seem to fit the form better than for the dome.

V. CONCLUSION

In this experiment we were able to measure both bubble statistics and bubble dynamics of a foam on a curved two-dimensional surface of radius R . For bubble statistics we find that bubbles with few sides are under-represented compared to a two-dimensional foam in flat space. We also find that the Aboav-Weaire law generally holds, though not quite as well for the dome as for the flat cell. For bubble growth rates, in general, it is difficult to observe the effect of the term

added to von Neumann's law by Avron and Levine to account for substrate curvature. Our maximum bubble size is around $A_{\max} = 3.5 \text{ cm}^2$, compared to $R^2 = 41.6 \text{ cm}^2$; therefore, for all our bubbles $3A/(\pi R^2) \ll |n - 6|$ holds, except for $n = 6$. This is why all the data in Fig. 4 lie at x values near $(n - 6)$. Even if we managed to get a single bubble of 20 cm^2 to completely fill our viewing area, the rate of area change would be $dA/dt = K_o[(n - 6) + 0.46]$, so that a seven-sided bubble would have less than a 50% increase in growth rate. For these reasons, the clearest signal we see of curvature effects is that six-sided bubbles grow systematically and do so more rapidly for larger bubbles. The coarsening data as a whole are consistent with Avron and Levine's modification to von Neumann's law to account for coarsening on a curved surface.

ACKNOWLEDGMENTS

We thank J. Rieser for work on preliminary stages of this experiment. This work was supported by NASA Microgravity Fluid Physics Grant No. NNX07AP20G

-
- [1] D. Weaire and S. Hutzler, *The Physics of Foams* (Oxford University Press, New York, 1999).
  - [2] J. Glazier and D. Weaire, *J. Phys.: Condens. Matter* **4**, 1867 (1992).
  - [3] J. Stavans, *Rep. Prog. Phys.* **56**, 733 (1993).
  - [4] J. von Neumann, in *Metal Interfaces* (American Society for Metals, Cleveland, OH, 1952), pp. 108–110.
  - [5] J. E. Avron and D. Levine, *Phys. Rev. Lett.* **69**, 208 (1992).
  - [6] C. W. J. Beenakker, *Phys. Rev. Lett.* **57**, 2454 (1986).
  - [7] J. P. Kermode and D. Weaire, *Comput. Phys. Commun.* **60**, 75 (1990).
  - [8] J. Glazier, M. Anderson, and G. Grest, *Philos. Mag. B* **62**, 615 (1990).
  - [9] T. Herdtle and H. Aref, *J. Fluid Mech.* **241**, 233 (1992).
  - [10] D. Segel, D. Mukamel, O. Krichevsky, and J. Stavans, *Phys. Rev. E* **47**, 812 (1993).
  - [11] H. Flyvbjerg, *Phys. Rev. E* **47**, 4037 (1993).
  - [12] J. Stavans, *Physica A* **194**, 307 (1993).
  - [13] L. Neubert and M. Schreckenberg, *Physica A* **240**, 491 (1997).
  - [14] A. D. Rutenberg and M. B. McCurdy, *Phys. Rev. E* **73**, 011403 (2006).
  - [15] P. Peczak, G. S. Grest, and D. Levine, *Phys. Rev. E* **48**, 4470 (1993).
  - [16] J. A. Glazier, S. P. Gross, and J. Stavans, *Phys. Rev. A* **36**, 306 (1987).
  - [17] J. Stavans and J. A. Glazier, *Phys. Rev. Lett.* **62**, 1318 (1989).
  - [18] J. Stavans, *Phys. Rev. A* **42**, 5049 (1990).
  - [19] K. J. Stine, S. A. Rauser, B. G. Moore, J. A. Wise, and C. M. Knobler, *Phys. Rev. A* **41**, 6884 (1990).
  - [20] B. Berge, A. J. Simon, and A. Libchaber, *Phys. Rev. A* **41**, 6893 (1990).
  - [21] O. Krichevsky and J. Stavans, *Phys. Rev. B* **46**, 10579 (1992).
  - [22] M. de Icaza, A. Jiménez-Ceniceros, and V. M. Castaño, *J. Appl. Phys.* **76**, 7317 (1994).
  - [23] M. E. Rosa and M. A. Fortes, *Philos. Mag. A* **79**, 1871 (1999).
  - [24] A. E. Rosa, L. Afonso, and M. A. Fortes, *Philos. Mag. A* **82**, 2953 (2002).
  - [25] A. E. Roth, C. D. Jones, and D. J. Durian, *arXiv:1207.0146v1*.
  - [26] D. Levine, J. E. Avron, and A. Brokman, *Mater. Sci. Forum* **94-96**, 281 (1992).
  - [27] M. A. Palmer, M. E. Glicksman, and K. Rajan, *Philos. Mag. A* **79**, 763 (1999).
  - [28] G. Kellaway, *Map Projections* (Methuen, London, 1946).