



7-1-1975

Nature of the "Griffiths" Singularity in Dilute Magnets

A. Brooks Harris

University of Pennsylvania, harris@sas.upenn.edu

Follow this and additional works at: https://repository.upenn.edu/physics_papers

 Part of the [Physics Commons](#)

Recommended Citation

Harris, A. (1975). Nature of the "Griffiths" Singularity in Dilute Magnets. *Physical Review B*, 12 (1), 203-207.
<http://dx.doi.org/10.1103/PhysRevB.12.203>

At the time of publication, author A. Brooks Harris was affiliated with Sandia Laboratories, Albuquerque, NM. Currently, he is a faculty member in the Department of Physics at the University of Pennsylvania.

This paper is posted at ScholarlyCommons. https://repository.upenn.edu/physics_papers/401
For more information, please contact repository@pobox.upenn.edu.

Nature of the "Griffiths" Singularity in Dilute Magnets

Abstract

The nature of the singular behavior pointed out by Griffiths for $H=0$ in dilute magnets is investigated. It is argued that for concentration p less than that for formation of an infinite cluster, all derivatives of $M(H)$ are finite. The nonanalyticity in $M(H)$ is due to a branch cut along the imaginary H axis having weight $\exp[-(\text{const})/|H|]$ for $|H|\rightarrow 0$, and is thus too weak to be experimentally observable. Some numerical and exact analytic results for the dilute magnet on a Bethe lattice are presented.

Disciplines

Physics

Comments

At the time of publication, author A. Brooks Harris was affiliated with Sandia Laboratories, Albuquerque, NM. Currently, he is a faculty member in the Department of Physics at the University of Pennsylvania.

Nature of the "Griffiths" singularity in dilute magnets*

A. Brooks Harris[†]

Sandia Laboratories, Albuquerque, New Mexico 87115

(Received 29 October 1974)

The nature of the singular behavior pointed out by Griffiths for $H = 0$ in dilute magnets is investigated. It is argued that for concentration p less than that for formation of an infinite cluster, all derivatives of $M(H)$ are finite. The nonanalyticity in $M(H)$ is due to a branch cut along the imaginary H axis having weight $\exp[-(\text{const})/|H|]$ for $|H| \rightarrow 0$, and is thus too weak to be experimentally observable. Some numerical and exact analytic results for the dilute magnet on a Bethe lattice are presented.

I. INTRODUCTION

Recently there has been great interest in the precise nature of the singularity in the critical behavior of dilute magnets.¹⁻⁴ Some years ago Griffiths² showed that the free energy of a dilute ferromagnet is a nonanalytic function of magnetic field H at $H = 0$ for all temperatures below the transition temperature T_c^0 of the undiluted system. As yet, this singularity has not been detected either by high-temperature expansions^{5,6} or by renormalization-group methods.^{3,4}

Lifshitz⁷ has studied a related problem, namely, the density of states $g(\omega)$ of an electron in a random potential. If the potential at each site V is distributed uniformly over an interval $0 \leq V \leq V_0$, then the density of states near the lower band edge ω_0 is of the form

$$g(\omega) \sim e^{(\omega - \omega_0)^{-\xi}}, \quad (1)$$

where ξ was determined by a single dimensional argument to be $\frac{3}{2}$ for a three-dimensional system. This form results from the relatively infrequent occurrence of large regions having arbitrarily small values of V .

For a ferromagnet it is known^{8,9} that the singularities in the free energy occur at imaginary H , with a density at $H = 0$ which is proportional to the spontaneous magnetization.⁸ As we shall see, an argument similar to that of Lifshitz shows that the magnetization of a randomly dilute ferromagnet has a singularity of the form

$$M(H) \sim \int_{-i\infty}^{+i\infty} \frac{\rho(z)}{H - z} dz, \quad (2)$$

where $\rho(z) \sim e^{-A/|z|}$ for small z and A is nonzero for $T < T_c^0$.¹⁰ In contrast, Domb¹¹ has recently proposed that the Griffiths singularity is a much stronger one, leading to a discontinuity in d^2M/dH^2 at $H = 0$, whereas the arguments we give suggest that all derivatives of $M(H)$ are smooth at $H = 0$ for $p < p_c$, where p_c is the critical concentration for the formation of an infinitely large cluster. Our analysis and conclusions are very similar to those

given by Fisher¹² in his treatment of the cluster theory of condensation.

II. ANALYTIC PROPERTIES

Domb's approach, which we follow here, is to write the magnetization for $p < p_c$ as that of an assembly of separate finite clusters,

$$M(H) = \sum_n W_n(p) M_n(H), \quad (3)$$

where $W_n(p)$ is the probability per site that a cluster of size n is formed and $M_n(H)$ is the corresponding magnetization. Initially we will treat the case $kT \ll J$, where $2J$ is the energy difference between parallel and antiparallel alignment of a pair of spins. Then $M_n(H)$ depends only on n and not on the shape of the cluster. For an Ising system of spin $\frac{1}{2}$ one has

$$M_n(H) = \frac{1}{2} n \tanh(nH/2kT), \quad (4)$$

whereas for a Heisenberg system of spin $\frac{1}{2}$

$$M_n(H) = \frac{1}{2} \left((n+1) \coth \frac{(n+1)H}{2kT} - \coth \frac{H}{2kT} \right). \quad (5)$$

Thus, for an Ising system one has

$$M(H) = \frac{1}{2} \sum_{n=1}^{\infty} W_n(p) n \tanh \frac{nH}{2kT}. \quad (6)$$

Clearly, the terms in Eq. (6) with n finite are analytic for $H = 0$, so it is only the arbitrarily large clusters which produce the Griffiths' singularity. To study the nature of the singularity at $H = 0$ we only need to know the asymptotic form of $W(n)$ for large n . One can write $W_n(p)$ in the form^{11,12}

$$W_n(p) = \sum_s g(n, s) p^n (1-p)^s, \quad (7)$$

where $g(n, s)$ is the number of cluster configurations per site having n sites and s bounding-surface sites. It is clear that $\sum_s g(n, s)$ is less than the corresponding quantity for a Bethe lattice (Cayley tree) having the same number of bonds entering

a vertex. Thus it seems clear that

$$W_n(p) \sim e^{-An} \quad (8)$$

In three dimensions one would expect s in Eq. (7) to be of order $n^{2/3}$, producing a factor $e^{-Bn^{2/3}}$ in Eq. (8) which we drop, since the factor in Eq. (8) gives correctly the dominant behavior.

In fact, Fisher and Essam¹³ give the exact result for the Bethe lattice as

$$W_n(p) = \frac{\sigma(\sigma+1)[n\sigma-1]!}{(n-1)![n\sigma-n+2]!} p^n (1-p)^{\sigma n - n + 2}, \quad (9)$$

where $(\sigma+1)$ is the number of bonds which meet at each site. For large n one obtains Eq. (8) with

$$A = (1-\sigma) \ln \left(\frac{1-p}{1-\sigma p} \right) - \ln p \sigma \quad (10)$$

so that $A > 0$ for $p \neq p_c$, where $p_c = \sigma^{-1}$.

Thus the analytic properties of $M(H)$ are determined by using Eq. (8) in Eq. (6). A convergent power series for $M(H)$ at $H=0$ does not exist, because $M(H)$ has a branch cut along the imaginary H axis caused by the poles in $\tanh(nH/2kT)$ at rational values of $H/\pi kT$. However, an asymptotic expansion for $M(H)$ at $H=0$ can be generated by expanding $\tanh(nH/2kT)$ in powers of H/kT . To proceed further we replace the sum over n in Eq. (6) by an integral over n from $n=0$ to $n=\infty$. This replacement will not affect the nonanalytic contribution from large n . By suitable changes of variables one then obtains Eq. (2) with

$$\rho(z) = i \pi (kT)^2 Q^{1/2} (1+Q)(1-Q)^{-2} z^{-2}, \quad (11)$$

where $Q(z) = e^{-2A\pi kT/|z|}$, with A given in Eq. (10). This result again shows that all derivatives of $M(H)$ are finite at $H=0$. In the Appendix we show the error in the analysis of Ref. 11 which leads to a different result.

A crude analysis of Eq. (6) can be made by recognizing that $n \tanh(nH/2kT)$ is proportional to $n^2 H$ for $nH \ll kT$ and to n for $nH \gg kT$. No matter how small H may be, this crossover behavior creates an anomalous variation in $M(H)$, thus causing a singularity. At finite temperatures for sufficiently large n one will still have $M_n(H) \sim n^2 H$ for $nH \ll kT$ and $nH \ll J$ providing $T < T_c^0$. Thus, for large n we set

$$M_n(H) \sim n^2 H [M_0(T)/M_0(0)]^2, \quad nH \ll kT, \quad (12)$$

where $M_0(T)$ is the spontaneous magnetization of the infinite system. Since $M_0(T)$ is only nonzero for $kT \lesssim J$, the condition $nH \ll J$ is redundant in Eq. (12). Equation (12) remains valid for the Heisenberg model, so we suggest that the form

of the singularity in Eqs. (2) and (11) is appropriate for both Heisenberg and Ising models for $T < T_c^0$.

III. RESULTS FOR THE BETHE LATTICE

In this section we present several analytic and numerical results for the Bethe lattice. While the Bethe lattice does have some properties uncharacteristic of three-dimensional lattices, the general trend of the results we obtain seems appropriate for three-dimensional systems.

Since $W_n(p)$ is known exactly, the zero-temperature value of any order derivative of $M(H)$ at $H=0$ can in principle be evaluated in closed form. We have calculated χ and $d^2\chi/dH^2$ at $H=0$ for the Ising (I) and Heisenberg (H) systems, using Eqs. (4) and (5), respectively. We write results as

$$kT\chi_I = \frac{1}{4} \langle n^2 \rangle, \quad (13)$$

$$kT\chi_H = \frac{1}{12} [\langle n^2 \rangle + 2\langle n \rangle], \quad (14)$$

$$(kT)^3 \frac{d^2\chi_I}{dH^2} = -\frac{1}{8} \langle n^4 \rangle, \quad (15)$$

$$(kT)^3 \frac{d^2\chi_H}{dH^2} = -\frac{1}{120} [\langle n^4 \rangle + 4\langle n^3 \rangle + 6\langle n^2 \rangle + 4\langle n \rangle], \quad (16)$$

where $\langle n^r \rangle = \sum_n W_n(p) n^r$. Using Eq. (9) for $W_n(p)$ one finds

$$\langle n \rangle = p, \quad (17)$$

$$\langle n^2 \rangle = p(1+p)/(1-\sigma p), \quad (18)$$

$$\langle n^3 \rangle = p(1+3p-3\sigma p^2-\sigma p^3)/(1-\sigma p)^3, \quad (19)$$

$$\langle n^4 \rangle = \langle n^3 \rangle(1+p)/(1-\sigma p) + 3(\sigma+1)p^2(1-p)(1-\sigma p^2)/(1-\sigma p)^5. \quad (20)$$

[To obtain these results it is convenient to evaluate derivatives of $K^s(x, y)$ given in Ref. 13.] The values of $\hat{\chi} \equiv kT\chi$ and $\hat{\chi}'' \equiv -(kT)^3 d^2\chi/dH^2$ at $H=0$ are shown as a function of p for $\sigma+1=6$ in Fig. 1.

There one sees the striking divergence in $d^2\chi/dH^2$ as $p \rightarrow p_c$. In fact, from Eq. (10) one sees that $A \sim |p_c - p|^2$ for $p \rightarrow p_c$, so that $\langle n^{r+1} \rangle / \langle n^r \rangle \sim |p_c - p|^{-2}$. We have explicitly verified that $\langle n^r \rangle \sim |p_c - p|^{2r-3}$ as $p \rightarrow p_c$ for $r \geq 2$. Thus, succeeding even-order derivatives of χ diverge increasingly strongly as $p \rightarrow p_c$. Even for p fixed the zero-field derivatives for large r can be estimated to obey $(d^{2r}\chi/dH^{2r}) / (d^{2r-2}\chi/dH^{2r-2}) \sim r^4$ in view of the asymptotic form $\langle n^r \rangle \sim A^{-r} r!$.¹⁴

These results are illustrated by the numerical evaluations shown in Figs. 2 and 3. There one sees that $-(kT)^3 d^2\chi/dH^2$ [i. e., the slope of $M''(H)$] is an order of magnitude larger than $kT\chi$ [the slope of $M(H)$]. Also, the region in which M is nearly a linear function of H is very much larger than the corresponding linear region for d^2M/dH^2 . This effect becomes more pronounced as p approaches p_c . Still higher derivatives will be larger and have

smaller linear regimes. So, if measurements were taken for $H > H_0$, then for some r depending on the size of H_0 one would find an apparent discontinuity in $d^{2r}M/dH^{2r}$. Nonetheless, the true analytic behavior is that all derivatives of $M(H)$ are continuous and the even-order ones vanish as $H \rightarrow 0$. Also, in conformity with Eqs. (13)–(16) one sees from Figs. 2 and 3 that M and its derivatives are noticeably smaller for the Heisenberg model than for the Ising model.

Finally, we conclude this section by giving some exact analytic results for finite temperatures. The following discussion will be confined to the paramagnetic region, i. e., for¹⁵

$$p\sigma \tanh(J/kT) < 1, \tag{21}$$

which, for $p < p_c = \sigma^{-1}$, includes all temperatures. In Ref. 15 we give the exact result for $\chi(T, H=0)$ as

$$4kT\chi(T, H=0) = p(1 + pt)/(1 - p\sigma t) \tag{22}$$

in the units of the present paper, where $t = \tanh(J/kT)$.

We now evaluate $d^2\chi/dH^2$ at $H=0$ as a function of




FIG. 1. Zero-field and zero-temperature values of the reduced susceptibility $\hat{\chi} \equiv kT\chi$, solid line, and $\hat{\chi}'' \equiv -(kT)^3 d^2\chi/dH^2$, broken line, for the Ising (I) and Heisenberg (H) models as a function of concentration for a Bethe lattice with $\sigma+1=6$. Note the difference in scales: that for $\hat{\chi}''$ for the Ising model is on the right; that for the other curves is on the left.




FIG. 2. Zero-temperature values of $M(H)$, full line, and $M''(H) \equiv -(kT)^2 d^2M/dH^2$, broken line, versus H for the Ising model on a Bethe lattice with $\sigma+1=6$ for $p=0, 10$ and $p=0, 14$ ($p_c=0.2$). The scale for $M(H)$ is on the left; that for $M''(H)$ is on the right.




FIG. 3. As in Fig. 2, but for the Heisenberg model.




FIG. 4. Topological structure of the various terms contributing to $d^2\chi/dH^2$ for $H=0$ for the Bethe lattice. The forks [viz., in diagrams (h)–(k)] indicate that the paths between two or more of the labeled sites overlap in part.

$$\begin{aligned}
 -16(kT)^3 \frac{d^2\chi}{dH^2} = & 2p + \frac{8p^2t(\sigma+1)}{1-\sigma pt} + \frac{6p^2t^2(\sigma+1)}{1-\sigma pt^2} + \frac{12p^3t^2(\sigma+1)\sigma}{1-\sigma pt} \left[\frac{1}{1-\sigma pt} + \frac{2t}{1-\sigma pt^2} \right] \\
 & + \frac{24p^4t^3\sigma(\sigma^2-1)}{(1-\sigma pt)^2(1-\sigma pt^2)} \left[\frac{1}{3} \frac{1-\sigma pt^2}{1-\sigma pt} + \frac{\sigma}{\sigma-1} + \frac{1}{2} \right] + 2p^5t^5\sigma(\sigma^2-1) \left[\frac{12\sigma}{(1-\sigma pt)^3} \frac{\sigma-2}{1-\sigma pt^2} + \frac{\sigma-2}{1-\sigma pt} \right] + \frac{6p^6t^6\sigma^2(\sigma^2-1)(\sigma-1)}{(1-\sigma pt)^4(1-\sigma pt^2)}. \quad (26)
 \end{aligned}$$

These terms represent the contributions of diagrams 4(a)–4(k), respectively. Some numerical evaluations of Eqs. (22) and (26) are shown in Fig. 5. As expected, both $kT\chi$ and $(kT)^3 d^2\chi/dH^2$ are monotonic functions of both p and T .

IV. CONCLUSION

We conclude that $M(H)$ has a branch cut along the imaginary axis with exponentially small weight near $H=0$. All derivatives of $M(H)$ are finite at $H=0$. Numerically, the higher-order derivatives become large, particularly near $p=p_c$, so that an experimental determination of the exact nature of the sin-

temperature. For $H=0$ we may write

$$\begin{aligned}
 16(kT)^3 \frac{d^2\chi}{dH^2} = & \sum_{ijkl} [\langle \sigma_i \sigma_j \sigma_k \sigma_l \rangle - \langle \sigma_i \sigma_j \rangle \langle \sigma_k \sigma_l \rangle \\
 & - \langle \sigma_i \sigma_k \rangle \langle \sigma_j \sigma_l \rangle - \langle \sigma_i \sigma_l \rangle \langle \sigma_j \sigma_k \rangle], \quad (23)
 \end{aligned}$$

since all odd-power averages of σ 's vanish. We classify the terms on the right-hand side of Eq. (23) into 11 topologically distinct classes, as shown in Fig. 4. We use the result for the Bethe lattice for $p=1$,

$$\langle \sigma_i \sigma_j \rangle = t^{d_{ij}}, \quad (24)$$

where d_{ij} is the distance between sites i and j . For averages of four σ 's we have similar results. For terms having the topological structure of diagrams shown in Figs. 4(d)–4(g) we have, respectively,

$$\langle \sigma_i \sigma_j \sigma_k \sigma_l \rangle = t^{d_{jkh}} \quad (25a)$$

$$= t^{d_{jkh}} \quad (25b)$$

$$= t^{d_{ij}+d_{ik}+d_{il}} \quad (25c)$$

$$= t^{d_{ij}+d_{kl}}. \quad (25d)$$

To evaluate Eq. (23) we combine these results for the averages at $p=1$ with a factor of p for each site in the diagram. The counting of diagrams then proceeds as usual for a high-temperature expansion. Since we sum all terms in the high-temperature expansion our results are valid throughout the paramagnetic region. We find that

gularity at $H=0$ is probably impossible.

ACKNOWLEDGMENTS

I would like to thank Dr. W. J. Camp for suggesting the use of the results of Ref. 13 for the Bethe lattice. His critical comments and those of Dr. P. M. Richards are gratefully acknowledged.

APPENDIX: ANALYSIS OF REF. 11

In Ref. 11 Domb writes a set of equations, Eq. (1) and (4), equivalent to

$$kT \frac{\partial M}{\partial H} = \sum_n W_n(p) n^2 \operatorname{sech}^2 nH/kT, \quad (A1)$$




FIG. 5. Zero-field values of $\hat{\chi} = kT\chi$, solid line, and $\hat{\chi}'' = -(kT)^2 d^2\chi/dH^2$, broken line, for the Ising model on a Bethe lattice with $(\sigma+1)=6$ versus concentration at various temperatures. For the pure ($\phi=1$) system the transition temperature is $kT_c \approx 4.9J$.

which he approximates as

$$kT \frac{\partial M}{\partial H} = \sum_n W_n(\phi) n^2 e^{-2n|H|/kT}. \quad (A2)$$

We claim that this approximation is inappropriate for analyzing the singularity for $H \rightarrow 0$. To see this, differentiate Eq. (A2):

$$-(kT)^2 \frac{\partial^2 M}{\partial H^2} = 2\theta(H) \sum_n W_n(\phi) n^3 e^{-2n|H|/kT}, \quad (A3)$$

where $\theta(H) = H/|H|$. According to Eq. (A3) $\partial^2 M/\partial H^2$

has a discontinuity for $H \rightarrow 0$ given by

$$\left. \frac{\partial^2 M}{\partial H^2} \right|_{H \rightarrow 0^+} - \left. \frac{\partial^2 M}{\partial H^2} \right|_{H \rightarrow 0^-} = -\frac{4}{(kT)^2} \sum_n W_n(\phi) n^3. \quad (A4)$$

On general grounds we know that the contribution to the left-hand side of Eq. (A4) from finite-sized clusters is zero. However, the right-hand side of Eq. (A4) incorrectly has nonvanishing contributions from finite-sized clusters. Thus we conclude that Eq. (A2) does not correctly represent the low-field singularity contained in Eq. (A1).

*This work supported by the U. S. Atomic Energy Commission.

[†]Permanent address: Department of Physics, University of Pennsylvania, Philadelphia, Pa. 19174.

¹B. M. McCoy and T. T. Wu, Phys. Rev. Lett. **21**, 549 (1968).

²R. B. Griffiths, Phys. Rev. Lett. **23**, 17 (1968).

³A. Luther and G. Grinstein, AIP Conf. Proc. **18**, 876 (1974).

⁴A. B. Harris and T. C. Lubensky, Phys. Rev. Lett. **33**, 1540 (1974).

⁵D. C. Rapaport, J. Phys. C **5**, 1830 (1972).

⁶G. S. Rushbrooke, R. A. Muse, R. L. Stephenson, and K. Pirnie, J. Phys. C **5**, 3371 (1972).

⁷I. M. Lifshitz, Adv. Phys. **13**, 483 (1964).

⁸T. D. Lee and C. N. Yang, Phys. Rev. **87**, 410 (1952).

⁹T. Asano, Phys. Rev. Lett. **24**, 1409 (1970); J. Phys. Soc. Jpn. **29**, 350 (1970).

¹⁰In the range of temperature and concentration for which the spontaneous magnetization $M_0(T)$ is nonzero, i.e., for $T < T_c(\phi)$, there will of course be a singular contribution from the infinite cluster of the same form as for the pure system, viz., $M(H, T) \sim (H/|H|)M_0(T)$ for $H \rightarrow 0$. For an Ising system for $T \neq T_c(\phi)$, $\chi(H)$ is probably

finite as $H \rightarrow 0$, and one can separate the singular contributions of the infinite cluster from those of the finite clusters considered in the text. In contrast, for a Heisenberg model $\chi(H)$ diverges as $H \rightarrow 0$, so such a separation may be impossible.

¹¹C. Domb, J. Phys. C **7**, 2677 (1974). In the note added in proof Domb cites an observation by Fisher that the approximation Domb used is inadequate for a discussion of the Griffiths singularity and that a more careful treatment indicates no discontinuity but rather an essential singularity in d^2M/dH^2 . Technically speaking, the singularity we find in Eqs. (2) and (11) is not an essential singularity inasmuch as it is not an isolated one. In a very recent paper M. Wortis obtained conclusions similar to those obtained here. See M. Wortis, Phys. Rev. B **10**, 4665 (1974).

¹²M. E. Fisher, Physics **3**, 255 (1967).

¹³M. E. Fisher and J. W. Essam, J. Math. Phys. **2**, 609 (1961).

¹⁴Note that since $\tanh(nH/2kT)$ has a pole at $H = i\pi kT/n$, we may estimate that for $H = 0$, $(d/dH)^{2r+1} \tanh(nH/2kT) \sim (2r+1)! (n/\pi kT)^{2r+1}$ for large r . Thus $(d/dH)^{2r} \chi \sim (2r+1)! \langle n^{2r+2} \rangle \sim (2r+1)! (2r+2)!$.

¹⁵A. B. Harris, J. Phys. C **7**, 1671 (1974).