12-2011

On the Minimum Size of a Point Set Containing a 5-Hole and a Disjoint 4-Hole

Bhaswar B. Bhattacharya
University of Pennsylvania

Sandip Das

Follow this and additional works at: https://repository.upenn.edu/statistics_papers

Part of the [Applied Mathematics Commons](#), [Business Commons](#), [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

Recommended Citation

Bhattacharya, B. B., & Das, S. (2011). On the Minimum Size of a Point Set Containing a 5-Hole and a Disjoint 4-Hole. *Studia Scientiarum Mathematicarum Hungarica*, 48 (4), 445-457. <http://dx.doi.org/10.1556/SScMath.2011.1173>

This paper is posted at ScholarlyCommons. https://repository.upenn.edu/statistics_papers/652
For more information, please contact repository@pobox.upenn.edu.

On the Minimum Size of a Point Set Containing a 5-Hole and a Disjoint 4-Hole

Abstract

Let $H(k, l)$, $k \leq l$ denote the smallest integer such that any set of $H(k, l)$ points in the plane, no three on a line, contains an empty convex k -gon and an empty convex l -gon, which are disjoint, that is, their convex hulls do not intersect. Hosono and Urabe [*JCDCC*, LNCS 3742, 117–122, 2004] proved that $12 \leq H(4, 5) \leq 14$. Very recently, using a Ramsey-type result for disjoint empty convex polygons proved by Aichholzer et al. [*Graphs and Combinatorics*, Vol. 23, 481–507, 2007], Hosono and Urabe [*Kyoto CGGT*, LNCS 4535, 90–100, 2008] improve the upper bound to 13. In this paper, with the help of the same Ramsey-type result, we prove that $H(4, 5) = 12$.

Keywords

primary 52C10, 52A10, convex hull, discrete geometry, empty convex polygons, Erdős-Szekeres theorem, Ramsey-type results

Disciplines

Applied Mathematics | Business | Mathematics | Statistics and Probability

On the Minimum Size of a Point Set Containing a 5-Hole and a Disjoint 4-Hole

Bhaswar B. Bhattacharya¹ and Sandip Das²

¹ Indian Statistical Institute, Kolkata, India, bhaswar.bhattacharya@gmail.com

² Advanced Computing and Microelectronics Unit, Indian Statistical Institute, Kolkata, India, sandipdas@isical.ac.in

Abstract. Let $H(k, l)$ denote the smallest integer such that any set of $H(k, l)$ points in the plane, no three on a line, contains an empty convex k -gon and an empty convex l -gon, which are disjoint, that is, their convex hulls do not intersect. Hosono and Urabe [*JCD CG*, LNCS 3742, 117-122, 2004] proved that $12 \leq H(4, 5) \leq 14$. Very recently, using a Ramsey-type result for disjoint empty convex polygons proved by Aichholzer et al. [*Graphs and Combinatorics*, Vol. 23, 481-507, 2007], Hosono and Urabe [*Kyoto CGGT*, LNCS 4535, 90-100, 2008] improve the upper bound to 13. In this paper, with the help of the same Ramsey-type result, we prove that $H(4, 5) = 12$.

Keywords. Convex hull, Discrete geometry, Empty convex polygons, Erdős-Szekeres theorem, Ramsey-type results.

1 Introduction

The famous Erdős-Szekeres theorem [7] states that for every positive integer m , there exists a smallest integer $ES(m)$, such that any set of at least $ES(m)$ points in the plane, no three on a line, contains m points which lie on the vertices of a convex polygon. Evaluating the exact value of $ES(m)$ is a long standing open problem. A construction due to Erdős [8] shows that $ES(m) \geq 2^{m-2} + 1$, which is also conjectured to be sharp. It is known that $ES(4) = 5$ and $ES(5) = 9$ [15]. Following a long computer search, Szekeres and Peters [19] recently proved that $ES(6) = 17$. The value of $ES(m)$ is unknown for all $m > 6$. The best known upper bound for $m \geq 7$ is due to Toth and Valtr [20]: $ES(m) \leq \binom{2m-5}{m-3} + 1$. For a more detailed description of the Erdős-Szekeres theorem and its numerous ramifications, see the surveys by Bárány and Károlyi [4] and Morris and Soltan [16].

In 1978 Erdős [6] asked whether for every positive integer k , there exists a smallest integer $H(k)$, such that any set of at least $H(k)$ points in the plane, no three on a line, contains k points which lie on the vertices of convex polygon whose interior contains no points of the set. Such a subset is called an *empty convex k -gon* or a *k -hole*. Esther Klein showed $H(4) = 5$ and Harborth [10] proved that $H(5) = 10$. Horton [11] showed that it is possible to construct arbitrarily large set of points without a 7-hole, thereby proving that $H(k)$ does not exist for $k \geq 7$. Recently, after a long wait, the existence of $H(6)$ has been proved by Gerken [9] and independently by Nicolás [17]. Later Valtr [22] gave a simpler version of Gerken's proof.

The problems concerning disjoint holes, that is, empty convex polygons with disjoint convex hulls, was first studied by Urabe [21] while addressing the problem of partitioning of planar point sets. For any set S of points in the plane, denote by $CH(S)$ the *convex hull* of S . Given a set S of n points in the plane, no three on a line, a *disjoint convex partition* of S is a partition of S into subsets $S_1, S_2, \dots, S_t$, with $\sum_{i=1}^t |S_i| = n$, such that for each $i \in \{1, 2, \dots, t\}$, $CH(S_i)$ forms a $|S_i|$ -gon and $CH(S_i) \cap CH(S_j) = \emptyset$, for any pair of indices

i, j . Observe that in any disjoint convex partition of S , the set S_i forms a $|S_i|$ -hole and the holes formed by the sets S_i and S_j are disjoint for any pair of distinct indices i, j . If $F(S)$ denote the minimum number of disjoint holes in any disjoint convex partition of S , then $F(n) = \max_S F(S)$, where the maximum is taken over all sets S of n points, is called the *disjoint convex partition number* for all sets of fixed size n . The disjoint convex partition number $F(n)$ is bounded by $\lceil \frac{n-1}{4} \rceil \leq F(n) \leq \lceil \frac{5n}{18} \rceil$. The lower bound is by Urabe [21] and the upper bound by Hosono and Urabe [14]. The proof of the upper bound uses the fact that every set of 7 points in the plane contains a 3-hole and a disjoint 4-hole. Later, Xu and Ding [25] improved the lower bound to $\lceil \frac{n+1}{4} \rceil$.

Another class of related problems arise if the condition of disjointness is relaxed. Given a set S of n points in the plane, no three on a line, a *empty convex partition* of S is a partition of S into subsets $S_1, S_2, \dots, S_t$, with $\sum_{i=1}^t |S_i| = n$, such that for each $i \in \{1, 2, \dots, t\}$, $CH(S_i)$ forms a $|S_i|$ -hole in S . In this case, $CH(S_i)$ and $CH(S_j)$ may intersect for some pair of distinct indices i and j . If $G(S)$ denote the minimum number of holes in any empty convex partition of S , then the *empty convex partition number* for all sets of fixed size n is $G(n) = \max_S G(S)$, where the maximum is taken over all sets S of n points. Urabe [21] proved that $\lceil \frac{n-1}{4} \rceil \leq G(n) \leq \lceil \frac{3n}{11} \rceil$. Xu and Ding [25] improved the bounds to $\lceil \frac{n+1}{4} \rceil \leq G(n) \leq \lceil \frac{5n}{14} \rceil$. The upper bound bound was further improved to $\lceil \frac{9n}{34} \rceil$ by Ding et al. [5].

In [14], Urabe defined the function $F_k(n) = \min_S F_k(S)$, where $F_k(S)$ is the maximum number of k -holes in a disjoint convex partition of S , and the minimum being taken over all sets S of n points. Using the fact that the minimum size of a point set containing two disjoint 4-holes is 9, they showed that $F_4(n) \geq \lfloor \frac{5n}{22} \rfloor$. Recently, Wu and Ding [23] defined $G_k(n) = \min_S G_k(S)$, where $G_k(S)$ is the maximum number of k -holes in a empty convex partition of S , and the minimum being taken over all sets S of n points. They proved that $G_4(n) \geq \lfloor \frac{9n}{38} \rfloor$. The problem of obtaining non-trivial lower bounds on $F_5(n)$ and $G_5(n)$ remains open.

Hosono and Urabe [13] also introduced the function $H(k, l)$, $k \leq l$, which denotes the smallest integer such that any set of $H(k, l)$ points in the plane, no three on a line, contains both a k -hole and a l -hole which are disjoint. Clearly, $H(3, 3) = 6$ and Horton's result [11] implies that $H(k, l)$ does not exist for all $l \geq 7$. Urabe [21] showed that $H(3, 4) = 7$, while Hosono and Urabe [14] showed that $H(4, 4) = 9$. Hosono and Urabe [13] also proved that $H(3, 5) = 10$ and $12 \leq H(4, 5) \leq 14$. The results $H(3, 4) = 7$ and $H(4, 5) \leq 14$ were later reconfirmed by Wu and Ding [24]. Very recently, using a Ramsey-type result for disjoint empty convex polygons proved by Aichholzer et al. [1], Hosono and Urabe [12] proved that $12 \leq H(4, 5) \leq 13$, thus improving upon their earlier result.

In this paper, using the same Ramsey-type result, we evaluate the exact value of $H(4, 5)$, thereby improving upon the result of Hosono and Urabe [12], as stated in the following theorem.

Theorem 1. $H(4, 5) = 12$.

While addressing the problem of pseudo-convex decomposition, Aichholzer et al. [1] proves the following theorem with the help of the order type data base ([2], [3]). Here, we use this result to prove Theorem 1.

Theorem 2. [1] *Every set of 11 points in the plane, no three on a line, contains either a 6-hole or a 5-hole and a disjoint 4-hole.*

The outline of the proof of Theorem 1 is as follows. Consider a set S of 12 points in the plane, no three on a line. Theorem 2 implies that S always contains a 6-hole or a 5-

hole and a 4-hole, which are disjoint. If S contains a 5-hole and a disjoint 4-hole, we are done. Therefore, it suffices to assume that S contains a 6-hole. Next, we show that if S contains a 7-hole, then S contains a 5-hole and a disjoint 4-hole. Thus, we assume that S contains a 6-hole, which cannot be extended to a 7-hole. Then we consider a subdivision of the exterior of the 6-hole and prove the existence a 5-hole and a disjoint 4-hole for all the different possible distributions of the remaining 6 points in the regions formed by the subdivision. The formal proof of Theorem 1 is presented in Section 3.

2 Definitions and Notations

We first introduce the definitions and notations required for the remaining part of the paper. Let S be a finite set of points in the plane in general position, that is, no three on a line. Denote the *convex hull* of S by $CH(S)$. The boundary vertices of $CH(S)$, and the points of S in the interior of $CH(S)$ are denoted by $\mathcal{V}(CH(S))$ and $\mathcal{I}(CH(S))$, respectively. A region R in the plane is said to be *empty* in S if R contains no elements of S in its interior. Moreover, for any set T , $|T|$ denotes the cardinality of T .

By $P := p_1p_2 \dots p_k$ we denote a convex k -gon with vertices $\{p_1, p_2, \dots, p_k\}$ taken in anti-clockwise order. $\mathcal{V}(P)$ denotes the set of vertices of P and $\mathcal{I}(P)$ the interior of P . The collection of all points $q \in \mathbb{R}^2$ such that $\{q\} \cup \mathcal{V}(P)$ form a convex $(k+1)$ -gon is called the *forbidden zone* of P . The forbidden zone of the pentagon $P := p_1p_2p_3p_4p_5$ is the shaded region as shown in Figure 1(a).

For any three points $p, q, r \in S$, $\mathcal{H}(pq, r)$ denotes the open halfplane bounded by the line pq containing the point r . Similarly, $\overline{\mathcal{H}}(pq, r)$ is the open halfplane bounded by pq not containing the point r . Moreover, if $\angle rpq < \pi$, $\text{Cone}(rpq)$ denotes the interior of the angular domain $\angle rpq$. A point $s \in \text{Cone}(rpq) \cap S$ is called the *nearest angular neighbor* of $\overrightarrow{pq}$ in $\text{Cone}(rpq)$ if $\text{Cone}(spq)$ is empty in S . Similarly, for any convex region R a point $s \in R \cap S$ is called the *nearest angular neighbor* of $\overrightarrow{pq}$ in R if $\text{Cone}(spq) \cap R$ is empty in S . More generally, for any positive integer k , a point $s \in S$ is called the *k -th angular neighbor* of $\overrightarrow{pq}$ whenever $\text{Cone}(spq) \cap R$ contains exactly $k-1$ points of S in its interior.

Fig. 1. (a) Forbidden zone of a pentagon P , (b) 11 points without a 5-hole and a 4-hole which are disjoint [13], and (c) Illustration of the proof of Observation 1.

3 Proof of Theorem 1

Urabe and Hosono [13] constructed a set of 11 points not containing an 4-hole and a disjoint 5-hole, which is shown in Figure 1(b). This implies that $H(4, 5) \geq 12$. Therefore, for proving the theorem it suffices to show that $H(4, 5) \leq 12$.

Let S be a set of 12 points in general position in the plane. We say S is *admissible* whenever S contains a 4-hole and 5-hole which are disjoint.

First, consider that S does not contain a 6-hole. Then Theorem 2 implies that S must contain a 5-hole and a disjoint 4-hole. Therefore, assume that S contains a 6-hole.

We now have the following observation:

Observation 1 *If S contains a 7-hole, then S is admissible.*

Proof. Let $H := s_1 s_2 s_3 s_4 s_5 s_6 s_7$ be a 7-hole in S . For $i \in \{1, 2, \dots, 7\}$, let Q_i denote the region $\text{Cone}(s_{i+3} s_i s_{i+4}) \setminus \mathcal{I}(s_{i+3} s_i s_{i+4})$ (Figure 1(c)), with indices taken modulo 7. If $|Q_1 \cap S| = 0$, then by the pigeon-hole principle either $|\mathcal{H}(s_1 s_4, s_2) \cap S| \geq 5$ or $|\mathcal{H}(s_1 s_5, s_2) \cap S| \geq 5$. Without loss of generality, let $|\mathcal{H}(s_1 s_4, s_2) \cap S| \geq 5$. Then $\mathcal{H}(s_1 s_4, s_2) \cap S$ contains a 4-hole, since $H(4) = 5$. This 4-hole is disjoint from the 5-hole $s_1 s_4 s_5 s_6 s_7$. Therefore, whenever $|Q_i \cap S| = 0$ for some $i \in \{1, 2, \dots, 7\}$, then S is admissible. However, $|Q_i \cap S| \geq 1$ for all $i \in \{1, 2, \dots, 7\}$ implies, $\sum_{i=1}^7 |Q_i \cap S| \geq 7 > 5 = |S| - |\mathcal{V}(H)|$, which is a contradiction. $\square$

Fig. 2. The subdivision of the exterior of the 6-hole $s_1 s_2 s_3 s_4 s_5 s_6$.

Let $B := s_1s_2s_3s_4s_5s_6$ be a 6-hole in S . In light of Observation 1, it can be assumed that the forbidden zone of B is empty in S , that is, B cannot be extended to a 7-hole. Hereafter, while indexing the points of $\mathcal{V}(B)$, we identify the indices modulo 6.

We begin with a simple observation:

Observation 2 *If for some $s_i \in \mathcal{V}(B)$, $|\overline{\mathcal{H}}(s_i s_{i+1}, s_{i+2}) \cap S| \geq 4$, then S is admissible.*

Proof. If $|\overline{\mathcal{H}}(s_i s_{i+1}, s_{i+2}) \cap S| \geq 4$, then $(\overline{\mathcal{H}}(s_i s_{i+1}, s_{i+2}) \cap S) \cup \{s_i\}$ contains a 4-hole, as $H(4) = 5$. This 4-hole is disjoint from the 5-hole formed by $\mathcal{V}(B) \setminus \{s_i\}$. Hence S is admissible. $\square$

Consider the subdivision of the exterior of the hexagon B into regions R_i and $R_i R_j$, as shown in Figure 2. The regions of the type R_i are disjoint from each other, but the regions of the type $R_i R_j$ may overlap with each other but are disjoint from regions of the type R_i . Observe that in Figure 2, the deeply shaded region R is the intersection of the regions $R_2 R_3$ and $R_1 R_{12}$. $|R_i|$ or $|R_i R_j|$ denotes the number of points of S in R_i or $R_i R_j$, respectively. Also, let s, s' be two points on the extended line $s_1 s_4$ as shown in Figure 2.

Fig. 3. Illustration of the proof of (a) Observation 3 and (b) Observation 4.

Note, Observation 2 implies that for all $s_i \in \mathcal{V}(B)$, $|\overline{\mathcal{H}}(s_i s_{i+1}, s_{i+2}) \cap S| \leq 3$. In particular, $|R_i| \leq 3$ for all $i \in \{1, 2, \dots, 12\}$. Now, we have the following observations:

Observation 3 *S is admissible, if $|R_i| = 3$ for some $i \in \{1, 2, \dots, 12\}$.*

Proof. Without loss of generality assume $|R_2| = 3$. Let $s_\alpha \in S$ be the nearest angular neighbor of $\overrightarrow{s_2 s_1}$ in R_2 . Observation 2 implies that S is admissible unless $|(\mathcal{H}(s_1 s_2, s_\alpha) \setminus R_2) \cap S| = |(\mathcal{H}(s_1 s_6, s_\alpha) \setminus R_2) \cap S| = 0$.

Case 1: *The forbidden zone of the 5-hole $s_\alpha s_2 s_3 s_4 s_1$ is empty in $R_2 \cap S$. Let s_β and s_γ be the other two points in $R_2 \cap S$ such that s_β is the nearest angular neighbor of $\overrightarrow{s_2 s_\alpha}$*

in $R_2 \cap S$. If $s_\gamma \in \mathcal{H}(s_\alpha s_\beta, s_1)$, then $s_1 s_\alpha s_\beta s_\gamma$ is a 4-hole which is disjoint from the 5-hole $s_2 s_3 s_4 s_5 s_6$ (see Figure 3(a)). Otherwise, $s_\gamma \in \mathcal{H}(s_\alpha s_\beta, s_2)$, and $s_2 s_\alpha s_\beta s_\gamma$ is a 4-hole disjoint from 5-hole $s_1 s_3 s_4 s_5 s_6$.

Case 2: There exists $s_\beta \in R_2 \cap S$ such that $s_\alpha s_\beta s_2 s_3 s_4 s_1$ is a 6-hole. If $|Cone(ss_1 s_3) \cap S| \geq 5$, $Cone(ss_1 s_3) \cap S$ contains a 4-hole, since $H(4) = 5$. This 4-hole and the 5-hole $s_1 s_3 s_4 s_5 s_6$ are disjoint (see Figure 3(a)). Otherwise, $|Cone(ss_1 s_3) \cap S| \leq 4$, and so $|Cone(s_6 s_1 s_3) \cap S| \geq 5$. This implies that the 4-hole contained $Cone(s_6 s_1 s_3) \cap S$ is disjoint from 5-hole $s_\alpha s_\beta s_2 s_3 s_1$. $\square$

Observation 4 S is admissible, if $|R_i| = 2$ for some $i \in \{1, 2, \dots, 12\}$.

Proof. Without loss of generality assume $|R_2| = 2$. Let $R_2 \cap S = \{s_\alpha, s_\beta\}$, where s_α is the nearest angular neighbor of $\overrightarrow{s_2 s_1}$ in R_2 . There are two cases:

Case 1: s_α lies inside the triangle $s_1 s_2 s_\beta$. Let s^*, t^* be as shown in Figure 3(b). If there exists a point $s_\gamma \in S \setminus \{s_\alpha, s_\beta\}$ in the halfplane $\mathcal{H}(s_1 s_2, s_\alpha)$ or $\mathcal{H}(s_1 s_6, s_\alpha)$, then either $s_1 s_\alpha s_\beta s_\gamma$ or $s_2 s_\alpha s_\beta s_\gamma$ is a 4-hole, and the admissibility of S is immediate. Hence, assume that s_α and s_β are the only points of S in these two halfplanes. Observe that $|Cone(ss_1 s_3) \cap S| \geq 3$. Since $H(4) = 5$, $Cone(ss_1 s_3) \cap S$ contains a 4-hole whenever $|Cone(ss_1 s_3) \cap S| \geq 5$. This 4-hole is then disjoint from 5-hole $s_1 s_3 s_4 s_5 s_6$. Therefore, assume that $3 \leq |Cone(ss_1 s_3) \cap S| \leq 4$.

Case 1.1: $|Cone(ss_1 s_3) \cap S| = 4$. This implies that $|Cone(s_2 s_3 t^*) \cap S| = 1$. Suppose, $Cone(s_2 s_3 t^*) \cap S = \{s_\gamma\}$. If $s_\gamma \in Cone(s_\beta s_\alpha s_2)$, then $s_\beta s_\alpha s_2 s_\gamma$ is a 4-hole which is disjoint from the 5-hole $s_1 s_3 s_4 s_5 s_6$. Otherwise, $s_2 s_\alpha s_1 s_3 s_\gamma$ is a 5-hole which is disjoint from the 4-hole contained in $Cone(s_3 s_1 s_6) \cap S$, since $|Cone(s_3 s_1 s_6) \cap S| \geq 5$.

Case 1.2: $|Cone(ss_1 s_3) \cap S| = 3$. If $|\mathcal{H}(s_3 s_5, s_4) \cap S| \geq 5$, $H(4) = 5$ immediately implies the admissibility of S . Hence, assume that $|\mathcal{H}(s_3 s_5, s_4) \cap S| = \lambda \leq 4$. Now, since $\overline{\mathcal{H}}(s_1 s_2, s_4) \cap S = \overline{\mathcal{H}}(s_1 s_6, s_4) \cap S = \{s_\alpha, s_\beta\}$ and $Cone(s_2 s_3 t^*) \cap S$ is empty, we have $|\overline{\mathcal{H}}(s_3 s_5, s_4) \cap (R_{10} \cap S)| = 5 - \lambda$. Then there exists $s_\gamma \in \overline{\mathcal{H}}(s_3 s_5, s_4) \cap (R_{10} \cap S)$ such that $s_1 s_2 s_3 s_\gamma s_6$ is a 5-hole which is disjoint from the 4-hole contained in $\mathcal{H}(s_3 s_\gamma, s_4)$.

Case 2: $s_1, s_2, s_\beta, s_\alpha$ are in convex position. Then $s_1 s_3 s_4 s_5 s_6$ and $s_1 s_\alpha s_\beta s_2 s_3$ are two 5-holes sharing the edge $s_1 s_3$. Now, since $|S \setminus \{s_1, s_3\}| = 10$, by the pigeonhole principle either $|\mathcal{H}(s_1 s_3, s_2) \cap S| \geq 5$ or $|\overline{\mathcal{H}}(s_1 s_3, s_2) \cap S| \geq 5$. Therefore, the 4-hole contained in $\mathcal{H}(s_1 s_3, s_2) \cap S$ or $\overline{\mathcal{H}}(s_1 s_3, s_2) \cap S$ is disjoint from the 5-hole $s_1 s_3 s_4 s_5 s_6$ or $s_1 s_\alpha s_\beta s_2 s_3$, respectively. $\square$

Equipped with these three observations we proceed with the proof of Theorem 1. For every point $s_i \in \mathcal{V}(B)$, the diagonal $d := s_i s_{i+3}$ is called a *dividing diagonal* of B . A dividing diagonal d of B is called an (a, b) -*splitter* of S , where $a \leq b$ are integers, if either $|\mathcal{H}(s_i s_{i+3}, s_{i+1}) \cap S \setminus \mathcal{V}(B)| = a$ and $|\overline{\mathcal{H}}(s_i s_{i+3}, s_{i+1}) \cap S \setminus \mathcal{V}(B)| = b$, or $|\mathcal{H}(s_i s_{i+3}, s_{i+1}) \cap S \setminus \mathcal{V}(B)| = a$ and $|\mathcal{H}(s_i s_{i+3}, s_{i+1}) \cap S \setminus \mathcal{V}(B)| = b$.

From Observations 3 and 4, we have $|R_i| \leq 1$ for all $i \in \{1, 2, \dots, 12\}$. Now, if some dividing diagonal of B , say $s_1 s_4$, is a $(0, 6)$ -splitter of S with $|\mathcal{H}(s_1 s_4, s_2) \cap S \setminus \mathcal{V}(B)| = 6$, then from Observation 2, $|\overline{\mathcal{H}}(s_2 s_3, s_1) \cap S| \leq 3$, and hence $|R_2| + |R_7| \geq 3$. Then, either $|R_2| \geq 2$ or $|R_7| \geq 2$, and the admissibility of S is immediate from Observations 3 and 4. Therefore, no dividing diagonal of B is a $(0, 6)$ -splitter of S . The only cases which remain to be considered are:

Fig. 4. (a) s_1s_4 is a $(1, 5)$ -splitter of S such that $|R_2| = |R_7| = 1$, (b) s_1s_4 is a $(2, 4)$ -splitter of S such that $|R_2| = 1$.

Case 1: s_1s_4 is a $(1, 5)$ -splitter of S with $|\mathcal{H}(s_1s_4, s_2) \cap S \setminus \mathcal{V}(B)| = 5$. Observations 2, 3, and 4 imply that $|R_2| = |R_7| = 1$. Let s'', s^*, u, v be as shown in Figure 4(a). Let $s_\alpha \in R_2 \cap S$ and $s_r \in R_7 \cap S$. Now, $|\overline{\mathcal{H}}(s_2s_3, s_1) \cap S| = 3$ and Observation 2 implies that either $|\text{Cone}(us_2s'') \cap S| = 2$ or $|\text{Cone}(vs_3s^*) \cap S| = 2$. Without loss of generality assume, $|\text{Cone}(us_2s'') \cap S| = 2$. Let $s_\beta, s_\gamma \in \text{Cone}(us_2s'') \cap S$, such that s_β is the nearest angular neighbor of $\overline{s_2u}$ in $\text{Cone}(us_2s'')$. If $s_\alpha \in \mathcal{I}(s_2s_1s_\beta)$, then either $s_2s_\alpha s_\beta s_\gamma$ or $s_1s_\alpha s_\beta s_\gamma$ is a 4-hole which is disjoint from the 5-hole $s_1s_3s_4s_5s_6$ or $s_2s_3s_4s_5s_6$, respectively. Otherwise, $s_\alpha \notin \mathcal{I}(s_2s_1s_\beta)$ and $s_1s_\alpha s_\beta s_2$ is a 4-hole. It can be assumed that the forbidden zone of $s_1s_\alpha s_\beta s_2$ is empty in $\overline{\mathcal{H}}(s_2s_3, s_1) \cap S$. Then, $s_\beta \in \mathcal{I}(s_2s_\alpha s_\gamma)$ (see Figure 4(a)). If $s_\delta \in S$ is the remaining point in $\mathcal{H}(s_2s_3, s_\gamma) \cap S$, then either $s_\alpha s_\beta s_\gamma s_\delta$ or $s_2s_\beta s_\gamma s_\delta$ is a 4-hole disjoint from the 5-hole $s_1s_3s_4s_5s_6$.

Case 2: s_1s_4 is a $(2, 4)$ -splitter of S with $|\mathcal{H}(s_1s_4, s_2) \cap S \setminus \mathcal{V}(B)| = 4$. Without loss of generality, suppose $|R_2| = 1$. Let $s_\alpha \in R_2 \cap S$. Refer to Figure 4(b). If $|R_7| \neq 0$, there exists $s_\beta \in R_7 \cap S$ such that $s_\alpha s_2 s_3 s_\beta s_4$ is a 5-hole which is disjoint from the 4-hole contained in $(\mathcal{H}(s_1s_4, s_5) \cap S) \cup \{s_1\}$. Therefore, assume that $|R_7| = 0$. Similarly, it can be shown that S is admissible unless $|R_1| = |R_8| = 0$. So, $|\overline{\mathcal{H}}(s_5s_6, s_1) \cap S| = 2$. Let $\overline{\mathcal{H}}(s_5s_6, s_1) \cap S = \{s_q, s_r\}$. If s_5, s_6, s_q, s_r are in convex position, then this 4-hole is disjoint from the 5-hole $s_\alpha s_2 s_3 s_4 s_1$. Therefore, assume that $s_q \in \mathcal{I}(s_5s_6s_r)$. This implies that either $s_q, s_r \in \overline{\mathcal{H}}(s_1s_6, s_4)$ or $s_q, s_r \in \overline{\mathcal{H}}(s_4s_5, s_1)$. If $s_q, s_r \in \overline{\mathcal{H}}(s_1s_6, s_4)$, then $s_\alpha s_6 s_q s_r$ is a 4-hole which is disjoint from the 5-hole $s_1s_2s_3s_4s_5$. Therefore, let $s_q, s_r \in \overline{\mathcal{H}}(s_4s_5, s_1) \cap S$ (see Figure 4(b)). Again, S is admissible unless R_6R_7 is empty in S . If $|\text{Cone}(ss_1s_2) \cap \overline{\mathcal{H}}(s_2s_3, s_1) \cap S| \geq 3$, then $|\overline{\mathcal{H}}(s_1s_2, s_3) \cap S| \geq 4$, and the admissibility of S follows from Observation 2. Again, if $|\text{Cone}(ss_1s_2) \cap \overline{\mathcal{H}}(s_2s_3, s_1) \cap S| \leq 1$, then $|R_5| + |R_6| \geq 2$. From Observations 3 and 4, it suffices to consider $|R_5| = |R_6| = 1$. Then, $\{s_1, s_2, s_3, s_6\} \cup (R_5 \cap S)$ forms a 5-hole which is disjoint from the 4-hole $s_r s_q s_5 s_4$. Therefore, assume that $|\text{Cone}(ss_1s_2) \cap \overline{\mathcal{H}}(s_2s_3, s_1) \cap S| = 2$. Let $s_\beta, s_\gamma \in \text{Cone}(us_2s'') \cap S$, such that s_β is the nearest angular neighbor of $\overline{s_2u}$ in $\text{Cone}(us_2s'')$. Refer to Figure 4(a). If $s_\alpha \in \mathcal{I}(s_2s_1s_\beta)$, then either $s_2s_\alpha s_\beta s_\gamma$ or $s_1s_\alpha s_\beta s_\gamma$ is a 4-hole which is disjoint from the 5-hole $s_1s_3s_4s_5s_6$ or $s_2s_3s_4s_5s_6$, respectively. Otherwise, $s_\alpha \notin \mathcal{I}(s_2s_1s_\beta)$ and $s_1s_\alpha s_\beta s_2$ is a

4-hole. It can be assumed that the forbidden zone of $s_1 s_\alpha s_\beta s_2$ is empty in $\overline{\mathcal{H}}(s_2 s_3, s_1) \cap S$. Then, $s_\beta \in \mathcal{I}(s_2 s_\alpha s_\gamma)$. If $s_\delta \in S$ is the remaining point in $\mathcal{H}(s_2 s_3, s_\gamma) \cap S$, then either $s_\alpha s_\beta s_\gamma s_\delta$ or $s_2 s_\beta s_\gamma s_\delta$ is a 4-hole disjoint from the 5-hole $s_1 s_3 s_4 s_5 s_6$.

Fig. 5. All the dividing diagonals of B are $(3,3)$ -splitters of S : (a) Case 3.1, (b) Case 3.2

Case 3: All the dividing diagonals of the hexagon B are $(3,3)$ -splitters of S . Suppose $R_1 \cap S$ is non-empty. Then, there exists $s_\alpha \in R_1 \cap S$ such that $s_\alpha s_1 s_4 s_5 s_6$ is a 5-hole. This 5-hole is disjoint from the 4-hole contained in $\mathcal{H}(s_1 s_4, s_2) \cap S$, since $H(4) = 5$. Therefore, it can be assumed that $|R_i| = 0$ for $i \in \{1, 2, \dots, 12\}$. Observation 2 and the fact that $|S \setminus \mathcal{V}(B)| = 6$ now implies that for any point $s_i \in \mathcal{V}(B)$, $|\overline{\mathcal{H}}(s_i s_{i+1}, s_{i+2}) \cap S| = 3$. Therefore, regions in the exterior of the hexagon B , where the regions of the type $R_i R_j$ intersect must be empty in S . Now, we consider the following two cases:

Case 3.1: $|R_i R_j| \leq 2$ for all pair of indices i, j . Observe that either $|R_2 R_3| + |R_4 R_5| \geq 2$ or $|R_4 R_5| + |R_6 R_7| \geq 2$. Without loss of generality assume that $|R_2 R_3| + |R_4 R_5| \geq 2$. To begin with, suppose that $|R_2 R_3| + |R_4 R_5| = 2$, with $(R_2 R_3 \cup R_4 R_5) \cap S = \{s_\alpha, s_\beta\}$ and $R_6 R_7 \cap S = \{s_\gamma\}$. If the four points $s_2, s_\alpha, s_\beta, s_\gamma$ form a convex quadrilateral, S is clearly admissible. Otherwise, let $s_\beta \in \mathcal{I}(s_2 s_\alpha s_\gamma)$ and $R_1 R_{12} \cap S = \{s_\delta\}$. Depending on the position of the point s_δ , either $s_2 s_\beta s_\alpha s_\delta$ or $s_\delta s_\alpha s_\beta s_\gamma$ is a 4-hole which is disjoint from the 5-hole $s_1 s_3 s_4 s_5 s_6$ (see Figure 5(a)). Now, let $|R_2 R_3| + |R_4 R_5| = 3$ and without loss of generality, assume $|R_2 R_3| = 2$ and $|R_4 R_5| = 1$. Then $|R_1 R_{12}| = 0$ and $|R_{10} R_{11}| = 1$, since all three dividing diagonals of B are $(3,3)$ -splitters of S . From symmetry, it is the same case as before.

Case 3.2: $|R_i R_j| = 3$ some pair of indices i, j . Without loss of generality assume that $|R_1 R_{12}| = 3$. This implies that $|R_6 R_7| = 3$. Let s_0 be as shown in Figure 5(b) and $s_\alpha, s_\beta, s_\gamma$ be the first, second, and third angular neighbors of $\overline{s_3 s_0}$ in $R_6 R_7$, respectively. If $s_\alpha \in \mathcal{I}(s_3 s_4 s_\beta)$ the admissibility of S follows from the fact that $s_3 s_\alpha s_\beta s_\gamma$ or $s_4 s_\alpha s_\beta s_\gamma$ is a 4-hole which is disjoint from the 5-hole $s_1 s_2 s_4 s_5 s_6$ or $s_1 s_2 s_3 s_5 s_6$, respectively. Otherwise, $s_\alpha \notin \mathcal{I}(s_3 s_4 s_\beta)$, and similarly $s_\gamma \notin \mathcal{I}(s_2 s_3 s_\beta)$.

Then, either $s_1s_3s_\beta s_\alpha s_4$ or $s_1s_3s_\beta s_\gamma s_2$ is a 5-hole which is disjoint from the 4-hole contained in $(R_1R_{12} \cap S) \cup \{s_5, s_6\}$. $\square$

4 Remarks and Conclusions

In this paper we proved that $H(4, 5) = 12$, that is, every set of 12 points in plane in general position contains a 4-hole and a disjoint 5-hole, thus improving a result of Hosono and Urabe [12]. The proof uses a Ramsey type result for 11 points proposed by Aichholzer et al. [1].

The most important case that remains to be settled is that of $H(5, 5)$. Urabe and Hosono [13] proved that $16 \leq H(5, 5) \leq 20$, and later improved the lower bound to 17 [12]. There is still a substantial gap between the upper and lower bounds of $H(5, 5)$. We believe that a new Ramsey-type result similar to Theorem 2 might be useful in obtaining better bounds on $H(5, 5)$.

However, we are still far from establishing non-trivial bounds on $H(6, l)$, for $0 \leq l \leq 6$, since the exact value of $H(6) = H(6, 0)$ is still unknown. The best known bounds are, $30 \leq H(6) \leq ES(9) \leq 1717$. The lower bound is due to Overmars [18] and the upper bound due to Gerken [9].

Acknowledgement

We are grateful to the anonymous referee for the valuable comments, which has helped us in improving the quality of the paper.

References

1. O. Aichholzer, C. Huemer, S. Kappes, B. Speckmann, C. D. Tóth, Decompositions, partitions, and coverings with convex polygons and pseudo-triangles, *Graphs and Combinatorics*, Vol. 23, 481–507, 2007.
2. O. Aichholzer, F. Aurenhammer, H. Krasser, Enumerating order types for small point sets with applications, *Order*, Vol. 19, 265–281, 2002.
3. O. Aichholzer, H. Krasser, Abstract order type extensions and new results on the rectilinear crossing number, *Proc. 21st Symposium on Computational Geometry*, ACM Press, 91–98, 2005.
4. I. Bárány, G. Károlyi, Problems and results around the Erdős-Szekeres convex polygon theorem, *JCD CG*, LNCS 2098, 91–105, 2000.
5. Ren Ding, Kiyoshi Hosono, Masatsugu Urabe, Changqing Xu, Partitioning a planar point set into empty convex polygons, *JCD CG*, LNCS 2866, 129–134, 2002.
6. P. Erdős, Some more problems on elementary geometry, *Australian Mathematical Society Gazette*, Vol. 5, 52–54, 1978.
7. P. Erdős, G. Szekeres, A combinatorial problem in geometry, *Compositio Mathematica*, Vol. 2, 463–470, 1935.
8. P. Erdős, G. Szekeres, On some extremum problems in elementary geometry, *Ann. Univ. Sci. Budapest, Eötvös, Sect. Math.* 3/4, 53–62, 1960–61.
9. T. Gerken, Empty convex hexagons in planar point sets, *Discrete and Computational Geometry*, Vol. 39, 239–272, 2008.
10. H. Harborth, Konvexe Funfecke in ebenen Punktmengen, *Elemente der Mathematik*, Vol. 33(5), 116–118, 1978.
11. J.D. Horton, Sets with no empty convex 7-gons, *Canadian Mathematical Bulletin*, Vol. 26, 482 – 484, 1983.
12. K. Hosono, M. Urabe, A minimal planar point set with specified disjoint empty convex subsets, *Kyoto CGGT*, LNCS 4535, 90–100, 2008.
13. K. Hosono, M. Urabe, On the minimum size of a point set containing two non-intersecting empty convex polygons, *JCD CG*, LNCS 3742, 117–122, 2004.

14. K. Hosono, M. Urabe, On the number of disjoint convex quadrilaterals for a planar point set, *Computational Geometry: Theory and Applications*, Vol. 20, 97 – 104, 2001.
15. J. D. Kalbfleisch, J. G. Kalbfleisch, R. G. Stanton, A combinatorial problem on convex regions, *Proc. Louisiana Conf. Combinatorics, Graph Theory and Computing, Louisiana State Univ., Baton Rouge, La., Congr. Numer.*, Vol. 1, 180–188, 1970.
16. W. Morris, V. Soltan, The Erdős-Szekeres problem on points in convex position - A survey, *Bulletin of the American Mathematical Society*, Vol. 37(4), 437 – 458, 2000.
17. C. M. Nicolás, The empty hexagon theorem, *Discrete and Computational Geometry*, Vol. 38, 389–397, 2007.
18. M. Overmars, Finding sets of points without empty convex 6-gons, *Discrete and Computational Geometry*, Vol. 29, 153– 158, 2003.
19. G. Szekeres, L. Peters, Computer solution to the 17-point Erdős-Szekeres problem, *ANZIAM Journal*, Vol. 48, 151–164, 2006.
20. G. Tóth, P. Valtr, The Erdős-Szekeres theorem: upper bounds and related results, in *J. E. Goodman, J. Pach, and E. Welzl, Combinatorial and Computational Geometry*, MSRI Publications 52, 557–568, 2005.
21. M. Urabe, On a partition into convex polygons, *Discrete Applied Mathematics*, Vol. 64, 179–191, 1996.
22. P. Valtr, On empty hexagons, in *J. E. Goodman, J. Pach, R. Pollack, Surveys on Discrete and Computational Geometry, Twenty Years Later*, AMS, 433–441, 2008.
23. L. Wu, R. Ding, On the number of empty convex quadrilaterals of a finite set in the plane, *Applied Mathematics Letters* Vol. 21, 966–973, 2008.
24. L. Wu, R. Ding, Reconfirmation of two results on disjoint empty convex polygons, *CJCDGCGT*, LNCS 4381, 216–220, 2005.
25. C. Xu, R. Ding, On the empty convex partition of a finite set in the plane, *Chinese Annals of Mathematics*, Vol. 23 (4), 487–494, 2002.