



Publicly Accessible Penn Dissertations

1-1-2016

On Decomposition of the Product of Demazure atoms and Demazure Characters

Ying Anna Pun

University of Pennsylvania, annapunying@gmail.com

Follow this and additional works at: <http://repository.upenn.edu/edissertations>

 Part of the [Mathematics Commons](#)

Recommended Citation

Pun, Ying Anna, "On Decomposition of the Product of Demazure atoms and Demazure Characters" (2016). *Publicly Accessible Penn Dissertations*. 1959.

<http://repository.upenn.edu/edissertations/1959>

This paper is posted at Scholarly Commons. <http://repository.upenn.edu/edissertations/1959>

For more information, please contact libraryrepository@pobox.upenn.edu.

On Decomposition of the Product of Demazure atoms and Demazure Characters

Abstract

This thesis studies the properties of Demazure atoms and characters using linear operators and also tableaux-combinatorics. It proves the atom-positivity property of the product of a dominating monomial and an atom, which was an open problem. Furthermore, it provides a combinatorial proof to the key-positivity property of the product of a dominating monomial and a key using skyline fillings, an algebraic proof to the key-positivity property of the product of a Schur function and a key using linear operator and verifies the first open case for the conjecture of key-positivity of the product of two keys using linear operators and polytopes.

Degree Type

Dissertation

Degree Name

Doctor of Philosophy (PhD)

Graduate Group

Mathematics

First Advisor

James Haglund

Keywords

decomposition of products, Demazure atoms, Demazure characters, key polynomials, skyline fillings, SSAF

Subject Categories

Mathematics

ON DECOMPOSITION OF THE PRODUCT OF DEMAZURE ATOMS AND
DEMAZURE CHARACTERS

Anna, Ying Pun

A DISSERTATION

in

Mathematics

Presented to the Faculties of the University of Pennsylvania

in

Partial Fulfillment of the Requirements for the

Degree of Doctor of Philosophy

2016

Supervisor of Dissertation

James Haglund, Professor of Mathematics

Graduate Group Chairperson

David Harbater, Professor of Mathematics

Dissertation Committee:

James Haglund, Professor of Mathematics

Greta Panova, Assistant Professor of Mathematics

Jonathan Block, Professor of Mathematics

Acknowledgments

I would like to express my genuine gratitude to my advisor Prof. James Haglund for his guidance and encouragement throughout the period of my graduate study. I would also like to thank every member in the CAGE seminar group, especially Prof. Jennifer Morse, Prof. Linda Chen, Prof. Elizabeth Milicevic and Per Alexandersson for their ideas and support. Thanks Paula, Janet, Monica, Robin and Reshma in the Math Department office for their great help.

Thanks to my classmates and friends Max, Soum, Hua, Jin, Spencer, Martin, Siggy, Zachary and all others that I met in Hong Kong, especially Kei Yuen Chan and Ming Ho Ng for their wonderful friendship. Thanks to Brett for always helping me to take care of my dogs when I was away for conferences. I am also indebted to James Sundstrom for helping me through the oral examination and for his relentless support.

Special thanks to Prof. M.K. Siu, Dr. P.P.W. Wong and Dr. G. Han from the HKU for their advices throughout these years, and to Prof. Richard Stanley for ideas and advices in Math and comics and let me have a chance to relax and move on whenever I struggled with my study.

Fianlly my greatest thanks to my family: My adorable dogs Lan Jai (靚仔) and Lan Lui(靚女) for accompanying me during these 5 years in the US and giving me great emotional support, and my mother, who is the greatest person I have ever met. She has always been there to encourage me to do my best, and has shown lots of love and care when I am under stress. It would not be possible for me to finish my graduate study without her.

ABSTRACT
ON DECOMPOSITION OF THE PRODUCT OF DEMAZURE ATOMS AND DEMAZURE
CHARACTERS

Anna, Ying Pun

James Haglund

This thesis studies the properties of Demazure atoms and characters using linear operators and also tableaux-combinatorics. It proves the atom-positivity property of the product of a dominating monomial and an atom, which was an open problem. Furthermore, it provides a combinatorial proof to the key-positivity property of the product of a dominating monomial and a key using skyline fillings, an algebraic proof to the key-positivity property of the product of a Schur function and a key using linear operator and verifies the first open case for the conjecture of key-positivity of the product of two keys using linear operators and polytopes.

Contents

Introduction	1
1 Symmetric group S_n	4
2 Demazure atoms and characters	12
2.1 Linear operators	12
2.2 Semi-Standard Augmented filling	25
2.3 Demazure atoms and Demazure characters	28
3 Decomposition of products of Demazure atoms and characters	31
3.1 Convert a column word to a row word	33
3.2 Convert a Column Recording Tableau to a Row Recording Tableau	51
3.3 Decomposition of the product of a dominating monomial and an atom into a positive sum of atoms	70
3.4 Decomposition of the product of a dominating monomial and a key into a positive sum of keys	74
3.5 Decomposition of the product of a Schur function and a Demazure character	76
4 Atom positivity of the product of two key polynomials whose basements have length at most 3	79

4.1	Polytopes	80
4.1.1	Demazure characters and polytopes	81
4.1.2	Demazure atoms and polytopes	91
4.2	Products of two Demazure characters	102
4.2.1	$\pi_1(x_1^m x_2^n) \times \pi_2(x_1^k x_2^l)$	102
4.2.2	$\pi_1(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l)$	107
4.2.3	$\pi_2(x_1^m x_2^n) \times \pi_{12}(x_1^k x_2^l)$	112
4.2.4	$\pi_{12}(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l)$	112
4.2.5	All other cases in the table	113
	Appendix	114
	A Bijection between LRS and LRK	115

Introduction

Macdonald [9] defined a family of non-symmetric polynomials, called non-symmetric Macdonald polynomials,

$$\{E_\gamma(x_1, \dots, x_n; q, t) \mid \gamma \text{ is a weak composition with } n \text{ parts, } n \in \mathbb{N}\}$$

which shares many properties with the family of symmetric Macdonald polynomials [8]

$$\{P_\lambda(x_1, \dots, x_n; q, t) \mid \lambda \text{ is a partition with } n \text{ parts, } n \in \mathbb{N}\}.$$

Haglund, Haiman and Loehr [2] obtained a combinatorial formula for $E_\gamma(X; q, t)$ where $X = (x_1, \dots, x_n)$, using fillings of augmented diagram of shape γ , called *skyline fillings*, satisfying certain constraints.

Marshall[10] studied the family of non-symmetric Macdonald polynomials using another notation $\hat{E}_\gamma(x_1, \dots, x_n; q, t) := E_{\bar{\gamma}}(x_n, \dots, x_1; \frac{1}{q}, \frac{1}{t})$. In particular, by setting $q = t = 0$ in \hat{E}_γ , one can obtain Demazure atoms (first studied by Lascoux and Schützenberger[6]) $\mathcal{A}_\gamma = \hat{E}_\gamma(x_1, \dots, x_n; 0, 0) = E_{\bar{\gamma}}(x_n, \dots, x_1; \infty, \infty)$. Similarly, one can obtain Demazure characters (key polynomials) by setting $q = t = 0$ in E_γ , *i.e.*, $\kappa_\gamma = E_\gamma(x_1, \dots, x_n; 0, 0) = \hat{E}_{\bar{\gamma}}(x_n, \dots, x_1; \infty, \infty)$. The set of all Demazure atoms forms a basis for the polynomial ring, as does the set of all key polynomials.

Haglund, Luoto, Mason, Remmel and van Willigenburg [3], [4] further studied the combinatorial formulas for Demazure atoms and Demazure characters given by the skyline fillings and obtained results which generalized those for Schur functions like the Pieri Rule, the Robinson-Schensted-Knuth (RSK) algorithm, and the Littlewood-Richardson (LR) rule.

It is a classical result in Algebraic Geometry that the product of two Schubert polynomials can be written as a positive sum of Schubert polynomials. A representation theoretic proof is also given recently by using Kraskiewicz-Pragacz modules [13]. However a combinatorial proof of the positivity property of Schubert polynomials has long been open.

Since every Schubert polynomial is a positive sum of key polynomials [7], the product of two Schubert polynomials is a positive sum of product of two key polynomials. This suggests one to study the product of two key polynomials. It is known that the product of two key polynomials is not key-positive in general. However, it is still a conjecture that whether the product is atom-positive. This provides a possible approach to a combinatorial proof of the positivity property of Schubert polynomials by trying to recombine the atoms into keys and hence into Schubert polynomials.

Also, since key polynomials are positive sum of atoms [6], one can study the atom-positivity properties of the products between atoms and keys or even atoms and atoms to try to prove the conjecture by recombining the atoms back to keys. In this thesis, we prove that the product of a dominating monomial and an atom is always atom positive and that the product of a dominating monomial and a key is always key positive (and hence atom-positive) by using the insertions introduced in [11] and [3].

In Chapter 1, we will give a brief summary on notations and some results in symmetric groups. We will introduce Demazure atoms and Demazure keys in Chapter 2 by first defining them using linear operators and then define them using semi-standard augmented fillings. We will then study some properties of atoms and characters using both definitions. We also study some properties among the linear operators and obtain certain useful identities for the proofs in later Chapters.

In Chapter 3, we will set up the tools, namely, words and recording tableaux, that we need to prove the main results of this thesis in the first 2 sections in the chapter and give the proof in Section 3.3. We then give alternative proofs to known results, namely, the key-positivity of the product of a dominating monomial and a key in Section 3.4 using results in Section 6 of [3] and the key-positivity of the product of a Schur function and a key in Section 3.5.

We will check the first open case of the conjecture of the key-positivity of the product of two key polynomials in Chapter 4. We first introduce a geometric interpretation of Demazure atoms and characters in Section 4.1. We then verify the key-positivity of the product of every pair of keys in this open case in Section 4.2.

We will give a brief summary of the materials from [3] that we use in Section 3.4 in the Appendix.

Chapter 1

Symmetric group S_n

This chapter gives a brief summary of the terminologies, notations, lemmas and theorems that will be used in later chapters.

Let $[n] = \{1, \dots, n\}$ be the set of all positive integers not greater than n . Let S_n be the group of all permutations on $[n]$, i.e. $S_n = \{\sigma : [n] \rightarrow [n] \mid \sigma \text{ is bijective}\}$ with identity id such that $id(j) = j$ for all $j \in [n]$, and the group product is defined as the composition of functions, that is, for all $\sigma_1, \sigma_2 \in S_n$, $\sigma_1\sigma_2(j) = \sigma_1(\sigma_2(j))$ for all $j \in [n]$.

Definition 1.1. Let n be a positive integer and $1 \leq k \leq n$. A cycle of length k , denoted as (a_1, a_2, \dots, a_k) , where a_1, a_2, \dots, a_k are k distinct integers in $[n]$, is a permutation $\sigma \in S_n$ such that

$$\begin{cases} \sigma(a_i) = a_{i+1} & \text{for } 1 \leq i < k \\ \sigma(a_k) = a_1 \\ \sigma(j) = j & \text{if } j \neq a_i \text{ for any } 1 \leq j \leq k \end{cases}$$

Proof. For any k -cycle, we have $(a_1, a_2, \dots, a_k) = (a_1, a_k)(a_1, a_{k-1}) \cdots (a_1, a_2)$. Also, for $3 \leq j \leq k$, $(a_1, a_j) = (a_{j-1}, a_j)(a_1, a_{j-1})(a_{j-1}, a_j)$. Hence every cycle can be written as a product of simple transpositions.

As a result, by Theorem 1.2, every permutation is a product of simple transpositions and thus $S_n = \langle s_1, s_2, \dots, s_{n-1} \rangle$. □

There are several ways to represent a permutation $\sigma \in S_n$:

1. Two-line notation: $\sigma := \begin{pmatrix} 1 & 2 & \cdots & n \\ \sigma(1) & \sigma(2) & \cdots & \sigma(n) \end{pmatrix}$

2. One-line notation: $\sigma = \sigma(1), \sigma(2), \sigma(3), \dots, \sigma(n)$

3. Product of disjoint cycles: This follows by Theorem 1.2.

4. Product of simple transpositions: This follows by Theorem 1.3.

Example 2. Consider the permutation σ in Example 1, we can write it as:

1. Two-line notation: $\sigma := \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 5 & 2 & 4 & 3 \end{pmatrix}$

2. One-line notation: $\sigma = 1, 5, 2, 4, 3$

3. Product of disjoint cycles: $\sigma = (1)(4)(2, 5, 3)$.

4. Product of simple transpositions: $\sigma = (3, 4)(4, 5)(3, 4)(2, 3) = s_3 s_4 s_3 s_2$.

From now on, we will use one-line notation to represent a permutation, i.e.

$$\sigma = \sigma(1), \sigma(2), \sigma(3), \dots, \sigma(n)$$

unless stated otherwise.

Note that applying a transposition s_i on the left of a permutation σ means interchanging i and $i+1$ in the one-line notation of σ while applying s_i on the right interchanges entries $\sigma(i)$ and $\sigma(i+1)$ in the one line notation of σ .

By Theorem 1.3, every permutation σ can be written as a product of simple transpositions. Hence we can find a decomposition with the shortest length (that is, with the smallest number of transpositions). For $\sigma \neq id$, we call such a decomposition a reduced decomposition of σ .

Definition 1.3. Let $n \geq 2$ be an integer and $\sigma \in S_n \setminus \{id\}$. Let $\sigma = s_{i_1} s_{i_2} \cdots s_{i_k}$ be a reduced decomposition of σ . We call $i_1 i_2 \dots i_k$ a reduced word of σ .

Lemma 1.4. Every consecutive substring of a reduced word is also a reduced word.

Proof. Let $i_1 i_2 \dots i_k$ be a reduced word. If there is a consecutive substring with length $m (\geq 2)$, say $i_{r+1} \dots i_{r+m}$ which is not reduced, then $l(s_{i_{r+1}} s_{i_{r+2}} \cdots s_{i_{r+m}}) < m$ implying that there exist j_1, \dots, j_t , where $t = l(s_{i_{r+1}} s_{i_{r+2}} \cdots s_{i_{r+m}})$ such that $s_{i_{r+1}} s_{i_{r+2}} \cdots s_{i_{r+m}} = s_{j_1} s_{j_2} \cdots s_{j_t}$. Then $s_{i_1} s_{i_2} \cdots s_{i_k}$ can be written as a product of $k - m + t$ simple transpositions by replacing $s_{i_{r+1}} s_{i_{r+2}} \cdots s_{i_{r+m}}$ by $s_{j_1} s_{j_2} \cdots s_{j_t}$, which contradicts the fact that $i_1 i_2 \dots i_k$ is reduced since $k - m + t < k$. \square

Definition 1.4. Let n be any positive integer and a permutation $\sigma \in S_n \setminus \{id\}$. Define the length of σ , denoted as $l(\sigma)$, as the number of simple transpositions in a reduced decomposition. Define $l(id) = 0$.

Note that reduced decomposition of a permutation is not unique. For instance, $s_1 s_3 s_2 s_3 = s_1 s_2 s_3 s_2 = s_3 s_1 s_2 s_3$. By Tit's Theorem, any reduced word can be obtained by applying a sequence of *braid relations* (i.e. item (iii) in Proposition 1.1) on any other reduced word representing the same permutation.

Definition 1.5. Let n be a positive integer and $\sigma \in S_n$ be a permutation. The pair (i, j) is called an inversion of σ if $i < j$ and $\sigma(i) > \sigma(j)$. Denote $\text{inv}(\sigma)$ as the number of inversions of σ .

Lemma 1.5. Let $n \geq 2$ be an integer. For any permutation $\sigma \in S_n$ and a simple transposition s_i (

$1 \leq i \leq n-1$), we have

$$\text{inv}(s_i\sigma) - \text{inv}(\sigma) = \begin{cases} -1 & \text{if } (\sigma^{-1}(i+1), \sigma^{-1}(i)) \text{ is an inversion pair of } \sigma \\ 1 & \text{else} \end{cases}.$$

Proof. Let $\{x, y\} = \{\sigma^{-1}(i), \sigma^{-1}(i+1)\}$ where $x < y$.

Note that $\sigma(w) = s_i\sigma(w)$ for $w \in [n] \setminus \{x, y\}$. Also if $\sigma(w) > i$, then as $w \neq x, y$, $\sigma(w) \neq i+1$ and hence $\sigma(w) > i+1 > i$. As a result, either $s_i\sigma(w) = \sigma(w) > i+1 > i$ or $i+1 > i > \sigma(w) = s_i\sigma(w)$ for $w \neq x, y$. This means that any pair of inversion (j_1, j_2) of σ , where $\{j_1, j_2\} \neq \{x, y\}$, is also an inversion in $s_i\sigma$. Hence the only difference between $\text{inv}(s_i\sigma)$ and $\text{inv}(\sigma)$ comes from the pair (x, y) . Since (x, y) is an inversion in exactly one of σ and $s_i\sigma$, we have

$$\text{inv}(s_i\sigma) - \text{inv}(\sigma) = \begin{cases} -1 & \text{if } (\sigma^{-1}(i+1), \sigma^{-1}(i)) \text{ is an inversion pair of } \sigma \\ 1 & \text{else} \end{cases}.$$

□

Proposition 1.6. *Let $\sigma = s_{i_1}s_{i_2}\dots s_{i_k}$ (not necessarily reduced). Then $k \equiv \text{inv}(\sigma) \pmod{2}$.*

Proof. By Lemma 1.5, we have

$$\begin{aligned} \text{inv}(s_{i_1}s_{i_2}\dots s_{i_k}) &\equiv \text{inv}(s_{i_2}s_{i_3}\dots s_{i_k}) + 1 \pmod{2} \\ &\equiv \text{inv}(s_{i_3}s_{i_4}\dots s_{i_k}) + 2 \pmod{2} \\ &\vdots \\ &\equiv \text{inv}(s_{i_k}) + k - 1 \pmod{2} \\ &\equiv k \pmod{2} \end{aligned}$$

and hence $k \equiv \text{inv}(\sigma) \pmod{2}$. □

Lemma 1.7. *Let $n > 1$ be an integer and $\sigma \in S_n$ be a permutation. Let s_i be a transposition in S_n , where $1 \leq i \leq n-1$. Then $|\text{inv}(s_i\sigma) - \text{inv}(\sigma)| = 1$.*

Proof. By Theorem 1.3 and Definition 1.4, we can write σ as a product of $l(\sigma)$ simple transpositions. Hence by Lemma 1.5, we have $l(\sigma) \equiv \text{inv}(\sigma) \pmod{2}$. Similarly, $l(s_i\sigma) \equiv \text{inv}(s_i\sigma) \pmod{2}$. Therefore, $l(s_i\sigma) - l(\sigma) \equiv \text{inv}(s_i\sigma) - \text{inv}(\sigma) \equiv 1 \pmod{2}$ and we get $l(s_i\sigma) \neq l(\sigma)$.

Let $s_i\sigma = s_{i_1}s_{i_2} \cdots s_{i_{l(s_i\sigma)}}$ be a reduced decomposition. If $l(s_i\sigma) < l(\sigma) - 1$, then $\sigma = s_i(s_i\sigma) = s_i s_{i_1}s_{i_2} \cdots s_{i_{l(s_i\sigma)}}$ and hence $l(\sigma) \leq l(s_i\sigma) + 1 < l(\sigma) - 1 + 1 = l(\sigma)$ which leads to a contradiction. As a result, $l(s_i\sigma) \geq l(\sigma) - 1$. Together with the fact that $l(s_i\sigma) \leq l(\sigma) + 1$ and $l(s_i\sigma) \neq l(\sigma)$, we have $|l(s_i\sigma) - l(\sigma)| = 1$. \square

Lemma 1.8. *Let $n > 1$ be an integer and $\sigma \in S_n$. $l(s_i\sigma) = l(\sigma) - 1$ if and only if there exists a reduced decomposition $s_{r_1}s_{r_2} \cdots s_{r_{l(\sigma)}}$ such that $r_1 = i$.*

Proof. If σ has a reduced decomposition $s_{r_1}s_{r_2} \cdots s_{r_{l(\sigma)}}$ such that $r_1 = i$, then

$$s_i\sigma = s_i s_{r_1}s_{r_2} \cdots s_{r_{l(\sigma)}} = s_i^2 s_{r_2} \cdots s_{r_{l(\sigma)}} = s_{r_2} \cdots s_{r_{l(\sigma)}}$$

by item (i) in Proposition 1.1. By Lemma 1.4, we know that $s_{r_2} \cdots s_{r_{l(\sigma)}}$ is reduced and hence $l(s_i\sigma) = l(\sigma) - 1$.

If $l(s_i\sigma) = l(\sigma) - 1$, then consider a reduced decomposition of $s_i\sigma$, say $s_i\sigma = s_{i_1} \cdots s_{i_{l(s_i\sigma)-1}}$, by item (i) in Proposition 1.1, applying s_i on both sides gives $\sigma = s_i s_{i_1} \cdots s_{i_{l(s_i\sigma)-1}}$ with exactly $l(\sigma)$ transpositions, which implies $s_i s_{i_1} \cdots s_{i_{l(s_i\sigma)-1}}$ is a reduced decomposition of σ . Hence σ has a decomposition with s_i as the leftmost simple transposition. \square

Proposition 1.9. *$l(\sigma) = \text{inv}(\sigma)$ for any permutation σ .*

Proof. We first consider $\sigma^{-1}(1)$. If $\sigma^{-1}(1) \neq 1$, then all the integers before 1 in σ , i.e. $\sigma(1), \dots, \sigma(\sigma^{-1}(1) - 1)$, are all larger than 1, and hence $(r, \sigma^{-1}(1))$ are inversions of σ for all $1 \leq r < \sigma^{-1}(1)$. So by interchanging 1 with $\sigma(\sigma^{-1}(1) - 1)$, and then with $\sigma(\sigma^{-1}(1) - 2)$ until with $\sigma(1)$, we can put 1 to the leftmost of the new σ (the sigma after interchanging 1 with the $\sigma^{-1}(1) - 1$ integers). Indeed, by a previous note, the procedure described above is exactly applying transpositions on the right of σ , resulting in a new permutation, call it $\sigma^{(1)} = \sigma s_{\sigma^{-1}(1)-1} s_{\sigma^{-1}(1)-2} \cdots s_1$.

Note that each of the above procedure of moving 1 to the front decreases the the number of inversions by exactly 1.

We then use the same procedure by moving 2 to the second leftmost position of $\sigma^{(1)}$ by applying $\sigma^{(1)-1}(2) - 1$ simple transpositions on the right of $\sigma^{(1)}$ and get $\sigma^{(2)}$.

Continue this process until we get the $\sigma^{(n-1)}$ which has no inversion, i.e. $\sigma^{(n-1)} = id$. Since each time we apply the interchanging procedure, we are actually applying a simple transposition on the right and also decrease the number of inversion by exactly 1, we have performed exactly $\text{inv}(\sigma)$ interchanging procedures from σ to id . As a result, we get $\sigma s_{i_1} s_{i_2} \cdots s_{i_{\text{inv}(\sigma)}} = id$ and hence $\sigma = s_{i_{\text{inv}(\sigma)}} \cdots s_{i_1}$. (This also proves Theorem 1.3) which implies $l(\sigma) \leq \text{inv}(\sigma)$.

Let $s_{r_1} \cdots s_{r_{l(\sigma)}}$ be a reduced decomposition of σ . By Lemma 1.5, we know $\text{inv}(s_{r_1} \cdots s_{r_{l(\sigma)}}) \leq \text{inv}(s_{r_2} \cdots s_{r_{l(\sigma)}}) + 1 \leq \cdots \leq \text{inv}(s_{r_{l(\sigma)}}) + l(\sigma) - 1 = l(\sigma)$ and hence we get $\text{inv}(\sigma) \leq l(\sigma)$.

As a result, $l(\sigma) = \text{inv}(\sigma)$. □

Note that $n, n-1, \dots, 1$ has the longest length in S_n as it has the maximum number (namely, $\binom{n}{2}$) of inversions.

Corollary 1.10. *Let $\sigma = n, n-1, \dots, 1$. Then for any $i \in [n-1]$, there is a reduced word of σ starting with i .*

Proof. Let $i \in [n-1]$. Since $|l(s_i\sigma) - l(\sigma)| = 1$ and $l(\sigma) > l(s_i\sigma)$ as σ has the longest length among all permutations in S_n , we have $l(s_i\sigma) = l(\sigma) - 1$. Hence result follows by Lemma 1.8. □

There are several equivalent definitions of Bruhat order on S_n and we will use the reduced word definition. See [1] for further discussion.

Definition 1.6. *Let n be a positive integer. Define a partial ordering \leq on S_n such that $\sigma \leq \gamma$ if and only if there exists a reduced word of σ which is a substring (not necessarily consecutive) of some reduced word of γ .*

Lemma 1.11. *Let $k \geq 2$ be a positive integer and $i_1 i_2 \dots i_k$ be a reduced word. Let $\sigma' = s_{i_2} \dots s_{i_k}$ and $\sigma = s_{i_1} \sigma'$. Then $\{\tau | \tau \leq \sigma\} = \{s_{i_1} \gamma, \gamma | \gamma \leq \sigma', l(s_{i_1} \gamma) = l(\gamma) + 1\}$.*

Proof. First note that as $i_1 i_2 \dots i_k$ is reduced, by Lemma 1.4, $i_2 \dots i_k$ is a reduced word of σ' . Hence $\sigma' \leq \sigma$.

Consider γ such that $\gamma \leq \sigma'$ such that $l(s_{i_1} \gamma) = l(\gamma) + 1$.

Since $\gamma \leq \sigma'$ and $\sigma' \leq \sigma$, we have $\gamma \leq \sigma$. Also, $\gamma \leq \sigma'$ implies γ has a reduced word which is a substring of $i_2 \dots i_k$, say $i_{r_1} i_{r_2} \dots i_{r_{l(\gamma)}}$ where $2 \leq r_1 < r_2 < \dots < r_{l(\gamma)} \leq k$. Then $s_{i_1} \gamma = s_{i_1} s_{i_{r_1}} s_{i_{r_2}} \dots s_{i_{r_{l(\gamma)}}$ which is reduced as $l(s_{i_1} \gamma) = l(\gamma) + 1$. Hence $s_{i_1} \gamma \leq \sigma$ (as $i_1 i_{r_1} \dots i_{r_{l(\gamma)}}$ is a substring of $i_1 i_2 \dots i_k$).

We thus have $\{\tau | \tau \leq \sigma\} \supseteq \{s_{i_1} \gamma, \gamma | \gamma \leq \sigma', l(s_{i_1} \gamma) = l(\gamma) + 1\}$.

Now consider τ such that $\tau \leq \sigma$. Let $i_{j_1} i_{j_2} \dots i_{j_{l(\tau)}}$ be a reduced word of τ , where $1 \leq j_1 < j_2 < \dots < j_{l(\tau)} \leq k$. If $j_1 \neq 1$ for any reduced word of τ , then by Lemma 1.8, $l(s_{i_1} \tau) \neq l(\tau) - 1$. By Lemma 1.7, $l(s_{i_1} \tau) = l(\tau) + 1$. Also, $2 \leq j_1 < j_2 < \dots < j_{l(\tau)} \leq k$ implies $\tau \leq \sigma'$. As a result, $\tau \leq \sigma'$ and $l(s_{i_1} \tau) = l(\tau) + 1$ if $j_1 \neq 1$ for any reduced word $i_{j_1} i_{j_2} \dots i_{j_{l(\tau)}}$ of τ .

If $j_1 = 1$, then we can write $\tau = s_{i_1} \tau'$ where $\tau' = s_{i_{j_2}} s_{i_{j_3}} \dots s_{i_{j_{l(\tau)}}$ which is also a reduced decomposition by Lemma 1.4. Since $1 = j_1 < j_2$, we have $\tau' \leq \sigma'$. Also $l(\tau') = l(\sigma) - 1$ which implies $l(s_{i_1} \tau') = l(\sigma) = l(\tau') + 1$. Hence $\tau = s_{i_1} \tau'$ where $\tau' \leq \sigma', l(s_{i_1} \tau') = l(\tau') + 1$ if $j_1 = 1$ for some reduced word $i_{j_1} i_{j_2} \dots i_{j_{l(\tau)}}$ of τ .

Therefore $\{\tau | \tau \leq \sigma\} \subseteq \{s_{i_1} \gamma, \gamma | \gamma \leq \sigma', l(s_{i_1} \gamma) = l(\gamma) + 1\}$ and result follows. □

Chapter 2

Demazure atoms and characters

2.1 Linear operators

Let P be the polynomial ring $\mathbb{Z}[x_1, x_2, \dots]$ and S_∞ be the permutation group of the positive integers, acting on P by permuting the indices of the variables. For any positive integer i , define linear operators

$$\begin{aligned}\partial_i &:= \frac{1 - s_i}{x_i - x_{i+1}} \\ \pi_i &:= \partial_i x_i \\ \theta_i &:= x_{i+1} \partial_i\end{aligned}$$

where s_i is the elementary transposition $(i, i + 1)$ and 1 is the identity element in S_∞ .

Note that for $f \in P$, $(x_i - x_{i+1})$ is a factor of $(1 - s_i)f$ and hence $\partial_i f \in P$. Therefore, $\pi_i f, \theta_i f \in P$.

Example 3. Let $i = 2$ and consider the monomials $x_1^5 x_2^4 x_3$, $x_1^3 x_3^2$ and $x_1 x_2^2 x_3^2$, we have

$$1 \text{ a) } \partial_2(x_1^5 x_2^4 x_3) = \frac{x_1^5 x_2^4 x_3 - s_2(x_1^5 x_2^4 x_3)}{x_2 - x_3} = \frac{x_1^5 x_2^4 x_3 - x_1^5 x_2 x_3^4}{x_2 - x_3} = x_1^5(x_2^3 x_3 + x_2^2 x_3^2 + x_2 x_3^3)$$

$$\text{b) } \pi_2(x_1^5 x_2^4 x_3) = \partial_2 x_2(x_1^5 x_2^4 x_3) = \partial_2(x_1^5 x_2^5 x_3) = x_1^5(x_2^4 x_3 + x_2^3 x_3^2 + x_2^2 x_3^3 + x_2 x_3^4)$$

$$\text{c) } \theta_2(x_1^5 x_2^4 x_3) = x_3 \partial_2(x_1^5 x_2^4 x_3) = x_1^5(x_2^3 x_3^2 + x_2^2 x_3^3 + x_2 x_3^4)$$

$$2 \text{ a) } \partial_2(x_1^3 x_3^2) = \frac{x_1^3 x_3^2 - s_2(x_1^3 x_3^2)}{x_2 - x_3} = \frac{x_1^3 x_3^2 - x_1^3 x_2^2}{x_2 - x_3} = -x_1^3(x_2 + x_3)$$

$$\text{b) } \pi_2(x_1^3 x_3^2) = \partial_2 x_2(x_1^3 x_3^2) = \partial_2(x_1^3 x_2 x_3^2) = -x_1^3 x_2 x_3$$

$$\text{c) } \theta_2(x_1^3 x_3^2) = x_3 \partial_2(x_1^3 x_3^2) = -x_1^3(x_2 x_3 + x_3^2)$$

$$3 \text{ a) } \partial_2(x_1 x_2^2 x_3^2) = \frac{x_1 x_2^2 x_3^2 - s_2(x_1 x_2^2 x_3^2)}{x_2 - x_3} = \frac{x_1 x_2^2 x_3^2 - x_1 x_2^2 x_3^2}{x_2 - x_3} = 0$$

$$\text{b) } \pi_2(x_1 x_2^2 x_3^2) = \partial_2 x_2(x_1 x_2^2 x_3^2) = \partial_2(x_1 x_2^3 x_3^2) = x_1 x_2^2 x_3^2$$

$$\text{c) } \theta_2(x_1 x_2^2 x_3^2) = x_3 \partial_2(x_1 x_2^2 x_3^2) = 0$$

Proposition 2.1. *For any positive integer i , we have*

1. $\pi_i = \theta_i + 1$;
2. $\pi_i \theta_i = \theta_i \pi_i$;
3. $s_i \partial_i = -\partial_i s_i = \partial_i$, $s_i \pi_i = \pi_i$, $\pi_i s_i = -\partial_i x_{i+1}$, $s_i \theta_i = x_i \partial_i$, $\theta_i s_i = -\theta_i$;
4. $\partial_i \partial_i = 0$, $\partial_i \pi_i = \theta_i \partial_i = 0$, $\pi_i \partial_i = -\partial_i \theta_i = \partial_i$;
5. $\pi_i \pi_i = \pi_i$, $\theta_i \theta_i = -\theta_i$, $\pi_i \theta_i = \theta_i \pi_i = 0$.

Proof. Let $f \in P$. Then

$$\begin{aligned}
\pi_i f &= \partial_i x_i f = \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}} \\
&= \frac{x_i f - x_{i+1} s_i f}{x_i - x_{i+1}} \\
&= \frac{x_{i+1} f - x_{i+1} s_i f + (x_i - x_{i+1}) f}{x_i - x_{i+1}} \\
&= x_{i+1} \frac{f - s_i(f)}{x_i - x_{i+1}} + f \\
&= x_{i+1} \partial_i f + f \\
&= (\theta_i + 1) f
\end{aligned}$$

and hence $\pi_i = \theta_i + 1$.

By item 1, we have $\pi_i \theta_i = (\theta_i + 1) \theta_i = \theta_i \theta_i + \theta_i = \theta_i(\theta_i + 1) = \theta_i \pi_i$, proving item 2.

$$s_i \partial_i f = s_i \left(\frac{f - s_i f}{x_i - x_{i+1}} \right) = \frac{s_i f - s_i s_i f}{x_{i+1} - x_i} = \frac{s_i f - f}{x_{i+1} - x_i} = \frac{f - s_i f}{x_i - x_{i+1}} = \partial_i f.$$

$$\partial_i s_i f = \frac{s_i f - s_i(s_i f)}{x_i - x_{i+1}} = -\frac{f - s_i f}{x_i - x_{i+1}} = -\partial_i f.$$

$$s_i \pi_i f = s_i(\partial_i x_i f) = s_i \partial_i(x_i f) = \partial_i(x_i f) = \pi_i f.$$

$$\pi_i s_i f = \partial_i(x_i s_i f) = \partial_i s_i(x_{i+1} f) = -\partial_i x_{i+1} f.$$

$$s_i \theta_i f = s_i(x_{i+1} \partial_i f) = x_i s_i \partial_i f = x_i \partial_i f.$$

$$\theta_i s_i f = x_{i+1} \partial_i s_i f = -x_{i+1} \partial_i f = -\theta_i f.$$

Hence, $s_i \partial_i = \partial_i$, $\partial_i s_i = -\partial_i$, $s_i \pi_i = \pi_i$, $\pi_i s_i = -\partial_i x_{i+1}$, $s_i \theta_i = x_i \partial_i$, $\theta_i s_i = -\theta_i$ and item 3 follows.

By item 3, $\partial_i \partial_i f = \frac{\partial_i f - s_i \partial_i f}{x_i - x_{i+1}} = \frac{\partial_i f - \partial_i f}{x_i - x_{i+1}} = 0$.

Hence $\partial_i \pi_i f = \partial_i \partial_i x_i f = 0$ and $\theta_i \partial_i f = x_{i+1} \partial_i \partial_i f = 0$.

Then $\pi_i \partial_i f = (1 + \theta_i) \partial_i f = \partial_i f + \theta_i \partial_i f = \partial_i f$.

Also, $\partial_i \theta_i f = \partial_i (\pi_i - 1) f = \partial_i \pi_i f - \partial_i f = -\partial_i f$.

As a result, $\partial_i \partial_i = \partial_i \pi_i = \theta_i \partial_i = 0$, $\pi_i \partial_i = \partial_i$, $\partial_i \theta_i = -\partial_i$, proving item 4.

Now by item 4, we have

$$\pi_i \pi_i f = (\pi_i \partial_i)(x_i f) = \partial_i(x_i f) = \pi_i f,$$

$$\theta_i \theta_i f = (x_{i+1} \partial_i) \theta_i f = x_{i+1} (\partial_i \theta_i) f = -x_{i+1} \partial_i f = -\theta_i f,$$

$$\text{Therefore } \theta_i \pi_i f = \theta_i (\theta_i + 1) f = \theta_i \theta_i f + \theta_i f = -\theta_i f + \theta_i f = 0.$$

Furthermore, $\pi_i \theta_i f = (\theta_i + 1) \theta_i f = \theta_i \theta_i f + \theta_i f = 0$ and item 5 follows.

□

Proposition 2.2. $\partial_i \partial_j = \partial_j \partial_i$, $\pi_i \pi_j = \pi_j \pi_i$ and $\theta_i \theta_j = \theta_j \theta_i$ for $|i - j| \geq 2$.

Proof. Let $f \in P$ and $i, j \in \mathbb{N}$ such that $|i - j| \geq 2$. Then

$$\partial_i \partial_j f = \frac{\partial_j f - s_i \partial_j f}{x_i - x_{i+1}}.$$

Since $s_i s_j = s_j s_i$, we have

$$s_i \partial_j f = s_i \frac{f - s_j f}{x_j - x_{j+1}} = \frac{s_i f - s_i s_j f}{x_j - x_{j+1}} = \frac{s_i f - s_j s_i f}{x_j - x_{j+1}} = \partial_j s_i f.$$

Hence

$$\begin{aligned} \partial_i \partial_j f &= \frac{\partial_j f - \partial_j(s_i f)}{x_i - x_{i+1}} \\ &= \frac{\partial_j(f - s_i f)}{x_i - x_{i+1}} \\ &= \frac{(f - s_i f) - s_j(f - s_i f)}{(x_j - x_{j+1})(x_i - x_{i+1})} \\ &= \frac{f - s_i f - s_j f + s_i s_j f}{(x_i - x_{i+1})(x_j - x_{j+1})}. \end{aligned}$$

$$\text{Similarly } \partial_j \partial_i f = \frac{f - s_j f - s_i f + s_j s_i f}{(x_j - x_{j+1})(x_i - x_{i+1})} = \frac{f - s_i f - s_j f + s_i s_j f}{(x_i - x_{i+1})(x_j - x_{j+1})}.$$

Thus $\partial_i \partial_j = \partial_j \partial_i$.

Note that $x_i \partial_j f = x_i \frac{f - s_j f}{x_j - x_{j+1}} = \frac{x_i f - x_i s_j f}{x_j - x_{j+1}} = \frac{x_i f - s_j x_i f}{x_j - x_{j+1}} = \partial_j x_i f$, that is, $x_i \partial_j = \partial_j x_i$.

Similarly, $x_j \partial_i = \partial_i x_j$.

Hence we have

$$\pi_i \pi_j f = \partial_i x_i \partial_j x_j f = \partial_i \partial_j x_i x_j f = \partial_j \partial_i x_i x_j f = \partial_j \partial_i x_j x_i f = \partial_j x_j \partial_i x_i f = \pi_j \pi_i f.$$

$$\theta_i \theta_j = (\pi_i - 1)(\pi_j - 1) = \pi_i \pi_j - \pi_i - \pi_j + 1 = \pi_j \pi_i - \pi_i - \pi_j + 1 = (\pi_j - 1)(\pi_i - 1) = \theta_j \theta_i.$$

□

Proposition 2.3. *For any positive integer i , linear operators ∂_i , π_i and θ_i satisfy the braid relation.*

That is, $\partial_i \partial_{i+1} \partial_i = \partial_{i+1} \partial_i \partial_{i+1}$, $\pi_i \pi_{i+1} \pi_i = \pi_{i+1} \pi_i \pi_{i+1}$ and $\theta_i \theta_{i+1} \theta_i = \theta_{i+1} \theta_i \theta_{i+1}$.

Proof. Let $f \in P$ and $i \in \mathbb{N}$.

$$\begin{aligned} & \partial_i \partial_{i+1} f \\ = & \frac{\partial_{i+1} f - s_i \partial_{i+1} f}{x_i - x_{i+1}} \\ = & \frac{\frac{f - s_{i+1} f}{x_{i+1} - x_{i+2}} - \frac{s_i f - s_i s_{i+1} f}{x_i - x_{i+2}}}{x_i - x_{i+1}} \\ = & \frac{(x_i - x_{i+2})(f - s_{i+1} f) - (x_{i+1} - x_{i+2})s_i f + (x_{i+1} - x_{i+2})s_i s_{i+1} f}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})}. \end{aligned}$$

Hence we have

$$\begin{aligned}
& \partial_i \partial_{i+1} \partial_i f \\
= & \frac{(x_i - x_{i+2})(\partial_i f - s_{i+1} \partial_i f) - (x_{i+1} - x_{i+2}) s_i \partial_i f + (x_{i+1} - x_{i+2}) s_i s_{i+1} \partial_i f}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})} \\
= & \frac{(x_i - x_{i+2}) \partial_i f - (x_i - x_{i+2}) s_{i+1} \partial_i f - (x_{i+1} - x_{i+2}) \partial_i f + (x_{i+1} - x_{i+2}) s_i s_{i+1} \partial_i f}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})} \\
= & \frac{(x_i - x_{i+1}) \partial_i f - (x_i - x_{i+2}) s_{i+1} \partial_i f + (x_{i+1} - x_{i+2}) s_i s_{i+1} \partial_i f}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})} \\
= & \frac{f - s_i f - s_{i+1} f + s_{i+1} s_i f + s_i s_{i+1} f - s_i s_{i+1} s_i f}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})} \\
= & \frac{(1 - s_i)(1 - s_{i+1})(1 - s_i)}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})} f.
\end{aligned}$$

Also, we have

$$\begin{aligned}
& s_{i+1} \partial_i \partial_{i+1} f \\
= & \frac{(x_i - x_{i+1})(s_{i+1} f - f) - (x_{i+2} - x_{i+1})(s_{i+1} s_i f - s_{i+1} s_i s_{i+1} f)}{(x_{i+2} - x_{i+1})(x_i - x_{i+1})(x_i - x_{i+2})} \\
= & \frac{(x_{i+1} - x_i)(s_{i+1} f - f) - (x_{i+1} - x_{i+2})(s_{i+1} s_i f - s_{i+1} s_i s_{i+1} f)}{(x_{i+1} - x_{i+2})(x_i - x_{i+1})(x_i - x_{i+2})}
\end{aligned}$$

which implies

$$\begin{aligned}
& \partial_i \partial_{i+1} f - s_{i+1} \partial_i \partial_{i+1} f \\
= & \frac{(x_{i+1} - x_{i+2})(f - s_i f - s_{i+1} f + s_{i+1} s_i f + s_i s_{i+1} f - s_{i+1} s_i s_{i+1} f)}{(x_{i+1} - x_{i+2})(x_i - x_{i+1})(x_i - x_{i+2})}
\end{aligned}$$

and thus

$$\begin{aligned}
\partial_{i+1} \partial_i \partial_{i+1} f &= \frac{\partial_i \partial_{i+1} f - s_{i+1} \partial_i \partial_{i+1} f}{x_{i+1} - x_{i+2}} \\
&= \frac{(x_{i+1} - x_{i+2}) \frac{(f - s_i f - s_{i+1} f + s_{i+1} s_i f + s_i s_{i+1} f - s_{i+1} s_i s_{i+1} f)}{(x_{i+1} - x_{i+2})(x_i - x_{i+1})(x_i - x_{i+2})}}{x_{i+1} - x_{i+2}} \\
&= \frac{f - s_i f - s_{i+1} f + s_{i+1} s_i f + s_i s_{i+1} f - s_{i+1} s_i s_{i+1} f}{(x_{i+1} - x_{i+2})(x_i - x_{i+1})(x_i - x_{i+2})} = \partial_i \partial_{i+1} \partial_i f
\end{aligned}$$

as $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$.

As a result, we get

$$\begin{aligned}
\partial_i \partial_{i+1} \partial_i &= \frac{(1-s_i)(1-s_{i+1})(1-s_i)}{(x_{i+1}-x_{i+2})(x_i-x_{i+2})(x_i-x_{i+1})r} \\
&= \frac{1+s_i s_{i+1} + s_{i+1} s_i - s_i - s_{i+1} - s_i s_{i+1} s_i}{(x_{i+1}-x_{i+2})(x_i-x_{i+2})(x_i-x_{i+1})} \\
&= \frac{(1-s_{i+1})(1-s_i)(1-s_{i+1})}{(x_{i+1}-x_{i+2})(x_i-x_{i+2})(x_i-x_{i+1})} = \partial_{i+1} \partial_i \partial_{i+1}.
\end{aligned}$$

Next we prove $\pi_i \pi_{i+1} \pi_i = \pi_{i+1} \pi_i \pi_{i+1}$.

$$\pi_i f = \partial_i x_i f = \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}}, \quad \pi_{i+1} f = \partial_{i+1} x_{i+1} f = \frac{x_{i+1} f - s_{i+1}(x_{i+1} f)}{x_{i+1} - x_{i+2}}$$

and hence

$$\begin{aligned}
&\pi_i \pi_{i+1} f \\
&= \frac{x_i \pi_{i+1} f - s_i(x_i \pi_{i+1} f)}{x_i - x_{i+1}} \\
&= \frac{x_i \frac{x_{i+1} f - s_{i+1}(x_{i+1} f)}{x_{i+1} - x_{i+2}} - s_i \left(x_i \frac{x_{i+1} f - s_{i+1}(x_{i+1} f)}{x_{i+1} - x_{i+2}} \right)}{x_i - x_{i+1}} \\
&= \frac{\frac{x_i x_{i+1} f - s_{i+1}(x_i x_{i+1} f)}{x_{i+1} - x_{i+2}} - \frac{s_i(x_i x_{i+1} f) - s_i s_{i+1}(x_i x_{i+1} f)}{x_i - x_{i+2}}}{x_i - x_{i+1}} \\
&= \frac{(x_i - x_{i+2})x_i x_{i+1} f - s_{i+1}((x_i - x_{i+1})x_i x_{i+1} f)}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})} \\
&\quad - \frac{s_i((x_i - x_{i+2})x_i x_{i+1} f) - s_i s_{i+1}((x_i - x_{i+1})x_i x_{i+1} f)}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})} \\
&= \frac{(1-s_i) \left((x_i - x_{i+2})x_i x_{i+1} f - s_{i+1}((x_i - x_{i+1})x_i x_{i+1} f) \right)}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})}
\end{aligned}$$

and

$$\begin{aligned}
& \pi_{i+1}\pi_i f \\
= & \frac{x_{i+1}\pi_i f - s_{i+1}(x_{i+1}\pi_i f)}{x_{i+1} - x_{i+2}} \\
= & \frac{x_{i+1} \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}} - s_{i+1} \left(x_{i+1} \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}} \right)}{x_{i+1} - x_{i+2}} \\
= & \frac{\frac{x_i x_{i+1} f - s_i(x_i^2 f)}{x_i - x_{i+1}} - \frac{s_{i+1}(x_i x_{i+1} f) - s_{i+1} s_i(x_i^2 f)}{x_i - x_{i+2}}}{x_{i+1} - x_{i+2}} \\
= & \frac{(x_i - x_{i+2})x_i x_{i+1} f - s_i((x_{i+1} - x_{i+2})x_i^2 f)}{(x_i - x_{i+1})(x_i - x_{i+2})(x_{i+1} - x_{i+2})} \\
& - \frac{s_{i+1}((x_i - x_{i+2})x_i x_{i+1} f) - s_{i+1} s_i((x_{i+1} - x_{i+2})x_i^2 f)}{(x_i - x_{i+1})(x_i - x_{i+2})(x_{i+1} - x_{i+2})} \\
= & \frac{(1 - s_{i+1}) \left((x_i - x_{i+2})x_i x_{i+1} f - s_i((x_{i+1} - x_{i+2})x_i^2 f) \right)}{(x_i - x_{i+1})(x_i - x_{i+2})(x_{i+1} - x_{i+2})}.
\end{aligned}$$

So

$$\begin{aligned}
& \pi_i \pi_{i+1} \pi_i f \\
= & \frac{(1 - s_i) \left((x_i - x_{i+2})x_i x_{i+1} \pi_i f - s_{i+1}((x_i - x_{i+1})x_i x_{i+1} \pi_i f) \right)}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})} \\
= & \frac{(1 - s_i) \left((x_i - x_{i+2})x_i x_{i+1} \pi_i f - s_{i+1} \left((x_i - x_{i+1})x_i x_{i+1} \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}} \right) \right)}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})} \\
= & \frac{(1 - s_i) \left((x_i - x_{i+2})x_i x_{i+1} \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}} - x_i x_{i+2} (x_i s_{i+1} f - s_{i+1} s_i(x_i f)) \right)}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})}.
\end{aligned}$$

Since

$$\begin{aligned}
& s_i \left((x_i - x_{i+2})x_i x_{i+1} \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}} - x_i x_{i+2} (x_i s_{i+1} f - s_{i+1} s_i(x_i f)) \right) \\
= & (x_{i+1} - x_{i+2})x_i x_{i+1} \frac{s_i(x_i f) - x_i f}{x_{i+1} - x_i} - x_{i+1} x_{i+2} (x_{i+1} s_i s_{i+1} f - s_i s_{i+1} s_i(x_i f)) \\
= & (x_{i+1} - x_{i+2})x_i x_{i+1} \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}} - x_{i+1} x_{i+2} (x_{i+1} s_i s_{i+1} f - s_i s_{i+1} s_i(x_i f)),
\end{aligned}$$

we have

$$\begin{aligned}
& (1 - s_i) \left((x_i - x_{i+2}) x_i x_{i+1} \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}} - x_i x_{i+2} (x_i s_{i+1} f - s_{i+1} s_i(x_i f)) \right) \\
= & x_i x_{i+1} (x_i f - s_i(x_i f)) - x_i^2 x_{i+2} s_{i+1} f \\
& + x_i x_{i+2} s_{i+1} s_i(x_i f) + x_{i+1}^2 x_{i+2} s_i s_{i+1} f - x_{i+1} x_{i+2} s_i s_{i+1} s_i(x_i f) \\
= & (1 - s_i - s_{i+1} + s_i s_{i+1} + s_{i+1} s_i - s_i s_{i+1} s_i) (x_i^2 x_{i+1} f)
\end{aligned}$$

$$\text{and hence } \pi_i \pi_{i+1} \pi_i f = \frac{(1 - s_i - s_{i+1} + s_i s_{i+1} + s_{i+1} s_i - s_i s_{i+1} s_i) (x_i^2 x_{i+1} f)}{(x_{i+1} - x_{i+2})(x_i - x_{i+2})(x_i - x_{i+1})}.$$

$$\begin{aligned}
& \pi_{i+1} \pi_i \pi_{i+1} f \\
= & \frac{(1 - s_{i+1}) \left((x_i - x_{i+2}) x_i x_{i+1} \pi_{i+1} f - s_i \left((x_{i+1} - x_{i+2}) x_i^2 \pi_{i+1} f \right) \right)}{(x_i - x_{i+1})(x_i - x_{i+2})(x_{i+1} - x_{i+2})} \\
= & \frac{(1 - s_{i+1}) \left((x_i - x_{i+2}) x_i x_{i+1} \pi_{i+1} f - s_i \left((x_{i+1} - x_{i+2}) x_i^2 \frac{x_{i+1} f - s_{i+1}(x_{i+1} f)}{x_{i+1} - x_{i+2}} \right) \right)}{(x_i - x_{i+1})(x_i - x_{i+2})(x_{i+1} - x_{i+2})} \\
= & \frac{(1 - s_{i+1}) \left((x_i - x_{i+2}) x_i x_{i+1} \pi_{i+1} f - s_i \left(x_i^2 (x_{i+1} f - s_{i+1}(x_{i+1} f)) \right) \right)}{(x_i - x_{i+1})(x_i - x_{i+2})(x_{i+1} - x_{i+2})}.
\end{aligned}$$

Since

$$\begin{aligned}
& s_{i+1} \left((x_i - x_{i+2}) x_i x_{i+1} \pi_{i+1} f - s_i \left(x_i^2 (x_{i+1} f - s_{i+1}(x_{i+1} f)) \right) \right) \\
= & (x_i - x_{i+1}) x_i x_{i+2} \frac{s_{i+1}(x_{i+1} f) - x_{i+1} f}{x_{i+2} - x_{i+1}} - s_{i+1} s_i \left(x_i^2 (x_{i+1} f - s_{i+1}(x_{i+1} f)) \right) \\
= & (x_i - x_{i+1}) x_i x_{i+2} \frac{x_{i+1} f - s_{i+1}(x_{i+1} f)}{x_{i+1} - x_{i+2}} - s_{i+1} s_i \left(x_i^2 (x_{i+1} f - s_{i+1}(x_{i+1} f)) \right),
\end{aligned}$$

we have

$$\begin{aligned}
& (1 - s_{i+1}) \left((x_i - x_{i+2}) x_i x_{i+1} \pi_{i+1} f - s_i (x_i^2 (x_{i+1} f - s_{i+1} (x_{i+1} f))) \right) \\
= & x_i^2 (x_{i+1} f - s_{i+1} (x_{i+1} f)) - s_i (x_i^2 x_{i+1} f) + s_i (x_i^2 s_{i+1} (x_{i+1} f)) \\
& + s_{i+1} s_i \left(x_i^2 (x_{i+1} f - s_{i+1} (x_{i+1} f)) \right) \\
= & (1 - s_i - s_{i+1} + s_i s_{i+1} + s_{i+1} s_i - s_{i+1} s_i s_{i+1}) (x_i^2 x_{i+1} f)
\end{aligned}$$

$$\text{and hence } \pi_{i+1} \pi_i \pi_{i+1} f = \frac{(1 - s_i - s_{i+1} + s_i s_{i+1} + s_{i+1} s_i - s_{i+1} s_i s_{i+1}) (x_i^2 x_{i+1} f)}{(x_i - x_{i+1})(x_i - x_{i+2})(x_{i+1} - x_{i+2})}.$$

Now $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$ implies $\pi_i \pi_{i+1} \pi_i = \pi_{i+1} \pi_i \pi_{i+1}$ and result follows.

By item 1. and 5. of Proposition 2.1, we have

$$\begin{aligned}
& \theta_i \theta_{i+1} \theta_i \\
= & (\pi_i - 1)(\pi_{i+1} - 1)(\pi_i - 1) \quad (\text{by item 1.}) \\
= & \pi_i \pi_{i+1} \pi_i - \pi_i \pi_{i+1} - \pi_{i+1} \pi_i - \pi_i \pi_i + 2\pi_i + \pi_{i+1} - 1 \\
= & \pi_i \pi_{i+1} \pi_i - \pi_i \pi_{i+1} - \pi_{i+1} \pi_i - \pi_i + 2\pi_i + \pi_{i+1} - 1 \quad (\text{by item 5.}) \\
= & \pi_i \pi_{i+1} \pi_i - \pi_i \pi_{i+1} - \pi_{i+1} \pi_i + \pi_i + \pi_{i+1} - 1
\end{aligned}$$

and

$$\begin{aligned}
& \theta_{i+1} \theta_i \theta_{i+1} \\
= & (\pi_{i+1} - 1)(\pi_i - 1)(\pi_{i+1} - 1) \quad (\text{by item 1.}) \\
= & \pi_{i+1} \pi_i \pi_{i+1} - \pi_i \pi_{i+1} - \pi_{i+1} \pi_i - \pi_{i+1} \pi_{i+1} + 2\pi_{i+1} + \pi_i - 1 \\
= & \pi_{i+1} \pi_i \pi_{i+1} - \pi_i \pi_{i+1} - \pi_{i+1} \pi_i - \pi_{i+1} + 2\pi_{i+1} + \pi_i - 1 \quad (\text{by item 5.}) \\
= & \pi_{i+1} \pi_i \pi_{i+1} - \pi_i \pi_{i+1} - \pi_{i+1} \pi_i + \pi_{i+1} + \pi_i - 1
\end{aligned}$$

and by using the fact that $\pi_i \pi_{i+1} \pi_i = \pi_{i+1} \pi_i \pi_{i+1}$, we have $\theta_i \theta_{i+1} \theta_i = \theta_{i+1} \theta_i \theta_{i+1}$.

□

For any permutation $\sigma \neq id$, define $\partial_\sigma = \partial_{i_1} \dots \partial_{i_j}$ where $s_{i_1} \dots s_{i_j}$ is a reduced decomposition of σ . By the same argument in Proposition 2.2 and Proposition 2.3, we also have $\theta_\sigma = \theta_{i_1} \dots \theta_{i_j}$ and $\pi_\sigma = \pi_{i_1} \dots \pi_{i_j}$. We define $\partial_{id} = \theta_{id} = \pi_{id} = id$.

Lemma 2.4. *Let $n > 1$ be an integer and consider a permutation $\gamma \in S_n$. For $1 \leq i \leq n - 1$,*

$$\theta_i \theta_\gamma = \begin{cases} -\theta_\gamma & \text{if } l(s_i \gamma) = l(\gamma) - 1 \\ \theta_{s_i \gamma} & \text{if } l(s_i \gamma) = l(\gamma) + 1 \end{cases}$$

Proof. By Lemma 1.8, if $l(s_i \gamma) = l(\gamma) - 1$, then there exists a reduced decomposition of $\gamma = s_i s_{r_2} \dots s_{r_{l(\gamma)}}$ and hence $\theta_\gamma = \theta_i \theta_{r_2} \dots \theta_{r_{l(\gamma)}}$. By item 5. in Proposition 2.1, we have $\theta_i \theta_\gamma = (\theta_i \theta_i) \theta_{r_2} \dots \theta_{r_{l(\gamma)}} = (-\theta_i) \theta_{r_2} \dots \theta_{r_{l(\gamma)}} = -\theta_i \theta_{r_2} \dots \theta_{r_{l(\gamma)}} = -\theta_\gamma$. Otherwise if $l(s_i \gamma) = l(\gamma) + 1$, $s_i s_{i_1} s_{i_2} \dots s_{i_{l(\gamma)}}$ is a reduced decomposition of $s_i \gamma$ for any reduced decomposition $s_{i_1} s_{i_2} \dots s_{i_{l(\gamma)}}$ of γ . Thus $\theta_i \theta_\gamma = \theta_i \theta_{i_1} \theta_{i_2} \dots \theta_{i_{l(\gamma)}} = \theta_{s_i \gamma}$. □

Lemma 2.5. *For any permutation σ , $\pi_\sigma = \sum_{\gamma \leq \sigma} \theta_\gamma$.*

Proof. Let $k = l(\sigma)$. We prove the statement by induction on k .

For $\sigma = id$ (i.e. $k = 0$), $\pi_{id} = \theta_{id} = id$.

When $k = 1$, then $\sigma = s_i$ for some positive integer i . By item 1. in Proposition 2.1, we have $\pi_\sigma = \pi_{s_i} = \pi_i = 1 + \theta_i = \theta_{id} + \theta_{s_i}$. Hence the statement is true for $l(\sigma) = 1$.

Assume the statement is true for all non-negative integers $k \leq m$ for some $m \geq 1$.

Let $l(\sigma) = m + 1$. Let $s_{i_1} s_{i_2} \dots s_{i_{m+1}}$ be a reduced decomposition of σ . Let $\sigma' = s_{i_2} \dots s_{i_{m+1}}$ (which implies $l(\sigma') \leq m$ by definition) and hence $\sigma = s_{i_1} \sigma'$. Note that by Lemma 1.4, $s_{i_2} \dots s_{i_{m+1}}$ is a reduced decomposition of σ' .

By induction assumption, $\pi'_\sigma = \sum_{\gamma \leq \sigma'} \theta_\gamma$. Now we have

$$\begin{aligned}
& \pi_\sigma \\
&= \pi_{i_1} \pi_{i_2} \cdots \pi_{i_{m+1}} \\
&= \pi_{i_1} \pi_{\sigma'} \\
&= (1 + \theta_{i_1}) \sum_{\gamma \leq \sigma'} \theta_\gamma \\
&= (1 + \theta_{i_1}) \left(\sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) + 1}} \theta_\gamma + \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) - 1}} \theta_\gamma \right) \quad (\text{by Lemma 1.7}) \\
&= \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) + 1}} \theta_\gamma + \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) - 1}} \theta_\gamma + \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) + 1}} \theta_{i_1} \theta_\gamma + \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) - 1}} \theta_{i_1} \theta_\gamma \\
&= \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) + 1}} \theta_\gamma + \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) - 1}} \theta_\gamma + \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) + 1}} \theta_{s_{i_1} \gamma} + \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) - 1}} -\theta_\gamma \\
&\quad (\text{by Lemma 2.4}) \\
&= \sum_{\substack{\gamma \leq \sigma' \\ l(s_{i_1} \gamma) = l(\gamma) + 1}} (\theta_\gamma + \theta_{s_{i_1} \gamma}) \\
&= \sum_{\tau \leq \sigma} \theta_\tau \quad (\text{by Lemma 1.11})
\end{aligned}$$

and result follows by induction. □

Lemma 2.6. *For any $f, g \in P$ and $i \in \mathbb{N}$, we have*

1. $\partial_i(fg) = (\partial_i f)g + (s_i f)(\partial_i g)$;
2. $\theta_i(fg) = (\theta_i f)g + (s_i f)(\theta_i g)$;
3. $\pi_i(fg) = (\pi_i f)g + (s_i f)(\theta_i g)$.

Proof.

$$\begin{aligned} & \partial_i(fg) \\ &= \frac{fg - s_i(fg)}{x_i - x_{i+1}} \\ &= \frac{fg - (s_i f)g + (s_i f)g - s_i f s_i g}{x_i - x_{i+1}} \\ &= g \frac{f - s_i f}{x_i - x_{i+1}} + (s_i f) \frac{g - s_i g}{x_i - x_{i+1}} \\ &= g(\partial_i f) + (s_i f)(\partial_i g) \\ &= (\partial_i f)g + (s_i f)(\partial_i g). \end{aligned}$$

Therefore

$$\begin{aligned} & \theta_i(fg) \\ &= x_{i+1} \partial_i(fg) \\ &= x_{i+1} ((\partial_i f)g + (s_i f)(\partial_i g)) \\ &= (x_{i+1} \partial_i f)g + (s_i f)(x_{i+1} \partial_i g) \\ &= (\theta_i f)g + (s_i f)(\theta_i g). \end{aligned}$$

By item 1. of Proposition 2.1,

$$\begin{aligned} & \pi_i(fg) \\ &= (1 + \theta_i)(fg) \\ &= fg + \theta_i(fg) \\ &= fg + (\theta_i f)g + (s_i f)(\theta_i g) \\ &= ((1 + \theta_i)f)g + (s_i f)(\theta_i g) \\ &= (\pi_i f)g + (s_i f)(\theta_i g). \end{aligned}$$

□

2.2 Semi-Standard Augmented filling

Let \mathbb{N} (or \mathbb{Z}^+) be the set of all positive integers and $\mathbb{Z}_{\geq 0}$ be the set of non-negative integers. Also we denote $\epsilon_k = 12 \cdots k = id$ as the identity element (we write permutations in one line notation). For $n \in \mathbb{Z}_{\geq 0}$ and $k \in \mathbb{N}$, we say $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k) \in (\mathbb{Z}_{\geq 0})^k$ is a weak composition n (denoted as $\alpha \vDash n$) with k parts if $\sum_{i=1}^k \alpha_i = n$ and write $l(\alpha) = k$ to denote the length (the number of parts) of α . Furthermore, if $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_k \geq 0$, we call α a partition of n with k parts and write $\alpha \vdash n$ (usually we denote $l(\alpha) = \max\{i : \alpha_i > 0\}$ for α being a partition). We use $\text{Par}(n)$ to denote the set of all partitions of a nonnegative integer n and Par to denote the set of all partitions. For a weak composition α with k parts, define $x^\alpha := x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_k^{\alpha_k}$. We call x^α a dominating monomial if α is a partition.

We denote $\bar{\alpha}$ as the *reverse* of α , that is, $\bar{\alpha} = (\alpha_k, \dots, \alpha_1)$. (Note that in [4], they use α^* instead of $\bar{\alpha}$.) Similarly, we write \bar{X} as the *reverse* of X for any finite string of alphabets X . For example, $\overline{cacdba} = abdcac$ and $\overline{14D9c7} = 7c9D41$.

Define ω_α as the permutation of minimal length such that


$$\omega_\alpha(\alpha) := (\alpha_{\omega_\alpha(1)}, \alpha_{\omega_\alpha(2)}, \dots, \alpha_{\omega_\alpha(k)})$$

is a partition.


Given two weak compositions α and β , we write $\beta \geq \alpha$ if and only if $\omega_\beta \leq \omega_\alpha$ in the strong Bruhat order.

Let α be a weak composition. The augmented diagram of shape α is the figure with $|\alpha| + l(\alpha)$ cells (or boxes) where column i has $\alpha_i + 1$ cells. The bottom row is called the *basement* of the augmented diagram.

For example, if $\alpha = (1, 0, 1, 0, 0, 4, 0, 6, 5)$, then the augmented diagram of α is



Also we impose an order, called the *reading order*, on the cells of the diagram which starts from left to right, top to bottom. So the order of the above diagram is:



where the number in each cell represents the order of that cell in reading order.

A filling of an augmented diagram is an assignment of a positive integer to each cell in the diagram.

From now on we only consider fillings whose entries in each column are weakly decreasing from the bottom to the top.

For any two columns (including the basement cells) i and j with $i < j$, we pick three cells X, Y and Z , where cell X is immediately above cell Y in the ‘taller’ column k , where


$$k = \begin{cases} i & \text{if } \alpha_i \geq \alpha_j \\ j & \text{if } \alpha_i < \alpha_j \end{cases},$$

and cell Z from the ‘shorter’ column to form a *triple* (X, Y, Z) in the following way:

$$\begin{cases} \text{Type A triple: cell } Z \text{ is in the same row as cell } X & \text{if } \alpha_i \geq \alpha_j \\ \text{Type B triple: cell } Z \text{ is in the same row as cell } Y & \text{if } \alpha_i < \alpha_j \end{cases}.$$


Here are some examples of triples:

(The first two are type A triples and the last two are type B triples.)



We say (X, Y, Z) forms a *coinversion triple* if the filling F of the diagram assigns each cell in the triple a positive integer, say $F(X), F(Y), F(Z)$ respectively, in such a way that $F(X) \leq F(Z) \leq F(Y)$. Otherwise we call (X, Y, Z) an *inversion triple*.

For instance in the following examples, the second and the third ones are coinversion triples while the first and the last one are inversion triples.




Definition 2.1. A semi-standard augmented filling (SSAF) of an augmented diagram with shape being a weak composition $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$ is a filling satisfying:

1. the basement entries form a permutation σ (in one line notation) of $\{1, \dots, k\}$, i.e. $\sigma \in S_k$;
2. every (Type A or B) triple is an inversion triple.

We denote $SSAF(\sigma, \alpha)$ the set of all SSAF of an augmented diagram of shape $\alpha = (\alpha_1, \dots, \alpha_k)$ with basement entries (from left to right) being $\sigma \in S_k$ (i.e. basement of column i has entry $\sigma(i)$).

Example 4. The following SSAFs are all the elements in the set $SSAF(4132, 1032)$:



Given an SSAF F with basement of length n for some $n \in \mathbb{N}$, define the weight of F as $x^F := \prod_{i=1}^n x_i^{m_i(F)-1}$, where $m_i(F)$ is the number of i appearing in F for $1 \leq i \leq n$.

Example 5. $x^F = x_1^2 x_2^2 x_3^2$ for $F =$

2.3 Demazure atoms and Demazure characters

Definition 2.2. A Demazure atom of shape α , where α is a weak composition, is defined as

$$\mathcal{A}_\alpha := \sum_{F \in \text{SSAF}(\epsilon_k, \alpha)} x^F.$$

Definition 2.3. A Demazure character (key polynomial, or key) of shape α , where α is a weak composition, is defined as

$$\kappa_{\bar{\alpha}} := \sum_{F \in \text{SSAF}(\bar{\epsilon}_k, \alpha)} x^F.$$

Remark: See [12] for further discussion on key polynomials.

The following theorem gives different equivalent definitions of Demazure atoms (\mathcal{A}_α) and Demazure characters ($\kappa_{\bar{\alpha}}$) of shape α and is proved in [2, 6, 12].

Theorem 2.7. For $k \in \mathbb{N}$ and composition α with $l(\alpha) = k$,

$$E_{\bar{\alpha}}(x_k, \dots, x_1; \infty, \infty) = \sum_{F \in \text{SSAF}(\epsilon_k, \alpha)} x^F = \theta_{\omega_\alpha^{-1}} x^{\omega_\alpha(\alpha)}$$

$$E_{\bar{\alpha}}(x_1, \dots, x_k; 0, 0) = \sum_{F \in \text{SSAF}(\bar{\epsilon}_k, \alpha)} x^F = \pi_{\bar{\epsilon}_k \omega_\alpha^{-1}} x^{\omega_\alpha(\alpha)}$$

where $E_{\bar{\alpha}}(x_k, \dots, x_1; \infty, \infty)$ and $E_{\bar{\alpha}}(x_1, \dots, x_k; 0, 0)$ are the nonsymmetric Macdonald polynomials of shape $\bar{\alpha}$ with $q = t = \infty$ and $X = (x_k, \dots, x_1)$ and with $q = t = 0$ and $X = (x_1, \dots, x_k)$ respectively.

Example 6. Let $\alpha = (1, 0, 3)$. Then $\omega_\alpha = 231 = (12)(23) = s_1 s_2$.

$$\text{Hence } \mathcal{A}_{(1,0,3)} = \theta_2 \theta_1 x_1^3 x_2 = \theta_2 (x_1^2 x_2^2 + x_1 x_2^3) = x_1^2 x_2 x_3 + x_1^2 x_3^2 + x_1 x_2^2 x_3 + x_1 x_2 x_3^2 + x_1 x_3^3.$$

The following are all the SSAFs of $\text{SSAF}(123, (1, 0, 3))$:

$\begin{array}{ c } \hline 3 \\ \hline 3 \\ \hline 3 \\ \hline \end{array}$	$\begin{array}{ c } \hline 2 \\ \hline 3 \\ \hline 3 \\ \hline \end{array}$	$\begin{array}{ c } \hline 1 \\ \hline 3 \\ \hline 3 \\ \hline \end{array}$	$\begin{array}{ c } \hline 2 \\ \hline 2 \\ \hline 3 \\ \hline \end{array}$	$\begin{array}{ c } \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline \end{array}$
1 2 3	1 2 3	1 2 3	1 2 3	1 2 3
weights : $x_1 x_3^3$	$x_1 x_2 x_3^2$	$x_1^2 x_3^2$	$x_1 x_2^2 x_3$	$x_1^2 x_2 x_3$

As a result, the sum of all weights of the SSAFs gives $\mathcal{A}_{(1,0,3)}$.

Since $\bar{\epsilon}_3 \omega_\alpha^{-1} = (s_2 s_1 s_2)(s_2 s_1) = s_2$, we have $\kappa_{(3,0,1)} = \pi_{\bar{\epsilon}_3 \omega_\alpha^{-1}}(x_1^3 x_2) = \pi_2(x_1^3 x_2) = x_1^3 x_2 + x_1^2 x_3$.

The following are all the SSAFs of $SSAF(321, (1, 0, 3))$:

$$\begin{array}{cc}
 \begin{array}{|c|} \hline 1 \\ \hline \end{array} & \begin{array}{|c|} \hline 1 \\ \hline \end{array} \\
 \begin{array}{|c|} \hline 1 \\ \hline \end{array} & \begin{array}{|c|} \hline 1 \\ \hline \end{array} \\
 \begin{array}{|c|} \hline 3 \\ \hline \end{array} & \begin{array}{|c|} \hline 2 \\ \hline \end{array} & \begin{array}{|c|} \hline 1 \\ \hline \end{array} & \begin{array}{|c|} \hline 1 \\ \hline \end{array} \\
 3 & 2 & 1 & 3 & 2 & 1 \\
 \text{weights : } & x_1^3 x_3 & & x_1^3 x_2 & &
 \end{array}$$

As a result, the sum of all weights of the SSAFs gives $\kappa_{(3,0,1)}$.

The following theorem gives some properties of Demazure atoms and characters.

Theorem 2.8.

1. A key polynomial is a positive sum of Demazure atoms. In fact,

$$\kappa_{\bar{\alpha}} = \sum_{\beta \geq \alpha} \mathcal{A}_{\beta}.$$

2. A key polynomial with a partition shape λ , with $l(\lambda) = k$, is the Schur polynomial s_{λ} , i.e.,

$$\kappa_{\bar{\lambda}} = s_{\lambda}(x_1, \dots, x_k).$$

3. The set of all Demazure atoms $\{\mathcal{A}_{\gamma} : \gamma \vDash n, n \in \mathbb{Z}_{\geq 0}\}$ forms a basis for the polynomial ring, and so does the set of all key polynomials $\{\kappa_{\gamma} : \gamma \vDash n, n \in \mathbb{Z}_{\geq 0}\}$.

Proof. Item 1 follows directly from Theorem 2.5. We can describe combinatorially how to get the atoms from the key (a particular case of Proposition 6.1 in [12]):

Consider a filling $F \in SSAF(\bar{\epsilon}_k, \alpha)$ and an empty filling G_0 with basement ϵ_k . Consider the entries of the first row (from the bottom above the basement) of F , namely $a_{11} < a_{12} < \dots < a_{1r_1}$ where r_1 is the length of the first row. Create the first row of G_0 by placing a_{1i} in the cell right above a_i in the basement of G_0 (that is, a_{1i} is placed in the first row above the basement and also in the a_{1i}^{th} column of G) for $1 \leq i \leq r_1$. We call the new filling G_1

Now consider the entries $a_{21} < a_{22} < \dots < a_{2r_2}$ of the second row of F where r_2 is the length of the second row of F . Search in the top row of G_1 for the leftmost number not less than a_{2r_2} and place a_{2r_2} in the cell right above it. Then search for the leftmost available number (i.e. not chosen yet) in the top row of G_1 not less than a_{2,r_2-1} and place a_{2,r_2-1} in the cell right above it, and so on until a_{21} is placed. We now get a new filling with 2 rows above the basement and call it G_2 .

By repeating the same process until all entries of F are placed and we get a filling G_r with basement ϵ_k with shape less than or equal to α , where r is the number of rows in F .

Item 2 follows from Theorem 2.7 as $s_\lambda = E_\lambda(x_1, \dots, x_k; 0, 0)$ ([2]). It is also proved in [6]. A combinatorial proof can be found in Theorem 4.1 in [3] which uses the insertion algorithm discussed in [11, 12].

Item 3 also follows from Theorem 2.7 as $\{E_\alpha(x; q, t) : \alpha \vDash n, n \in \mathbb{Z}_{\geq 0}\}$ forms a basis for the polynomial ring over $\mathbb{Q}(q, t)$. Again, it is also proved in [6].

□

Chapter 3

Decomposition of products of Demazure atoms and characters

In this chapter, we study the decomposition of the products of Demazure atoms and characters with respect to the atom-basis $\{\mathcal{A}_\gamma : \gamma \vDash n, n \in \mathbb{Z}_{\geq 0}\}$ and key-basis $\{\kappa_\gamma : \gamma \vDash n, n \in \mathbb{Z}_{\geq 0}\}$.

Let λ, μ be partitions and α, β be weak compositions. Let $+_{\mathcal{A}}$ and $+_{\kappa}$ denote the property of being able to be decomposed into a positive sum of atoms and keys respectively. Note that by item 1 in Theorem 2.8, $+_{\kappa}$ implies $+_{\mathcal{A}}$. Otherwise, we put an \times in the cell. For example, a partition (μ) -shaped atom times a key of any shape (α) is key positive and hence we put $+_{\kappa}$ in the corresponding box.

		Atoms		Keys	
		λ	α	λ	α
Atoms	μ	$+_{\mathcal{A}}$	$+_{\mathcal{A}} \textcircled{1}$	$+_{\mathcal{A}}$	$+_{\kappa} \textcircled{2}$
	β		\times	$+_{\mathcal{A}}$	\times
Keys	μ			$+_{\kappa}$	$+_{\kappa}$
	β				open $\textcircled{3}$

Table 3.1: Decomposition of products of atoms and keys into atoms

The positive results in the table can be found in [3], except for the cells marked $\textcircled{1}$, $\textcircled{2}$

and $\textcircled{3}$.

We will prove $\textcircled{1}$ (which was previously open) in this chapter using words and insertion algorithm introduced in [3, 12]:

Theorem 3.1. *The product $\mathcal{A}_\mu \cdot \mathcal{A}_\alpha$ is atom-positive for any partition μ and weak composition α .*

The coefficients in the decomposition into atoms are actually counting the number of ways to insert words arising from an SSAF of shape α into an SSAF of shape μ and we will discuss properties of words and how to record different ways of insertion in Section 3.1 and Section 3.2. Also note that the product in the theorem is not key positive. A simple counter example would be just putting μ as the empty partition, that is, with all entries 0 and $\alpha = (0, 1)$ and hence $\mathcal{A}_\mu \cdot \mathcal{A}_\alpha = \mathcal{A}_\alpha = \theta_1(x_1) = (\pi_1 - id)(x_1) = \kappa_{(0,1)} - \kappa_{(1,0)}$.

$\textcircled{2}$ is proved in [5] (the proof involves crystals but does not involve SSAF). Both results $\textcircled{1}$ and $\textcircled{2}$ imply $+\mathcal{A}$ for the $\mathcal{A}_\mu \cdot \kappa_{\bar{\alpha}}$ cell. We will apply the bijection in the proof of Theorem 6.1 in [3] to Theorem 3.1 to give a tableau-combinatorial proof of $\textcircled{2}$ in Section 3.4.

As for the product of two keys of arbitrary shapes, that is, the cell marked with $\textcircled{3}$, there are examples showing that such a product is not a positive sum of keys. For example, $\kappa_{(0,2)} \cdot \kappa_{(1,0,2)} = \kappa_{(1,2,2)} + \kappa_{(1,3,1)} + \kappa_{(1,4,0)} + \kappa_{(2,3,0)} + \kappa_{(3,0,2)} - \kappa_{(3,2,0)} + \kappa_{(4,0,1)} - \kappa_{(4,1,0)}$. Thus it remains to check whether it is a positive sum of atoms, which is still open. Hence $\textcircled{3}$ gives the following conjecture (first appearing in an unpublished work of Victor Reiner and Mark Shimozono).

Conjecture 1. *Let α, β be weak compositions. Then the product of the key polynomials of shape $\bar{\alpha}$ and $\bar{\beta}$ can be written as a positive sum of atoms, i.e.,*

$$\kappa_\alpha \cdot \kappa_\beta = \sum_{\gamma \models |\alpha| + |\beta|} c_{\alpha\beta}^\gamma \mathcal{A}_\gamma$$

for some nonnegative integers $c_{\alpha\beta}^\gamma$.

We will verify Conjecture 1 for $l(\alpha), l(\beta) \leq 3$ in Chapter 4.

3.1 Convert a column word to a row word

Definition 3.1. A **word** is a sequence of positive integers.

Definition 3.2. Let $a, b, c \in \mathbb{N}$ and u, v be some fixed (can be empty) words. Define twisted Knuth relation \smile^* by:

1. $ubacv \smile^* ubcav$ if $c \leq b < a$
2. $uacbv \smile^* ucabv$ if $c < b \leq a$.

Then we say two words w and w' are twisted Knuth equivalent if w can be transformed to w' by repeated use of 1. and 2. and we write $w \smile^* w'$.

Definition 3.3. A word w is a **column word** if it can be broken down into k weakly decreasing subsequences of weakly decreasing lengths

$$w = a_{11} \dots a_{1c_1} | a_{21} \dots a_{2c_2} | \dots | a_{k1} \dots a_{kc_k}$$

where $c_1 \geq c_2 \geq \dots \geq c_k > 0$, $c_1, \dots, c_k \in \mathbb{N}$

$$\text{such that } \begin{cases} a_{ij} \geq a_{i,j+1} & \forall 1 \leq j < c_i, 1 \leq i \leq k \\ a_{i+1,c_{i+1}-j} > a_{i,c_i-j} & \forall 0 \leq j < c_i, 1 \leq i < k \end{cases}.$$

Definition 3.4. A word w is a **row word** if it can be broken down into k strictly increasing subsequences of weakly decreasing lengths

$$w = a_{11} \dots a_{1r_1} | a_{21} \dots a_{2r_2} | \dots | a_{k1} \dots a_{kr_k}$$

where $r_1 \geq r_2 \geq \dots \geq r_k > 0$, $r_1, \dots, r_k \in \mathbb{N}$

$$\text{such that } \begin{cases} a_{ij} < a_{i,j+1} & \forall 1 \leq j < r_i, 1 \leq i \leq k \\ a_{i+1,r_{i+1}-j} \leq a_{i,c_i-j} & \forall 0 \leq j < r_i, 1 \leq i < k \end{cases}.$$

Given an SSAF, one can get its column word by using the algorithm described in [3], while one can get its row word (which is the reverse reading word defined in [4] by reading the entries of each

row in ascending order, starting from the bottom row to the top row. We call a word a column (resp. row) word because when we insert each subsequence of the word using the insertion in [11], a new column (resp. row) will be created.

Example 7. $886531|97643|9764|5|6$ is a column word whose corresponding SSAF is:

							6	
							6	4
					3		7	5
					4		7	6
					5		8	9
1		3			6		8	9
1	2	3	4	5	6	7	8	9


and the corresponding row word is $13689|589|467|357|46|6$.

We describe an insertion algorithm for inserting an integer $c \leq k$ into an SSAF with basement ϵ_k . A detailed description can be found in [11].

Given an SSAF F with basement ϵ_k and an integer c to be inserted, we write $c \rightarrow F$ or $F \leftarrow c$ to represent the insertion of c into F (and similarly we denote $c_2c_1 \rightarrow F$ and $F \leftarrow c_1c_2$ as first inserting c_1 to F and then insert c_2 , and so on).

To insert c , we first find the cell A in F with the smallest order, say m , such that $F(A') < c \leq F(A)$, where A' is the cell immediately above A if it exists and assign $F(A') = 0$ if A is the top cell of a column and just treat A' as an empty cell to be filled in. If cell A is the top cell of a column, then we create a new cell immediately above A , i.e. A' , and assign c to the new cell and we are done. Otherwise, we replace the entry $y = F(A')$ by c and now insert y as we do for c , but now finding a cell B of the smallest order **larger than** m such that $F(B') < y \leq F(B)$ as treating B as A in the previous step. Repeat the process until a new cell is finally created.

Example 8. Let F be the SSAF in Example 7 where $k = 9$. We find the new SSAF created by inserting 7, i.e. $7 \rightarrow F$, as follows:



The green cell represents the cell A in each step. The yellow cell represents the cell A' whose value is changed after the insertion step (i.e. the original entry is bumped out by the number being inserted). The white cells are the cells under consideration in each step (i.e. candidates for the position of B , that is, the new A).

Lemma 3.2. Let $u := a_1 a_2 \dots a_n | b$ and $v := a_1 a_2 \dots a_n | b | c$ be two column words, where $c > b > a_n$, $n \in \mathbb{N}$. Then


1. $u \rightsquigarrow^* b' a'_1 \dots a'_n$ where $b' < a'_1$, $a'_1 \geq a'_2 \geq \dots \geq a'_n$, and $b' := a_t$ where $t = \min\{j : b > a_j\}$
2. $v \rightsquigarrow^* b' c' a''_1 a''_2 \dots a''_n$ where b' is defined as in 1., $b' < c' < a''_1$ and $a''_1 \geq a''_2 \geq \dots \geq a''_n$.

Proof. One can check that for $t = \min\{j : b > a_j\}$,

$$u \rightsquigarrow^* \begin{cases} a_1 b a_2 \dots a_n & \text{if } t = 1 \\ a_t a_1 \dots a_{t-1} b a_{t+1} \dots a_n & \text{if } 1 < t < n \text{ and 1. follows. Also note that } b' = a_t. \\ a_n a_1 \dots a_{n-1} b & \text{if } t = n \end{cases}$$

Indeed, we can visualize by applying the insertion in [11]. When we insert the first n integers of u , we get a column above the basement entry a_1 :


If $b > a_1$, we have



giving the row word $a_1 b a_2 \dots a_n$.


If $b \leq a_1$, then for $t = \min\{j : b > a_j\}$, we have:

For $t < n$




giving the row word $a_t a_1 \dots a_{t-1} b a_{t+1} \dots a_n$.

For $t = n$



giving the row word $a_n a_1 \dots a_{n-1} b$.


Since inserting v to an empty atom is the same as inserting c to the atom created by inserting u to an empty atom, we can inset c to the tableaux above and get the following:



giving the row word $a_1 b c a_2 \dots a_n$.


If $b \leq a_1$ and $a_{t-1} \geq c > b > a_t$, we have:

For $t < n$



giving the row word $a_t b a_1 \dots a_{t-1} c a_{t+1} \dots a_n$.

For $t = n$




giving the row word $a_n b a_1 \dots a_{n-1} c$.

If $b \leq a_1$ and $c > a_{t-1}$, we have:


For $t < n$, let $t' = \min\{j : c > a_j, 1 \leq j \leq t-1\}$, we have:

For $t' = 1$,




giving the row word $a_t a_1 c a_2 \dots a_{t-1} b a_{t+1} \dots a_n$.

For $1 < t' < t-1$,



giving the row word $a_t a_{t'} a_1 \dots a_{t'-1} c a_{t'+1} \dots a_{t-1} b a_{t+1} \dots a_n$.


For $t' = t - 1$,



giving the row word $a_t a_{t-1} a_1 \dots a_{t-2} c b a_{t+1} \dots a_n$.


For $t = n$, let $t' = \min\{j : c > a_j\}$, we have:

For $t' = 1$,




giving the row word $a_n a_1 c a_2 \dots a_{n-1} b$.

For $1 < t' < n - 1$,



giving the row word $a_n a_{t'} a_1 a_2 \dots a_{t'-1} c a_{t'+1} \dots a_{n-1} b$.

For $t' = n - 1$,



giving the row word $a_n a_{n-1} a_1 a_2 \dots a_{n-2} c b$.

Notice that for all the cases of inserting c , the first row has the least entry unaffected, meaning

that the row word has the same first entry after inserting c . The result follows by setting the row word of the tableau of inserting v into an empty atom as $b'c'a''_1 \dots a''_n$. \square

Example 9. *One can also prove Lemma 3.2 by repeated use of 1. and 2. in Definition 3.2, for example:*

$9764|5 \smile^* 97465 \smile^* 94765 \smile^* 4|9765$ by using 2. repeatedly, and similarly, we have $9764|5|6 \smile^* 4|9765|6 \smile^* 45|9766$.

Using SSAF, it means

$$65 \rightarrow \begin{array}{ccccccc} & & & & 4 & & \\ & & & & 6 & & \\ & & & & 7 & & \\ & & & & 9 & & \\ 1 & 2 & 3 & \dots & 8 & 9 & \end{array} \Rightarrow 6 \rightarrow \begin{array}{ccccccc} & & & & 5 & & \\ & & & & 6 & & \\ & & & & 7 & & \\ & & & 4 & 9 & & \\ 1 & 2 & 3 & 4 & \dots & 8 & 9 \end{array} \Rightarrow \begin{array}{ccccccc} & & & & 6 & & \\ & & & & 6 & & \\ & & & & 7 & & \\ & & & 4 & 5 & & 9 \\ 1 & \dots & 4 & 5 & \dots & 8 & 9 \end{array}$$

where $65 \rightarrow$ means 5 is inserted before 6.

Lemma 3.3. *Let $k \geq 2$ be an integer and i_1, \dots, i_k be nonnegative integers. Let*

$$w_0 = a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} | a_{21}^{(i_2)} \dots a_{2c_2}^{(i_2)} | \dots | a_{k1}^{(i_k)} \dots a_{kc_k}^{(i_k)}$$

be a column word. Then there exist $a_{rs}^{(i_r+1)}$ where $1 \leq s \leq c_r$, $1 \leq r < k$ and $a_{k1}^{(i_k+m)}$ where $1 \leq m < k$ satisfying

$$\left\{ \begin{array}{l} a_{k-j-1, c_{k-j-1}}^{(i_{k-j-1})} < a_{k1}^{(i_k+j)} < a_{k-j,1}^{(i_{k-j+1})} \quad \forall 1 \leq j < k-1 \quad (k > 3) \\ a_{k1}^{(i_k+k-1)} < a_{11}^{(i_1+1)} \\ a_{1,1}^{(i_1+1)} \dots a_{1,c_1}^{(i_1+1)} | \dots | a_{k-1,1}^{(i_{k-1}+1)} \dots a_{k-1,c_{k-1}}^{(i_{k-1}+1)} | a_{k2}^{(i_k)} \dots a_{kc_k}^{(i_k)} \text{ is a column word} \end{array} \right.$$

such that $w_0 \smile^* w_1 \smile^* \dots \smile^* w_{k-1}$, where

$$\left\{ \begin{array}{l} w_j := a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} \dots a_{k-j-1,1}^{(i_{k-j-1})} \dots a_{k-j-1,c_{k-j-1}}^{(i_{k-j-1})} a_{k1}^{(i_k+j)} a_{k-j,1}^{(i_{k-j+1})} \dots a_{k-j,c_{k-j}}^{(i_{k-j+1})} \dots \\ \dots a_{k-1,1}^{(i_{k-1+1})} \dots a_{k-1,c_{k-1}}^{(i_{k-1+1})} a_{k2}^{(i_k)} \dots a_{kc_k}^{(i_k)} \quad \forall 1 \leq j < k-1, \\ w_{k-1} := a_{k1}^{i_k+k-1} a_{11}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} \dots a_{k-1,1}^{(i_{k-1+1})} \dots a_{k-1,c_{k-1}}^{(i_{k-1+1})} a_{k2}^{(i_k)} \dots a_{kc_k}^{(i_k)}. \end{array} \right.$$

Proof. We prove this by induction on k .

When $k = 2$, we have $w_0 := a_{11}^{(i_1)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}$.

Let $t_{12} = \min_{1 \leq j \leq c_1} \{j : a_{21}^{(i_2)} > a_{ij}^{(i_1)}\}$.

By applying the proof of 1. in Lemma 3.2 on $a_{11}^{(i_1)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)}$, we have

$$\begin{aligned} w_0 \smile^* w_1 &:= a_{21}^{(i_2+1)} a_{11}^{(i_1+1)} a_{12}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)} \\ &:= \begin{cases} a_{11}^{(i_1)} a_{21}^{(i_2)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)} & \text{if } t_{12} = 1 \\ a_{t_{12}}^{(i_1)} a_{11}^{(i_1)} \dots a_{1,t_{12}-1}^{(i_1)} a_{21}^{(i_2)} a_{1,t_{12}+1}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)} & \text{if } t_{12} > 1 \end{cases} \end{aligned}$$

where $a_{21}^{(i_2+1)} < a_{11}^{(i_1+1)}$ and $a_{11}^{(i_1+1)} \geq \dots \geq a_{1c_1}^{(i_1+1)}$. Also note that $a_{21}^{(i_2+1)} = a_{1t_{12}}^{(i_1)}$.

For $t_{12} = 1$, we know that $a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}$ is a column word and hence

$$a_{12}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}.$$

For $t_{12} > 1$, if $c_2 - (c_1 - t_{12} + 1) > 0$, then

$$a_{21}^{(i_2)} < a_{1,t_{12}-1}^{(i_1)} = a_{1,c_1-(c_1-t_{12}+1)}^{(i_1)} < a_{2,c_2-(c_1-t_{12}+1)}^{(i_2)} \quad (\text{by Definition 3.3})$$

which leads to a contradiction as $a_{21}^{(i_2)} \geq a_{2j}^{(i_2)}$ for $1 \leq j \leq c_2$.

As a result, we have $c_2 - (c_1 - t_{12} + 1) \leq 0$ which implies $c_2 - 1 \leq c_1 - t_{12}$.

Thus $a_{1,t_{12}+1}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}$ is a column word, and hence

$$a_{12}^{(i_1)} \dots a_{1,t_{12}-1}^{(i_1)} a_{21}^{(i_2)} a_{1,t_{12}+1}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)} \text{ is also a column word.}$$

Therefore $a_{12}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}$ is a column word.

Note that $a_{11}^{(i_1+1)} \geq a_{12}^{(i_1+1)}$, we can conclude $a_{11}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}$ is a column word.

Hence the statement is true for $k = 2$.

For $k = 3$, we have $w_0 := a_{11}^{(i_1)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)} a_{31}^{(i_3)} a_{32}^{(i_3)} \dots a_{3c_3}^{(i_3)}$.

Consider $a_{21}^{(i_2)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)} a_{31}^{(i_3)} a_{32}^{(i_3)} \dots a_{3c_3}^{(i_3)}$, by $k = 2$ case, we know

$$a_{21}^{(i_2)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)} a_{31}^{(i_3)} a_{32}^{(i_3)} \dots a_{3c_3}^{(i_3)} \smile^* a_{31}^{(i_3+1)} a_{21}^{(i_2+1)} a_{22}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)} a_{32}^{(i_3)} \dots a_{3c_3}^{(i_3)}$$

where $a_{31}^{(i_3+1)} < a_{21}^{(i_2+1)}$ and $a_{21}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)} a_{32}^{(i_3)} \dots a_{3c_3}^{(i_3)}$ is a column word. Also, $a_{31}^{(i_3+1)} = a_{2t_{23}}^{(i_2)} \geq a_{2c_2}^{(i_2)} > a_{1c_1}^{(i_1)}$, $t_{23} := \min_{1 \leq j \leq c_2} \{j : a_{31}^{(i_3+1)} > a_{2j}^{(i_2)}\}$.

Hence, $a_{1c_1}^{(i_1)} < a_{31}^{(i_3+1)} < a_{21}^{(i_2+1)}$.

Furthermore, $a_{11}^{(i_1)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{31}^{(i_3+1)}$ is also a column word and by 1. in Lemma 3.2,

$$a_{11}^{(i_1)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{31}^{(i_3+1)} \smile^* a_{31}^{(i_3+2)} a_{11}^{(i_1+1)} a_{12}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)}$$

where $a_{31}^{(i_3+2)} < a_{11}^{(i_1+1)}$ and $a_{11}^{(i_1+1)} a_{12}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)}$ is a column word for $a_{11}^{(i_1+1)} \geq a_{12}^{(i_1+1)} \geq \dots \geq a_{1c_1}^{(i_1+1)}$. Also $a_{31}^{(i_3+2)} = a_{1t_{13}}^{(i_1)}$ where $t_{13} := \min_{1 \leq j \leq c_1} \{j : a_{31}^{(i_3+2)} > a_{1j}^{(i_1)}\}$.

It remains to check that $a_{11}^{(i_1+1)} a_{12}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{21}^{(i_2+1)} a_{22}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)}$ is a column word.

First note that by 1. in Lemma 3.2,

$$\begin{aligned} a_{21}^{(i_2+1)} a_{22}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)} &= \begin{cases} a_{31}^{(i_3)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)} & \text{if } t_{23} = 1 \\ a_{21}^{(i_2)} \dots a_{2,t_{23}-1}^{(i_2)} a_{31}^{(i_3)} a_{2,t_{23}+1}^{(i_2)} \dots a_{2c_2}^{(i_2)} & \text{if } t_{23} > 1 \end{cases}, \\ a_{11}^{(i_1+1)} a_{12}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} &= \begin{cases} a_{31}^{(i_3+1)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} & \text{if } t_{13} = 1 \\ a_{11}^{(i_1)} \dots a_{1,t_{13}-1}^{(i_1)} a_{31}^{(i_3+1)} a_{1,t_{13}+1}^{(i_1)} \dots a_{1c_1}^{(i_1)} & \text{if } t_{13} > 1 \end{cases} \\ &= \begin{cases} a_{2t_{23}}^{(i_2)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} & \text{if } t_{13} = 1 \\ a_{11}^{(i_1)} \dots a_{1,t_{13}-1}^{(i_1)} a_{2t_{23}}^{(i_2)} a_{1,t_{13}+1}^{(i_1)} \dots a_{1c_1}^{(i_1)} & \text{if } t_{13} > 1 \end{cases}. \end{aligned}$$

By definition of a column word, we have $a_{2t_{23}}^{(i_2)} > a_{1,c_1-(c_2-t_{23})}^{(i_1)}$. By definition of t_{13} , we have $t_{13} \leq c_1 - (c_2 - t_{23})$ which implies $c_2 - t_{23} \leq c_1 - t_{13}$.

Combining, we have the following four cases:

(i) $t_{13} = t_{23} = 1$:

$$a_{11}^{(i_1+1)} a_{12}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{21}^{(i_2+1)} a_{22}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)} = a_{21}^{(i_2)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{31}^{(i_3)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}.$$

Since $a_{31}^{(i_3)} > a_{21}^{(i_2)} = a_{11}^{(i_1+1)} \geq a_{1j}^{(i_1+1)} = a_{1j}^{(i_1)}$ for $1 < j \leq c_1$,

and $a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}$ is a column word, we can conclude that

$$a_{21}^{(i_2)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{31}^{(i_3)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}$$
 is also a column word.

(ii) $t_{13} = 1, t_{23} > 1$:

$$\begin{aligned} & a_{11}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{21}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)} \\ &= a_{2t_{23}}^{(i_2)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \dots a_{2,t_{23}-1}^{(i_2)} a_{31}^{(i_3)} a_{2,t_{23}+1}^{(i_2)} \dots a_{2c_2}^{(i_2)}. \end{aligned}$$

Since $a_{31}^{(i_3)} > a_{2t_{23}}^{(i_2)}$ and $a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{22}^{(i_2)} \dots a_{2,t_{23}-1}^{(i_2)} a_{2t_{23}}^{(i_2)} a_{2,t_{23}+1}^{(i_2)} \dots a_{2c_2}^{(i_2)}$ is a column word, we

know $a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{22}^{(i_2)} \dots a_{2,t_{23}-1}^{(i_2)} a_{31}^{(i_3)} a_{2,t_{23}+1}^{(i_2)} \dots a_{2c_2}^{(i_2)}$ is a column word.

Also $a_{21}^{(i_2)} \geq a_{2,t_{23}-1}^{(i_2)} \geq a_{31}^{(i_3)} > a_{2t_{23}}^{(i_2)} = a_{11}^{(i_1+1)} \geq a_{1j}^{(i_1+1)} = a_{1j}^{(i_1)}$ for $1 < j \leq c_1$,

thus $a_{2t_{23}}^{(i_2)} a_{12}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \dots a_{2,t_{23}-1}^{(i_2)} a_{31}^{(i_3)} a_{2,t_{23}+1}^{(i_2)} \dots a_{2c_2}^{(i_2)}$ is a column word.

(iii) $t_{13} > 1, t_{23} = 1$:

$$\begin{aligned} & a_{11}^{(i_1+1)} a_{12}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{21}^{(i_2+1)} a_{22}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)} \\ &= a_{11}^{(i_1)} \dots a_{1,t_{13}-1}^{(i_1)} a_{21}^{(i_2)} a_{1,t_{13}+1}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{31}^{(i_3)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}. \end{aligned}$$

Since $c_2 - 1 \leq c_1 - t_{13}$ which implies $c_2 \leq c_1 - (t_{13} - 1)$, we just need to consider

$$a_{21}^{(i_2)} a_{1,t_{13}+1}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{31}^{(i_3)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}$$

to conclude that

$$a_{11}^{(i_1)} \dots a_{1,t_{13}-1}^{(i_1)} a_{21}^{(i_2)} a_{1,t_{13}+1}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{31}^{(i_3)} a_{22}^{(i_2)} \dots a_{2c_2}^{(i_2)}$$

is a column word.

As $c_2 - 1 \leq c_1 - t_{13}$, $a_{1,t_{13}+1}^{(i_1)} \cdots a_{1c_1}^{(i_1)} a_{22}^{(i_2)} \cdots a_{2c_2}^{(i_2)}$ is a column word. Also $a_{31}^{(i_3)} > a_{21}^{(i_2)}$ for $t_{23} = 1$, and $a_{21}^{(i_2)} = a_{1t_{13}}^{(i_1+1)} \geq a_{1j}^{(i_1+1)} = a_{1j}^{(i_1)}$ for $t_{13} < j \leq c_1$ and hence $a_{21}^{(i_2)} a_{1,t_{13}+1}^{(i_1)} \cdots a_{1c_1}^{(i_1)} a_{31}^{(i_3)} a_{22}^{(i_2)} \cdots a_{2c_2}^{(i_2)}$ is a column word and result follows.

(iv) $t_{13} > 1, t_{23} > 1$:

$$\begin{aligned} & a_{11}^{(i_1+1)} a_{12}^{(i_1+1)} \cdots a_{1c_1}^{(i_1+1)} a_{21}^{(i_2+1)} a_{22}^{(i_2+1)} \cdots a_{2c_2}^{(i_2+1)} \\ &= a_{11}^{(i_1)} \cdots a_{1,t_{13}-1}^{(i_1)} a_{2t_{23}}^{(i_2)} a_{1,t_{13}+1}^{(i_1)} \cdots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \cdots a_{2,t_{23}-1}^{(i_2)} a_{31}^{(i_3)} a_{2,t_{23}+1}^{(i_2)} \cdots a_{2c_2}^{(i_2)}. \end{aligned}$$

Since $a_{11}^{(i_1)} \cdots a_{1,t_{13}-1}^{(i_1)} a_{1t_{13}}^{(i_1)} a_{1,t_{13}+1}^{(i_1)} \cdots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \cdots a_{2,t_{23}-1}^{(i_2)} a_{2t_{23}}^{(i_2)} a_{2,t_{23}+1}^{(i_2)} \cdots a_{2c_2}^{(i_2)}$ is a column word, it remains to check if $a_{1,c_1-(c_2-t_{23})}^{(i_1+1)} < a_{2t_{23}}^{(i_2+1)}$ and $a_{1t_{13}}^{(i_1+1)} < a_{2,c_2-(c_1-t_{13})}^{(i_2+1)}$ (We need to check the latter condition only when $c_2 - (c_1 - t_{13}) > 0$).

If $c_2 - t_{23} = c_1 - t_{13}$, then

$$a_{1,c_1-(c_2-t_{23})}^{(i_1+1)} = a_{1t_{13}}^{(i_1+1)} = a_{2t_{23}}^{(i_2)} < a_{31}^{(i_3)} = a_{2t_{23}}^{(i_2+1)} = a_{2,c_2-(c_1-t_{13})}^{(i_2+1)}.$$

If $c_2 - t_{23} < c_1 - t_{13}$, then $c_2 - (c_1 - t_{13}) < t_{23}$.

$$\text{Hence } a_{1t_{13}}^{(i_1+1)} = a_{2t_{23}}^{(i_2)} < a_{2,c_2-(c_1-t_{13})}^{(i_2)} = a_{2,c_2-(c_1-t_{13})}^{(i_2+1)}.$$

Similarly, $t_{13} < c_1 - (c_2 - t_{23})$ implies

$$a_{1,c_1-(c_2-t_{23})}^{(i_1+1)} = a_{1,c_1-(c_2-t_{23})}^{(i_1)} < a_{2,c_2-(c_2-t_{23})}^{(i_2)} = a_{2t_{23}}^{(i_2)} < a_{31}^{(i_3)} = a_{2t_{23}}^{(i_2+1)}.$$

$$\text{Set } \begin{cases} w_1 := a_{11}^{(i_1)} a_{12}^{(i_1)} \cdots a_{1c_1}^{(i_1)} a_{31}^{(i_3+1)} a_{21}^{(i_2+1)} a_{22}^{(i_2+1)} \cdots a_{2c_2}^{(i_2+1)} a_{32}^{(i_3)} \cdots a_{3c_3}^{(i_3)} \\ w_2 := a_{31}^{(i_3+2)} a_{11}^{(i_1+1)} a_{12}^{(i_1+1)} \cdots a_{1c_1}^{(i_1+1)} a_{21}^{(i_2+1)} a_{22}^{(i_2+1)} \cdots a_{2c_2}^{(i_2+1)} a_{32}^{(i_3)} \cdots a_{3c_3}^{(i_3)} \end{cases},$$

and we thus have $w_0 \smile^* w_1 \smile^* w_2$ and result follows.

Therefore the statement is true for $k = 3$.

Assume the statement is true for all $k = 2, \dots, m, m+1$ for some $m \geq 2$.

When $k = m+2$,

By considering $a_{m,1}^{(i_m)} \cdots a_{m,c_m}^{(i_m)} a_{m+1,1}^{(i_{m+1})} \cdots a_{m+1,c_{m+1}}^{(i_{m+1})} a_{m+2,1}^{(i_{m+2})} \cdots a_{m+2,c_{m+2}}^{(i_{m+2})}$ and apply the result in

$k = 3$, we get

$$\begin{aligned}
& a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \dots a_{2c_2}^{(i_2)} \dots a_{m+1,1}^{(i_{m+1})} \dots a_{m+1,c_{m+1}}^{(i_{m+1})} a_{m+2,1}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})} \\
\rightsquigarrow^* & a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \dots a_{2c_2}^{(i_2)} \dots a_{m+2,1}^{(i_{m+2}+1)} a_{m+1,1}^{(i_{m+1}+1)} \dots a_{m+1,c_{m+1}}^{(i_{m+1}+1)} a_{m+2,2}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})} \\
\rightsquigarrow^* & a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \dots a_{2c_2}^{(i_2)} \dots a_{m+2,1}^{(i_{m+2}+2)} a_{m,1}^{(i_{m+1})} \dots \\
& \dots a_{m,c_m}^{(i_{m+1})} a_{m+1,1}^{(i_{m+1}+1)} \dots a_{m+1,c_{m+1}}^{(i_{m+1}+1)} a_{m+2,2}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})}
\end{aligned}$$

where $a_{m,1}^{(i_{m+1})} \dots a_{m,c_m}^{(i_{m+1})} a_{m+1,1}^{(i_{m+1}+1)} \dots a_{m+1,c_{m+1}}^{(i_{m+1}+1)} a_{m+2,2}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})}$ is a column word, and $a_{m,c_m}^{(i_m)} < a_{m+2,1}^{(i_{m+2}+1)} < a_{m+1,1}^{(i_{m+1}+1)}$.

As $a_{m,c_m}^{(i_m)} < a_{m+2,1}^{(i_{m+2}+1)}$, we can apply the induction assumption on

$$a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} \dots a_{m1}^{(i_m)} \dots a_{mc_2}^{(i_m)} a_{m+2,1}^{(i_{m+2}+1)}$$

and hence we have

$$\begin{aligned}
& a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \dots a_{2c_2}^{(i_2)} \dots a_{m1}^{(i_m)} \dots a_{mc_m}^{(i_m)} a_{m+2,1}^{(i_{m+2}+1)} \\
\rightsquigarrow^* & a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \dots a_{2c_2}^{(i_2)} \dots a_{m-1,1}^{(i_{m-1})} \dots a_{m-1,c_{m-1}}^{(i_{m-1})} a_{m+2,1}^{(i_{m+2}+2)} a_{m1}^{(i_{m+1})} \dots a_{mc_m}^{(i_{m+1})} \\
\rightsquigarrow^* & a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} a_{21}^{(i_2)} \dots a_{2c_2}^{(i_2)} \dots a_{m-2,1}^{(i_{m-2})} \dots a_{m-2,c_{m-2}}^{(i_{m-2})} a_{m+2,1}^{(i_{m+2}+3)} a_{m-1,1}^{(i_{m-1}+1)} \dots \\
& \dots a_{m-1,c_{m-1}}^{(i_{m-1}+1)} a_{m1}^{(i_{m+1})} \dots a_{mc_m}^{(i_{m+1})} \\
& \vdots \\
\rightsquigarrow^* & a_{m+2,1}^{(i_{m+2}+m+1)} a_{11}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{21}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)} \dots a_{m1}^{(i_m+1)} \dots a_{mc_m}^{(i_m+1)}
\end{aligned}$$

$$\text{where } \begin{cases} a_{m-j,c_{m-j}}^{(i_{m-j})} < a_{m+2,1}^{(i_{m+2}+1+j)} < a_{m+1-j,1}^{(i_{m+1-j}+1)} & \forall 1 \leq j < m \\ a_{m+2,1}^{(i_{m+2}+m+1)} < a_{11}^{(i_1+1)} \\ a_{11}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{21}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)} \dots a_{m1}^{(i_m+1)} \dots a_{mc_m}^{(i_m+1)} \text{ is a column word} \end{cases} .$$

Combining the above results, we have

$$\begin{aligned}
& a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} \dots a_{m+1,1}^{(i_{m+1})} \dots a_{m+1,c_{m+1}}^{(i_{m+1})} a_{m+2,1}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})} \\
& \smile^* a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} \dots a_{m1}^{(i_m)} \dots a_{mc_m}^{(i_m)} a_{m+2,1}^{(i_{m+2}+1)} a_{m+1,1}^{(i_{m+1}+1)} \dots a_{m+1,c_{m+1}}^{(i_{m+1}+1)} a_{m+2,2}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})} \\
& \smile^* a_{11}^{(i_1)} \dots a_{1c_1}^{(i_1)} \dots a_{m-1,1}^{(i_{m-1})} \dots a_{m-1,c_{m-1}}^{(i_{m-1})} a_{m+2,1}^{(i_{m+2}+2)} a_{m,1}^{(i_m+1)} \dots a_{m,c_m}^{(i_m+1)} a_{m+1,1}^{(i_{m+1}+1)} \dots \\
& \dots a_{m+1,c_{m+1}}^{(i_{m+1}+1)} a_{m+2,2}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})} \\
& \smile^* a_{11}^{(i_1)} \dots a_{m-2,c_{m-2}}^{(i_{m-2})} a_{m+2,1}^{(i_{m+2}+3)} a_{m-1,1}^{(i_{m-1}+1)} \dots \\
& \dots a_{m-1,c_{m-1}}^{(i_{m-1}+1)} a_{m1}^{(i_m+1)} \dots a_{mc_m}^{(i_m+1)} a_{m+1,1}^{(i_{m+1}+1)} \dots a_{m+1,c_{m+1}}^{(i_{m+1}+1)} a_{m+2,2}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})} \\
& \vdots \\
& \smile^* a_{m+2,1}^{(i_{m+2}+m+1)} a_{11}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} \dots a_{m1}^{(i_m+1)} \dots a_{mc_m}^{(i_m+1)} a_{m+1,1}^{(i_{m+1}+1)} \dots \\
& \dots a_{m+1,c_{m+1}}^{(i_{m+1}+1)} a_{m+2,2}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})}
\end{aligned}$$

Both $a_{11}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} a_{21}^{(i_2+1)} \dots a_{2c_2}^{(i_2+1)} \dots a_{m1}^{(i_m+1)} \dots a_{mc_m}^{(i_m+1)}$ and

$a_{m,1}^{(i_m+1)} \dots a_{m,c_m}^{(i_m+1)} a_{m+1,1}^{(i_{m+1}+1)} \dots a_{m+1,c_{m+1}}^{(i_{m+1}+1)} a_{m+2,2}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})}$ are column words, so

$a_{11}^{(i_1+1)} \dots a_{1c_1}^{(i_1+1)} \dots a_{m1}^{(i_m+1)} \dots a_{mc_m}^{(i_m+1)} a_{m+1,1}^{(i_{m+1}+1)} \dots a_{m+1,c_{m+1}}^{(i_{m+1}+1)} a_{m+2,2}^{(i_{m+2})} \dots a_{m+2,c_{m+2}}^{(i_{m+2})}$

is a column word.

Also,

$$\begin{cases} a_{m-j,c_{m-j}}^{(i_{m-j})} < a_{m+2,1}^{(i_{m+2}+1+j)} < a_{m+1-j,1}^{(i_{m+1-j}+1)} & \forall 1 \leq j < m \\ a_{m,c_m}^{(i_m)} < a_{m+2,1}^{(i_{m+2}+1)} < a_{m+1,1}^{(i_{m+1}+1)} \end{cases}$$

implies $a_{m+1-j,c_{m+1-j}}^{(i_{m+1-j})} < a_{m+2,1}^{(i_{m+2}+j)} < a_{m+2-j,1}^{(i_{m+2-j}+1)} \quad \forall 1 \leq j < m+1$.

Therefore the statement is true for $k = m + 2$.

By Mathematical Induction, the statement is true for all integers $k \geq 2$.

□

Lemma 3.3 shows how to create the first entry of the corresponding row word, leaving the remaining part as a column word. We illustrate the Lemma by an example with $a_{k1}^{(i_k+j)}$ circled for

all $0 \leq j \leq k-1$.

Example 10. Let w_0 be the word in Example 7, hence $k = 5$, $c_1 = 6, c_2 = 5, c_3 = 4, c_4 = c_5 = 1$.

Set $i_j = 0$ for all j . Then we have

$$\left\{ \begin{array}{l} w_0 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | a_{31}^{(0)} \dots a_{34}^{(0)} | a_{41}^{(0)} | \textcircled{a_{51}^{(0)}} = 886531|97643|9764|5| \textcircled{6} \\ w_1 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | a_{31}^{(0)} \dots a_{34}^{(0)} | \textcircled{a_{51}^{(1)}} | a_{41}^{(1)} = 886531|97643|9764| \textcircled{5} | 6 \\ w_2 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | \textcircled{a_{51}^{(2)}} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = 886531|97643| \textcircled{4} | 9765|6 \\ w_3 = a_{11}^{(0)} \dots a_{16}^{(0)} | \textcircled{a_{51}^{(3)}} | a_{21}^{(1)} \dots a_{25}^{(1)} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = 886531| \textcircled{3} | 97644|9765|6 \\ w_4 = \textcircled{a_{51}^{(4)}} | a_{11}^{(1)} \dots a_{16}^{(1)} | a_{21}^{(1)} \dots a_{25}^{(1)} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = \textcircled{1} | 886533|97644|9765|6 \end{array} \right. .$$

Lemma 3.4. Let $w = a_{11} \dots a_{1c_1} \dots a_{k1} \dots a_{kc_k}$ be a column word. Then there exists a sequence $\{b_{ij}\}_{1 \leq j \leq c_i, 1 \leq i \leq k}$ such that

$$w \rightsquigarrow^* b_{k1} b_{k-1,1} \dots b_{11} b_{12} \dots b_{1c_1} b_{22} \dots b_{2c_2} \dots b_{k2} \dots b_{kc_k},$$

where $b_{11} > b_{21} > \dots > b_{k1}$ and $b_{11} \dots b_{1c_1} | b_{22} \dots b_{2c_2} | \dots | b_{k2} \dots b_{kc_k}$ is a column word (with lengths $c_1 > c_2 - 1 \geq \dots \geq c_k - 1$).

Proof. Apply Lemma 3.3 on $a_{11} \dots a_{1c_2} \dots a_{k1} \dots a_{kc_k}$, we have

$$\begin{aligned} & a_{11} \dots a_{1c_1} a_{21} \dots a_{2c_2} \dots a_{k1} \dots a_{kc_k} \\ \rightsquigarrow^* & a_{k1}^{(k-1)} a_{11}^{(1)} \dots a_{1c_1}^{(1)} \dots a_{k-1,1}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k} \end{aligned}$$

where $a_{11}^{(1)} \dots a_{1c_1}^{(1)} \dots a_{k-1,1}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k}$ is a column word.

Apply Lemma 3.3 on $a_{11}^{(1)} \dots a_{1c_1}^{(1)} \dots a_{k-1,1}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)}$, we have

$$\begin{aligned} & a_{11}^{(1)} \dots a_{1c_1}^{(1)} \dots a_{k-1,1}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} \\ \rightsquigarrow^* & a_{k-1,1}^{(k-1)} a_{11}^{(2)} \dots a_{1c_1}^{(2)} \dots a_{k-2,1}^{(2)} \dots a_{k-2,c_{k-2}}^{(2)} a_{k-1,2}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)}, \end{aligned}$$

where $a_{11}^{(2)} \dots a_{1c_1}^{(2)} \dots a_{k-2,1}^{(2)} \dots a_{k-2,c_{k-2}}^{(2)} a_{k-1,2}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)}$ is a column word.

Since by the $k = 2$ case in the proof of Lemma 3.3, we have $a_{k1}^{(1)} < a_{k-1,1}^{(1)}$, then by applying 2. of Lemma 3.2 inductively, we have $a_{k1}^{(k-1)} < a_{k-1,1}^{(k-1)}$.

Combining the above, we have

$$\begin{aligned}
& a_{11} \dots a_{1c_1} a_{21} \dots a_{2c_2} \dots a_{k1} \dots a_{kc_k} \\
& \rightsquigarrow^* a_{k1}^{(k-1)} a_{11}^{(1)} \dots a_{1c_1}^{(1)} \dots a_{k-1,1}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k} \\
& \rightsquigarrow^* a_{k1}^{(k-1)} a_{k-1,1}^{(k-1)} a_{11}^{(2)} \dots a_{1c_1}^{(2)} \dots a_{k-2,1}^{(2)} \dots a_{k-2,c_{k-2}}^{(2)} a_{k-1,2}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k}
\end{aligned}$$

with $a_{k1}^{(k-1)} < a_{k-1,1}^{(k-1)}$ and $a_{11}^{(2)} \dots a_{1c_1}^{(2)} \dots a_{k-2,1}^{(2)} \dots a_{k-2,c_{k-2}}^{(2)} a_{k-1,2}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k}$ is a column word.

Hence by induction, we have

$$\begin{aligned}
& a_{11} \dots a_{1c_1} a_{21} \dots a_{2c_2} \dots a_{k1} \dots a_{kc_k} \\
& \rightsquigarrow^* a_{k1}^{(k-1)} a_{11}^{(1)} \dots a_{1c_1}^{(1)} \dots a_{k-1,1}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k} \\
& \rightsquigarrow^* a_{k1}^{(k-1)} a_{k-1,1}^{(k-1)} a_{11}^{(2)} \dots a_{1c_1}^{(2)} \dots a_{k-2,1}^{(2)} \dots a_{k-2,c_{k-2}}^{(2)} a_{k-1,2}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k} \\
& \rightsquigarrow^* a_{k1}^{(k-1)} a_{k-1,1}^{(k-1)} a_{k-2,1}^{(k-1)} a_{11}^{(3)} \dots a_{1c_1}^{(3)} \dots a_{k-3,1}^{(3)} \dots a_{k-3,c_{k-3}}^{(3)} a_{k-2,1}^{(2)} \dots a_{k-2,c_{k-2}}^{(2)} a_{k-1,2}^{(1)} \dots \\
& \quad \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k} \\
& \quad \quad \quad \vdots \\
& \rightsquigarrow^* a_{k1}^{(k-1)} a_{k-1,1}^{(k-1)} \dots a_{11}^{(k-1)} a_{12}^{(k-1)} \dots a_{1c_1}^{(k-1)} a_{22}^{(k-2)} \dots a_{2c_2}^{(k-2)} \dots \\
& \quad \dots a_{k-1,2}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k}
\end{aligned}$$

where $a_{k1}^{(k-1)} < a_{k-1,1}^{(k-1)} < \dots < a_{11}^{(k-1)}$ and

$a_{11}^{(k-1)} \dots a_{1c_1}^{(k-1)} a_{22}^{(k-2)} \dots a_{2c_2}^{(k-2)} \dots a_{k-1,2}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k}$ is a column word.

Set $b_{i1} := a_{i1}^{(k-1)}$ and $b_{ij} := a_{ij}^{(k-i)}$ for $1 < j \leq c_i$, $1 \leq i \leq k$ and result follows.

□

Example 11. We use the same word in Example 10 to illustrate Lemma 3.4 by repeated use of Lemma 3.3. We first have

$$w_0 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | a_{31}^{(0)} \dots a_{34}^{(0)} | a_{41}^{(0)} | a_{51}^{(0)} = 886531|97643|9764|5|6$$

and by Example 10, we have

$$a_{51}^{(4)} | a_{11}^{(1)} \dots a_{16}^{(1)} | a_{21}^{(1)} \dots a_{25}^{(1)} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = 1 | 886533 | 97644 | 9765 | 6$$

and we apply Example 10 again on the latter part (marked by the rectangle), we have

$$\left\{ \begin{array}{l} a_{51}^{(4)} | a_{11}^{(1)} \dots a_{16}^{(1)} | a_{21}^{(1)} \dots a_{25}^{(1)} | a_{41}^{(2)} | a_{31}^{(2)} \dots a_{34}^{(2)} = 1 | 886533 | 97644 | 5 | 9766 \\ a_{51}^{(4)} | a_{11}^{(1)} \dots a_{16}^{(1)} | a_{41}^{(3)} | a_{21}^{(2)} \dots a_{25}^{(2)} | a_{31}^{(2)} \dots a_{34}^{(2)} = 1 | 886533 | 4 | 97654 | 9766 \\ a_{51}^{(4)} | a_{41}^{(4)} | a_{11}^{(2)} \dots a_{16}^{(2)} | a_{21}^{(2)} \dots a_{25}^{(2)} | a_{31}^{(2)} \dots a_{34}^{(2)} = 1 | 3 | 886543 | 97654 | 9766 \end{array} \right.$$

and again we apply Example 10 again on the latter part (marked by the rectangle), we have

$$\left\{ \begin{array}{l} a_{51}^{(4)} a_{41}^{(4)} | a_{11}^{(2)} \dots a_{16}^{(2)} | a_{21}^{(2)} \dots a_{25}^{(2)} | a_{31}^{(2)} | a_{32}^{(2)} a_{33}^{(2)} a_{34}^{(2)} = 13 | 886543 | 97654 | 9 | 766 \\ a_{51}^{(4)} a_{41}^{(4)} | a_{11}^{(2)} \dots a_{16}^{(2)} | a_{31}^{(3)} | a_{21}^{(3)} \dots a_{25}^{(3)} | a_{32}^{(2)} a_{33}^{(2)} a_{34}^{(2)} = 13 | 886543 | 7 | 99654 | 766 \\ a_{51}^{(4)} a_{41}^{(4)} | a_{31}^{(4)} | a_{11}^{(3)} \dots a_{16}^{(3)} | a_{21}^{(3)} \dots a_{25}^{(3)} | a_{32}^{(2)} a_{33}^{(2)} a_{34}^{(2)} = 13 | 6 | 887543 | 99654 | 766 \end{array} \right.$$

and by repeating the same process again, we get

$$\left\{ \begin{array}{l} a_{51}^{(4)} a_{41}^{(4)} a_{31}^{(4)} | a_{11}^{(3)} \dots a_{16}^{(3)} | a_{21}^{(3)} | a_{22}^{(3)} \dots a_{25}^{(3)} | a_{32}^{(2)} a_{33}^{(2)} a_{34}^{(2)} = 136 | 887543 | 9 | 9654 | 766 \\ a_{51}^{(4)} a_{41}^{(4)} a_{31}^{(4)} | a_{21}^{(4)} | a_{11}^{(4)} \dots a_{16}^{(4)} | a_{22}^{(3)} \dots a_{25}^{(3)} | a_{32}^{(2)} a_{33}^{(2)} a_{34}^{(2)} = 136 | 8 | 987543 | 9654 | 766 \end{array} \right.$$

and finally we get

$$a_{51}^{(4)} a_{41}^{(4)} a_{31}^{(4)} a_{21}^{(4)} | a_{11}^{(4)} a_{12}^{(4)} \dots a_{16}^{(4)} | a_{22}^{(3)} \dots a_{25}^{(3)} | a_{32}^{(2)} a_{33}^{(2)} a_{34}^{(2)} = 1368 | 987543 | 9654 | 766.$$

So we have:

$$b_{51} b_{41} b_{31} b_{21} | b_{11} b_{12} \dots b_{16} | b_{22} \dots b_{25} | b_{31} \dots b_{34} = 1368 | 987543 | 9654 | 766, \text{ where}$$

$$b_{11} b_{12} \dots b_{16} | b_{22} \dots b_{25} | b_{32} b_{33} b_{34} = 987543 | 9654 | 766 \text{ is a column word with lengths } 6, 4, 3.$$

Using the notation in Lemma 3.4, since

$$w \smile^* b_{k_1} b_{k-1,1} \dots b_{11} b_{12} \dots b_{1c_1} b_{22} \dots b_{2c_2} \dots b_{k2} \dots b_{kc_k},$$

where $b_{11} > b_{21} > \dots > b_{k1}$, the SSAF with basement being ϵ_n representing the word w is the same as that representing $b_{k_1} b_{k-1,1} \dots b_{11} b_{12} \dots b_{1c_1} b_{22} \dots b_{2c_2} \dots b_{k2} \dots b_{kc_k}$. Denote $F(w)$ as the SSAF created.

Since $b_{k_1} b_{k-1,1} \dots b_{11}$ is strictly increasing, by Lemma 15 in [4], they create new cells in ascending reading order (i.e. one after another) and hence is exactly the first entire row of $F(w)$ as the entries

are fixed when inserting $b_{k1}b_{k-1,1} \dots b_{11}$ into an empty atom, and there are k columns (as w has k subsequences) and so the first row has length k and hence the row reading word has exactly $b_{k1}, b_{k-1,1}, \dots, b_{11}$ as the first subsequence. That means we can apply Lemma 3.4 to find the first subsequence of the row reading word of $F(w)$.

Since $b_{12} \dots b_{1c_1} b_{22} \dots b_{2c_2} \dots b_{k2} \dots b_{kc_k}$ is a column word, we can apply Lemma 3.4 again and get the second subsequence of the row reading word of $F(w)$, and we can apply Lemma 3.4 repeatedly on the remaining b_{ij} 's until we get all the subsequences of the row reading word of $F(w)$. As a result, we can convert w into the row reading word of $F(w)$ by applying Lemma 3.4 repeatedly as described.

We illustrate this by using the example in Example 11:

Example 12. *From Example 7 and Example 11, we notice that $b_{51}b_{41}b_{31}b_{21}b_{11} = 13689$ which is exactly the first subsequence of the row word of the SSAF representing the column word $w = 886531|97643|9764|5|6$. Also one can check that $87543|9654|766$ is a column word (by Definition 3.3).*

We can apply Lemma 3.4 on $87543|9654|766$ as we did in Example 11 on $886531|97643|9764|5|6$ and get $589|7643|754|66$ which gives the second subsequence of the row word: 589 , and by repeating the same process, we get $467|753|64|6$, $357|64|6$, $46|6$ and lastly 6 .

As a result, we get 467 , 357 , 46 and 6 as the third to the last subsequence of the row word (see Example 7).

3.2 Convert a Column Recording Tableau to a Row Recording Tableau

This section gives an interpretation of the twisted Knuth equivalence using recording tableaux. We use the insertion in [11] and the generalized Littlewood-Richardson rule in [3]. We also use the

notation $(U \leftarrow W)$ for an SSAF U and a biword $W = \begin{pmatrix} x_1 & x_2 & \dots & x_n \\ y_1 & y_2 & \dots & y_n \end{pmatrix}$ for some positive integer

n to denote the pair (U', L) where U' is the SSAF obtained by $(U \leftarrow y_1 y_2 \dots y_n)$ while L is the recording tableau, i.e. by putting x_i into the cell created when y_i is being inserted. In particular, if $y_1 y_2 \dots y_n$ is a column (resp. row) word, then we call L as a column (resp. row) recording tableau. By abuse of notation, we sometimes refer $(U \leftarrow W)$ to either V or L (depending on the context).

(Note that if we change the basement ϵ_n into the large basement in [3], a column recording tableau is the same as an LRS defined in Section 4 of [3].)

Lemma 3.5. *Let U be an SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & 1 \\ a & b & c \end{pmatrix}$, $a \geq b, c > b$ (i.e. $ab|c$ is a column word.). Let $L = U \leftarrow W$ be the recording tableau. Let V be the SSAF representing the word abc and $a'b'c'$ be the row reading word of V (so $a' < b'$ and $c' \leq b'$ and $abc \rightsquigarrow^* a'b'c'$). Consider the biword $\widetilde{W} = \begin{pmatrix} 1 & 1 & 2 \\ a' & b' & c' \end{pmatrix}$ and let $\widetilde{L} = U \leftarrow \widetilde{W}$ be the recording tableau. Then L determines \widetilde{L} .*

Proof. There are two cases to consider: $a \geq c$ or $a < c$.

Case(I): $a \geq c > b$

Hence we have $a'b'c' = bac$.

We first consider L . Since $a > b$, when we insert ab into U , i.e. $\left(U \leftarrow \begin{pmatrix} 2 & 2 \\ a & b \end{pmatrix} \right)$, the cell created by b is strictly above that created by inserting a (by Lemma 15 in [4]):

- (i) The cell appears when inserting b is immediately above that of inserting a :

$$\begin{array}{|c|} \hline 2 \\ \hline 2 \\ \hline \end{array}$$

- (ii) The two cells are the top cells of two distinct columns (by Theorem 16 in [4])

and the cells appear in ascending reading order, (i.e. one after another):

Since $c > b$, by Lemma 15 in [4], the cell created when inserting c after inserting ab into U is after the first cell created by inserting b in reading order.

For (i), we have $L =$

$$(a) \begin{array}{|c|} \hline 2 \\ \hline 2 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 1 \\ \hline \end{array} \quad \text{or} \quad (b) \quad \begin{array}{|c|} \hline 2 \\ \hline 2 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 1 \\ \hline \end{array} \quad \text{or} \quad (c) \quad \begin{array}{|c|} \hline 2 \\ \hline 2 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 1 \\ \hline \end{array} \quad \text{or} \quad (d) \quad \begin{array}{|c|} \hline 2 \\ \hline 2 \\ \hline \end{array} \quad \begin{array}{|c|} \hline 1 \\ \hline \end{array}$$

(a) is impossible as the cell created by inserting c , i.e. $\begin{array}{|c|} \hline 1 \\ \hline \end{array}$, is not a removable cell (defined in [4]).

For (b), $\tilde{L} = \begin{array}{|c|} \hline 2 \\ \hline 1 \\ \hline \end{array}$ as the cell created by c' must be the top cell of some column and is strictly above that created by b' but there is only one such cell.

(c) and (d) can be considered as the same case by viewing the cell created by inserting c is after that created by inserting a (the second $\begin{array}{|c|} \hline 2 \\ \hline \end{array}$ in reading order). By the same argument as (b), we have $\tilde{L} = \begin{array}{|c|} \hline 2 \\ \hline 1 \\ \hline \end{array}$

For (ii), since the reading word of L is contre-lattice by [3] which means it should be 221 or 212, the cell created by inserting c is either the middle in reading order or the last one in reading order.

Suppose the cell created by inserting c is the middle cell among the three in reading order, i.e. the reading word of L is 212.


Since $c' = c \leq a = b' > a' = b$, so when performing $(U \leftarrow a'b'c' = bac)$ to get \tilde{L} , the cell created by inserting a is after both of that inserted by b and by c , by Lemma 15 of [4], the last cell must be created when a is inserted and hence marked as 1.

If the middle cell (in reading order) is immediately above the last cell, then it cannot be created by $(U \leftarrow b)$ as the last cell is not created if b is inserted before a , and so it must be marked as 2, i.e. when $c' = c$ is inserted.

$$\square \quad \begin{array}{|c|} \hline 2 \\ \hline 1 \\ \hline \end{array} \quad \text{or} \quad \begin{array}{|c|} \hline 2 \\ \hline 1 \\ \hline \end{array} \quad \square$$

Suppose the middle cell (in reading order) is not immediately above the last cell, then all three cells are the top cell of three distinct columns. Suppose the middle cell is

marked as 1 in \tilde{L} , meaning that when we insert b before inserting a , the cell created is after the cell created by b when we first insert a into U .



That means the bumping sequence for $(U \leftarrow a)$ involves some cell in the bumping sequence in $(U \leftarrow b)$ and this implies the cell created by $(U \leftarrow a)$ is the same as $(U \leftarrow b)$ (because the bumping sequence has the same ending subsequence starting from the common cell and hence creating the same final cell), and this leads to a contradiction.

Therefore we have the reading order (from first to last in reading order) of $\tilde{L} = 121$.

Now consider the case when L has the reading word 221.

By the same argument, we know that the last cell in \tilde{L} is created when $b' = a$ is inserted. If $a' = b$ creates the first cell among the three in reading order as b does in L after a is inserted, then the insertion of $(U \leftarrow a)$ has no common cell in the bumping sequence of $(U \leftarrow b)$. That means when we insert a after inserting b , the bumping sequence is the same as inserting a before inserting b (this is true because the bumping sequence of b is a decreasing sequence, so inserting b into U would not affect the first (in reading order) cell in U containing an entry larger than a as $a > b$). That means $(U \leftarrow ab)$ is the same as $(U \leftarrow ba)$. This leads to a contradiction as we assumed a create different cells in the two cases (the middle and the last cell respectively).

So we know the middle cell is marked as 1 in \tilde{L} . So the reading word of \tilde{L} is 211.

Case(II): $c > a \geq b$

Then $a'b'c' = acb$. Hence the first cells created in L and \tilde{L} are always the same. Note that $c > a$ meaning the cell created by inserting $a = a'$ is not the last (in reading order) cell, and $a \leq b$ meaning that the cell created by inserting a is not the first (in reading order) cell either. As a result, we know that the middle cell must be created

when a is inserted. Since the first (in reading order) is created in L when inserting b while the last (in reading order) cell is created in \tilde{L} when inserting $b' = c$ is inserted, we know the first cell of L is marked as 2 while the last cell in \tilde{L} must be marked as 1 and that means the reading word of L is 221 and that of \tilde{L} is 211.

This shows that if the reading word of L is 221 or when L is of the form as **Case(I)**(i) then the reading word of \tilde{L} is 211, otherwise the reading word of \tilde{L} is 121.

Hence L determines \tilde{L} .

□

Lemma 3.6. *Let U be an SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & \cdots & 2 & 1 \\ a_1 & a_2 & \cdots & a_k & b \end{pmatrix}$, $a_1 \geq a_2 \geq \cdots \geq a_k, b > a_k$ (i.e. $a_1 a_2 \dots a_k | b$ is a column word). Let $L = U \leftarrow W$ be the recording tableau. If $b > a_1$, then L has reading word $22 \dots 21$, i.e. b creates the last cell in reading order and the insertion of $a_2 \dots a_k$ is independent of the insertion of b .*

Proof. Since $a_1 a_2 \dots a_k b \rightsquigarrow^* a_1 b a_2 \dots a_k$, we have

$$(U \leftarrow a_1 a_2 \dots a_k b) = (U \leftarrow a_1 b a_2 \dots a_k).$$

When $k = 2$, the result follows by **Case(II)** in the proof of Lemma 3.5.

For $k > 2$, since $a_1 a_2 \dots a_k b \rightsquigarrow^* a_1 b a_2 \dots a_k$, we have $(U \leftarrow a_1 a_2 \dots a_k b) = (U \leftarrow a_1 b a_2 \dots a_k) = ((U \leftarrow a_1 b a_2) \leftarrow a_3 \dots a_k)$. By the case when $k = 2$, we know b creates a cell after both a_1 and a_2 (and $a_3 \dots a_k$ create cells with decreasing reading order each of which has an order smaller than that of the cell created by a_2) and result follows. □

Example 13. Let $W = \begin{pmatrix} 2 & 2 & 2 & 2 & 2 & 2 & 1 \\ 8 & 8 & 6 & 5 & 3 & 1 & 9 \end{pmatrix}$ where $a_1 = 8 < 9 = b$. Let $n = 9$ and U be an SSAF with basement $\epsilon_9 = 123456789$ and shape $\alpha = (1, 1, 0, 0, 3, 0, 2, 0, 6)$:

$$U = \begin{array}{cccccccccc} & & & & & & & & & 1 \\ & & & & & & & & & 2 \\ & & & & & & & & & 2 \\ & & & & & & & & & 6 \\ & & & & & 1 & & 4 & & 8 \\ & & & & 5 & 5 & & 7 & & 9 \\ 1 & 2 & & & & & & & & \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & \end{array}$$

Therefore, $(U \leftarrow W)$ means:

$$\begin{array}{cccccccccc} & & & & & & & & & 1 \\ & & & & & & & & & 2 \\ & & & & & & & & & 2 \\ & & & & & & & & & 6 \\ & & & & & 1 & & 4 & & 8 \\ & & & & 5 & 5 & & 7 & & 9 \\ 1 & 2 & & & & & & & & \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & \end{array} \leftarrow \begin{pmatrix} 2 & 2 & 2 & 2 & 2 & 2 & 1 \\ 8 & 8 & 6 & 5 & 3 & 1 & 9 \end{pmatrix}$$

which results in

$$\begin{array}{cccccccccc} & & & & & & & & & 1 \\ & & & & & & & & & 3 \\ & & & & & & & & & 5 \\ & & & & & & & & & 6 \\ & & & & & 1 & & 1 & & 8 \\ & & & & 2 & 2 & & 6 & & 9 \\ & & & & 5 & 5 & & 7 & & 9 \\ 1 & 2 & & & & & & & & \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & \end{array} \quad \begin{array}{cccccccccc} & & & & & & & & & 2 \\ & & & & & & & & & 2 \\ & & & & & & & & & 2 \\ & & & & & & & & & 2 \\ & & & & & & & & & 1 \\ & & & & & & & & & 9 \\ & & & & & & & & & \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & \end{array}$$

When 9 is inserted, it creates the last cell (we marked the cell red) in reading order among all cells created, and one can see this by reading the recording tableau where the cell is marked as 1, which is the last cell in reading order among all cells.

Also, note that the bumping route of 9 starts from the second row of U while those of $a_2, \dots, a_6 = 8, 6, 5, 3, 1$ starts from the third row. Hence the insertion of 9 does not affect the bumping routes of inserting a_2, \dots, a_6 .

Lemma 3.7. Let U be an SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & \cdots & 2 & 1 \\ a_1 & a_2 & \cdots & a_k & b \end{pmatrix}$, $a_1 \geq a_2 \geq \cdots \geq a_k, b > a_k$ (i.e. $a_1 a_2 \dots a_k | b$ is a column word). Let $L = (U \leftarrow W)$ be the recording tableau. If $i = \min_{1 \leq j \leq k} \{j : b > a_j\}$, then the cell created by inserting a_m for $m > i$ is not affected by the insertion of b .

Proof. Since $(U \leftarrow a_1 a_2 \dots a_k b) = ((U \leftarrow a_1 a_2 \dots a_{i-1}) \leftarrow a_i \dots a_k b)$. By applying Lemma 3.6 with the U in the lemma being $(U \leftarrow a_1 \dots a_{i-1})$ and result follows. □

Lemma 3.8. Let U be an SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & \cdots & 2 & 1 \\ a_1 & a_2 & \cdots & a_k & b \end{pmatrix}$, $a_1 \geq a_2 \geq \cdots \geq a_k, b > a_k$ (i.e. $a_1 a_2 \dots a_k | b$ is a column word). Let $L = (U \leftarrow W)$ be the recording tableau. Suppose $b \leq a_{k-1}$. Let $\tilde{L} = (U \leftarrow \tilde{W})$ where $\tilde{W} = \begin{pmatrix} 1 & 1 & 2 & \cdots & 2 & 2 \\ a_k & a_1 & a_2 & \cdots & a_{k-1} & b \end{pmatrix}$, then L determines \tilde{L} .

Proof. When $k = 2$, this is proved by **Case(I)** in Lemma 3.5.

For $k > 2$, since $a_1 \dots a_k b \smile^* a_k a_1 \dots a_{k-1} b$, L and \tilde{L} has the same shape.

Also, $a_1 \dots a_k b \smile^* a_1 \dots a_{k-1} a_k b$, we have


$$(U \leftarrow a_1 \dots a_{k-1} a_k b) = ((U \leftarrow a_1 \dots a_{k-2}) \leftarrow a_{k-1} a_k b).$$

By applying the case when $k = 2$ on $(U \leftarrow a_1 \dots a_{k-2})$ and $a_{k-1} a_k b$ being the length-3-word inserted, we know the order of the cell being created which represents $((U \leftarrow a_1 \dots a_{k-2}) \leftarrow a_{k-1} a_k b) = ((U \leftarrow a_1 \dots a_{k-2}) \leftarrow a_k a_{k-1} b)$ and so we know which one among the three is the last cell being created (by Lemma 3.5, the last cell being created may either be the first or second cell in reading order among the three cells created). By marking that cell as 2 in \tilde{L} and then consider the first two cells being created we know how the cells are being created by the insertion on $((U \leftarrow a_1 \dots a_{k-2}) \leftarrow a_k a_{k-1} b) = (U \leftarrow a_1 \dots a_{k-2} a_k a_{k-1} b) = (U \leftarrow a_k a_1 \dots a_{k-2} a_{k-1} b)$ by induction (as $a_{k-2} \leq a_{k-1} < a_k$ for $a_k < b \leq a_{k-1}$) and together with the last cell marked by 2 as mentioned, we know how to label the entries of the recording tableau \tilde{L} for $(U \leftarrow a_k a_1 \dots a_{k-2} a_{k-1} b)$. □


Example 14. Let U as in Example 13 and let $W = \begin{pmatrix} 2 & 2 & 2 & 2 & 1 \\ 9 & 7 & 6 & 4 & 5 \end{pmatrix}$.

$$\text{Then } \tilde{W} = \begin{pmatrix} 2 & 2 & 2 & 2 & 1 \\ 4 & 9 & 7 & 6 & 5 \end{pmatrix}.$$

$(U \leftarrow W)$ means:




which results in




We illustrate the proof of Lemma 3.8 to get \tilde{L} from L .

Consider the last three cells created which are marked green in L :



We get the first cell (in reading order) in \tilde{L} (the yellow cell) marked as 2 by applying Case (ii) in the proof of Lemma 3.5 as we know the green cells would have reading 211 in \tilde{L} which implies the first cell in reading order among the three green cells in L must be marked with 2 in \tilde{L} while the two cells appear in the order as circled (i.e. the one marked as $\textcircled{1}$ appear before the one marked with $\textcircled{2}$) when we insert $a_4a_3b = 465$ to get \tilde{L} .)


Hence we can treat the cell marked with $\textcircled{2}$ as the last cell created among the remaining 4 cells in \tilde{L} to be filled, and so we mark that cell as 1 (treating it as the new b being inserted as $4 < 6 < 7$). Now we have the three new lastly-created cells marked green:



Again, we get the second (in reading order) cell among the three green cells (the new yellow cell) marked as 2 by applying Case (ii) in the proof of Lemma 3.5 as we know the green cells would

have reading 121 in \tilde{L} which implies the second cell in reading order among the three green cells in L must be marked 2 in \tilde{L} while the two cells appear in the order as circled (i.e. the one marked with $\textcircled{1}$ appears before the one marked with $\textcircled{2}$ when we insert $a_4a_2a_3 = 476$ to get \tilde{L} .)

Hence we can, again, treat the cell marked with $\textcircled{2}$ as the last cell created among the remaining three cells in \tilde{L} to be filled, and so mark it as 1 and the other two cells as 2 by a similar argument as the previous step. Now we have the three new lastly-created cells marked green:



We get the remaining entries of \tilde{L} by applying Lemma 3.5 again as we know the reading word among these three cells would change from 212 to 121.

One can verify that \tilde{L} is indeed the recording tableau of $(U \leftarrow \tilde{W})$.

Lemma 3.9. Let U be an SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Consider the biword $W = \begin{pmatrix} 2 & 2 & \cdots & 2 & 1 \\ a_1 & a_2 & \cdots & a_k & b \end{pmatrix}$, $a_1 \geq a_2 \geq \cdots \geq a_k, b > a_k$ (i.e. $a_1a_2 \dots a_k|b$ is a column word). Let $L = (U \leftarrow W)$ be the recording tableau. Let $\tilde{L} = (U \leftarrow \tilde{W})$ where $\tilde{W} = \begin{pmatrix} 1 & 1 & 2 & \cdots & 2 \\ b' & a'_1 & a'_2 & \cdots & a'_k \end{pmatrix}$, such that $b' = a_i$, where $i = \min_{1 \leq j \leq k} \{j : b > a_j\}$ and $b' < a'_1$, $a'_1 \geq a'_2 \geq \cdots \geq a'_k$ (one can verify that $b'a'_1 \dots a'_k$ is indeed the row reading word of the SSAF representing the word $a_1a_2 \dots a_kb$ by 1. in Lemma 3.2), then L determines \tilde{L} .

Proof. If $i = k$, then we are done by Lemma 3.8.

If $i = 1$, then by 1. in Lemma 3.2, we have $b'a'_1 \dots a'_k = a_1ba_2 \dots a_k$ and hence by Lemma 3.6, b would create the last cell in reading order in \tilde{L} when being inserted and hence we know that which two cells are created by a_1 and b in \tilde{L} and hence we know how to label the entries by marking those two cells as 1 and the rest as 2.

Suppose $i < k$ then by 1. in Lemma 3.2, we have

$$b' a'_1 \dots a'_k = a_i a_1 \dots a_{i-1} b a_{i+1} \dots a_k.$$

Now by Lemma 3.7, we know all $a_j, j > i$ creates create the same cells in L and \tilde{L} and since $a_1 \dots a_{i-1} a_i b \rightsquigarrow^* a_i a_1 \dots a_{i-1} b$, we just need to first apply Lemma 3.8 on $(U \leftarrow a_1 \dots a_i b)$ as the insertion recorded by L in the lemma in order to find the first $i+1$ cells created in \tilde{L} by the insertion $(U \leftarrow a_i a_1 \dots a_{i-1} b)$ and then label the rest of cells created by $((U \leftarrow a_i a_1 \dots a_{i-1} b) \leftarrow a_{i+1} \dots a_k)$ as 2 and get the entries of \tilde{L} and result follows. \square

Lemma 3.9 gives a recording tableau interpretation of 1. in Lemma 3.2 using L and \tilde{L} .

Lemma 3.10. *Let U be an SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Let $w_0 = a_{11} \dots a_{1c_1} | a_{21} \dots a_{2c_2} | \dots | a_{k1} \dots a_{kc_k}$ with $k \leq n$ be a column word and using the notation in Lemma 3.3 while we assume $(i_j) = 0$ for $1 \leq j \leq c_r, 1 \leq r \leq k$, the recording tableau L of $(U \leftarrow W_0)$ determines the recording tableau \tilde{L}_m to $(U \leftarrow W_m)$ for $0 \leq m \leq k-1$, where $L := \tilde{L}_0$ and W_m is a biword with the lower word being w_m and the upper word has entry $k+1-t$ if the lower word entry just below it is $a_{ij}^{(s)}$ for $s = i_t, i_t + 1$, and $1_{(1)}$ if $m > 0$ and the entry is $a_{k1}^{i_k+m}$.*

Proof. Suppose $k = 2$ then

$$(U \leftarrow w_0) = ((U \leftarrow a_{11} \dots a_{1c_1} a_{21}) \leftarrow a_{22} \dots a_{2c_2}) = (((U \leftarrow a_{21}^{(1)}) \leftarrow a_{11}^{(1)} \dots a_{1c_1}^{(1)}) \leftarrow a_{22} \dots a_{2c_2})$$

and by Lemma 3.9 we know how the cells are created in $((U \leftarrow a_{21}^{(1)}) \leftarrow a_{11}^{(1)} \dots a_{1c_1}^{(1)})$. So giving the recording tableau L of $(U \leftarrow a_{11} \dots a_{1c_1} a_{21})$, we know how the cells are created by $(U \leftarrow a_{21}^{(1)} a_{11}^{(1)} \dots a_{1c_1}^{(1)})$. Now inserting the rest of the sequence $a_{22} \dots a_{2c_2}$ and which creates in \tilde{L}_1 the same last $(c_2 - 1)$ cells as in L , we know how to label the recording tableau \tilde{L}_1 .

Suppose $k > 2$, then

$$\begin{aligned} (U \leftarrow w_0) &= (U \leftarrow a_{11} \dots a_{1c_1} a_{21} \dots a_{2c_2} \dots a_{k1} \dots a_{kc_k}) \\ &= ((U \leftarrow a_{11} \dots a_{1c_1} \dots a_{k-2,1} \dots a_{k-2,c_{k-2}}) \leftarrow a_{k-1,1} \dots a_{k-1,c_{k-1}} a_{k1} \dots a_{kc_k}) \end{aligned}$$

and by the argument

with

$$(U \leftarrow a_{11} \dots a_{1c_1} \dots a_{k-2,1} \dots a_{k-2,c_{k-2}})$$

being the U to be considered for $k = 2$, we know how the cells are created in \tilde{L}_1 for

$$((U \leftarrow a_{11} \dots a_{1c_1} \dots a_{k-2,1} \dots a_{k-2,c_{k-2}}) \leftarrow a_{k1}^{(1)} a_{k-1,1}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k})$$

and by the same argument, we know how the cells are created in \tilde{L}_2 for

$$((U \leftarrow a_{11} \dots a_{1c_1} \dots a_{k-3,1} \dots a_{k-3,c_{k-3}}) \leftarrow a_{k1}^{(2)} a_{k-2,1}^{(1)} \dots a_{k-2,c_{k-2}}^{(1)} a_{k-1,1}^{(1)} \dots a_{k-1,c_{k-1}}^{(1)} a_{k2} \dots a_{kc_k})$$

and result follows by induction. □

Example 15. We use the words in Example 10 and the same U as in Example 13 to illustrate Lemma 3.10.

Recall that in Example 10, we have

$$\left\{ \begin{array}{l} w_0 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | a_{31}^{(0)} \dots a_{34}^{(0)} | a_{41}^{(0)} | \textcircled{a_{51}^{(0)}} = 886531|97643|9764|5| \textcircled{6} \\ w_1 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | a_{31}^{(0)} \dots a_{34}^{(0)} | \textcircled{a_{51}^{(1)}} | a_{41}^{(1)} = 886531|97643|9764| \textcircled{5} |6 \\ w_2 = a_{11}^{(0)} \dots a_{16}^{(0)} | a_{21}^{(0)} \dots a_{25}^{(0)} | \textcircled{a_{51}^{(2)}} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = 886531|97643| \textcircled{4} |9765|6 \\ w_3 = a_{11}^{(0)} \dots a_{16}^{(0)} | \textcircled{a_{51}^{(3)}} | a_{21}^{(1)} \dots a_{25}^{(1)} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = 886531| \textcircled{3} |97644|9765|6 \\ w_4 = \textcircled{a_{51}^{(4)}} | a_{11}^{(1)} \dots a_{16}^{(1)} | a_{21}^{(1)} \dots a_{25}^{(1)} | a_{31}^{(1)} \dots a_{34}^{(1)} | a_{41}^{(1)} = \textcircled{1} |886533|97644|9765|6 \end{array} \right. .$$

Therefore, we now have


$$\left\{ \begin{array}{l}
W_0 = \left(\begin{array}{c|c|c|c|c}
5 & 5 & 5 & 5 & 5 & 5 & 4 & 4 & 4 & 4 & 4 & 3 & 3 & 3 & 3 & 2 & 1 \\
8 & 8 & 6 & 5 & 3 & 1 & 9 & 7 & 6 & 4 & 3 & 9 & 7 & 6 & 4 & 5 & \textcircled{6}
\end{array} \right) \\
W_1 = \left(\begin{array}{c|c|c|c|c}
5 & 5 & 5 & 5 & 5 & 5 & 4 & 4 & 4 & 4 & 4 & 3 & 3 & 3 & 3 & 1_{(1)} & 2 \\
8 & 8 & 6 & 5 & 3 & 1 & 9 & 7 & 6 & 4 & 3 & 9 & 7 & 6 & 4 & \textcircled{5} & 6
\end{array} \right) \\
W_2 = \left(\begin{array}{c|c|c|c|c}
5 & 5 & 5 & 5 & 5 & 5 & 4 & 4 & 4 & 4 & 4 & 1_{(1)} & 3 & 3 & 3 & 3 & 2 \\
8 & 8 & 6 & 5 & 3 & 1 & 9 & 7 & 6 & 4 & 3 & \textcircled{4} & 9 & 7 & 6 & 5 & 6
\end{array} \right) \\
W_3 = \left(\begin{array}{c|c|c|c|c}
5 & 5 & 5 & 5 & 5 & 5 & 1_{(1)} & 4 & 4 & 4 & 4 & 4 & 3 & 3 & 3 & 3 & 2 \\
8 & 8 & 6 & 5 & 3 & 1 & \textcircled{3} & 9 & 7 & 6 & 4 & 4 & 9 & 7 & 6 & 5 & 6
\end{array} \right) \\
W_4 = \left(\begin{array}{c|c|c|c|c}
1_{(1)} & 5 & 5 & 5 & 5 & 5 & 4 & 4 & 4 & 4 & 4 & 3 & 3 & 3 & 3 & 2 \\
\textcircled{1} & 8 & 8 & 6 & 5 & 3 & 3 & 9 & 7 & 6 & 4 & 4 & 9 & 7 & 6 & 5 & 6
\end{array} \right)
\end{array} \right.$$

We now show how to get \tilde{L}_1 to \tilde{L}_4 from $L = \tilde{L}_0$.


For $U \leftarrow W_0$:

$$\left\{ \begin{array}{l}
\begin{array}{cccccccccc}
& & & & & & & & & & \boxed{1} \\
& & & & & & & & & & \boxed{2} \\
& & & & & & & & & & \boxed{2} \\
& & & & & & & & & & \boxed{6} \\
& & & & & & \boxed{1} & & & & \boxed{8} \\
& & & & & \boxed{5} & \boxed{4} & & & & \boxed{9} \\
& & & & & \boxed{5} & \boxed{7} & & & & \\
\boxed{1} & \boxed{2} & & & & & & & & & \\
1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & &
\end{array} \\
\leftarrow \left(\begin{array}{c|c|c|c|c}
5 & 5 & 5 & 5 & 5 & 5 & 4 & 4 & 4 & 4 & 4 & 3 & 3 & 3 & 3 & 2 & 1 \\
8 & 8 & 6 & 5 & 3 & 1 & 9 & 7 & 6 & 4 & 3 & 9 & 7 & 6 & 4 & 5 & 6
\end{array} \right)
\end{array} \right.$$


which results in




To get \tilde{L}_0 , we just consider the last (in reading order) cell containing 1 and all cells containing 2 and 1 and apply Lemma 3.10. Note that in this case since there are only two cells involved, we can simply move 1 to the cell with a smaller reading order cell and mark as $1_{(1)}$:




From \tilde{L}_1 we get \tilde{L}_2 by considering the cell containing $1_{(1)}$ and all the cells containing 3 and apply Lemma 3.10 to find the new position of $1_{(1)}$:



Then consider the cells with 4 and also the cell with $1_{(1)}$ in \tilde{L}_2 , we get \tilde{L}_3 by applying Lemma 3.10 to find the new position of $1_{(1)}$:



By the same argument and consider the cells with 5 and also the cell with $1_{(1)}$, we get \tilde{L}_4 :



We describe a more direct way to get $L = \tilde{L}_0$ from \tilde{L}_4 . Note that the reading word of L is a contre-lattice word, we can always find a 2 before (in reading order) 1. By the last part in the proof of Lemma 3.5, we know that, except for the case when L is of the form of **Case(I)(i)**, the first cell created in \tilde{L} (which is the first cell in reading order containing a 1) is the cell with largest reading order containing 2 in L before the cell containing 1 in L , and so we put $1_{(1)}$ into that cell in \tilde{L} .

In short, the red cell with $1_{(1)}$ is always interchanged with the last green cell before it in reading order, except for the case when that green cell is immediately above the next green cell (which must be after the red cell if it exists) in which the red cell stays the same and compare to the next set of green cells (if available).

Lemma 3.11. Let U be an SSAF with basement ϵ_n and shape α for some positive integer n and $l(\alpha) \leq n$. Let $w = a_{11} \dots a_{1c_1} | a_{21} \dots a_{2c_2} | \dots | a_{k1} \dots a_{kc_k}$ with $k \leq n$ be a column word and using the notation in Lemma 3.4, the recording tableau L of $(U \leftarrow W)$ determines the recording tableau \tilde{L}

of $(U \leftarrow \widetilde{W})$, where

$$W = \begin{pmatrix} k & k & \dots & k & \dots & 1 & 1 & \dots & 1 \\ a_{11} & a_{12} & \dots & a_{1c_1} & \dots & a_{k1} & a_{k2} & \dots & a_{kc_k} \end{pmatrix}$$

and

$$\widetilde{W} = \begin{pmatrix} 1_{(1)} & 2_{(1)} & \dots & k_{(1)} & k & k & \dots & k & \dots & 1 & 1 & \dots & 1 \\ b_{k1} & b_{k-1,1} & \dots & b_{11} & b_{12} & b_{13} & \dots & b_{1c_1} & \dots & b_{k2} & b_{k3} & \dots & b_{kc_k} \end{pmatrix}.$$

Proof. As in the proof of Lemma 3.4 which depends mostly on Lemma 3.3, we apply Lemma 3.10 (which is like the tableaux version of Lemma 3.3) to get the result.

Given L , by Lemma 3.10, we can get \widetilde{L}_{k-1} and if we ignore all the 1 entries (including $1_{(1)}$) in \widetilde{L}_{k-1} , we get a new column recording tableau L' with entries $2, \dots, k$, and we apply Lemma 3.10 again (treating r in L' as $r - 1$ in L when we apply Lemma 3.10) and get a \widetilde{L}'_{k-2} with an entry $2_{(1)}$. By changing the corresponding entries of \widetilde{L} with those in \widetilde{L}'_{k-2} , we have $2_{(2)}$ in a cell after the cell containing $1_{(1)}$ in reading order (because $a_{k1}^{(k-1)} < a_{k-1,1}^{(k-1)}$ and apply Lemma 15 of [4]). Result follows by repeating this process until $(k-1)_{(1)}$ is formed and then convert the first k as $k_{(1)}$. \square

Example 16. We use the same word in Example 7 and the same U in Example 13, which is also the U and the word we used as w_0 in Example 15, to illustrate Lemma 3.11.


By Example 11,


$$\widetilde{W} = \begin{pmatrix} 1_{(1)} & 2_{(1)} & 3_{(1)} & 4_{(1)} & 5_{(1)} & 5 & 5 & 5 & 5 & 5 & 4 & 4 & 4 & 4 & 3 & 3 & 3 \\ 1 & 3 & 6 & 8 & 9 & 8 & 7 & 5 & 4 & 3 & 9 & 6 & 5 & 4 & 7 & 6 & 6 \end{pmatrix}.$$

We continue Example 15 to illustrate Lemma 3.11 to get \widetilde{L} .

We mark the cell which is being moved and added subscript (1) as **green** and the the cells under consideration to get the new position of the green cell as **yellow** and mark the final cell of the subscripted green cell as **red**.

By Example 15, we get the position $1_{(1)}$ and we continue using the same process:







□


Example 17. We continue with the \tilde{L} in Example 16 to illustrate Lemma 3.12.

We mark the final position of the subscripted cells a different color for a different subscript.


Starting with what we get in Example 16:







Now by replacing the entry of each cell with the subscript number, we get



One can verify that R can be obtained by using the row word mentioned in Example 7.

3.3 Decomposition of the product of a dominating monomial and an atom into a positive sum of atoms

We now prove Theorem 3.1 mentioned in the beginning of this Chapter. We rephrase the Theorem in a more precise form as follows.

Theorem 3.13. *Let λ be a partition and α be a weak composition. Let \mathcal{A}_λ and \mathcal{A}_α be atoms of shape λ and α respectively. Then*

$$\mathcal{A}_\lambda \cdot \mathcal{A}_\alpha = \sum_{\beta=|\lambda|+|\alpha|, \lambda \subseteq \beta} c_{\lambda\alpha}^\beta \mathcal{A}_\beta$$

where $c_{\lambda\alpha}^\beta$ is the number of distinct LRS of shape β/λ created by column words whose corresponding SSAF has shape α .

Proof. Since $\mathcal{A}_\lambda = x^\lambda$ (there is exactly one SSAF with shape λ , denoted by U_λ) and $\mathcal{A}_\alpha = \sum_{F \in \text{SSAF}(\alpha)} x^F$, we have

$$\mathcal{A}_\lambda \cdot \mathcal{A}_\alpha = \sum_{F \in \text{SSAF}(\alpha)} x^\lambda x^F.$$

To prove the theorem, we only need to check that given a LRS L of shape β/λ created by column words whose corresponding SSAF has shape α , if there is some column word

$$w = a_{11} \dots a_{1c_1} | a_{21} \dots a_{2c_2} | \dots | a_{k1} \dots a_{kc_k}$$

and biword

$$W = \begin{pmatrix} k & k & \dots & k & \dots & 1 & 1 & \dots & 1 \\ a_{11} & a_{12} & \dots & a_{1c_1} & \dots & a_{k1} & a_{k2} & \dots & a_{kc_k} \end{pmatrix}$$

such that $(U_\lambda \leftarrow W)$ creates the same L , then the SSAF corresponding to w also has shape α .

First consider the last cell in reading order among all those containing entry k in L , that means it is the very first entry inserted. Since a_{11} must be inserted in a cell immediately above the cell (including those in basement) containing a_{11} , the position of that cell fixes the value of a_{11} .

Now consider all the cells with k and also the last cell in reading order among all those containing the entry $k - 1$, then these cells are created by $U_\lambda \leftarrow a_{11} \dots a_{1c_1} a_{21}$. By Lemma 3.12, we know the corresponding row recording tableau and hence we know which two cells are the first two entries being inserted using the corresponding row word. Note that the first row consists of distinct entries and is inserted in ascending order using the row word, then by Lemma 15 in [4], we know the cells are created in ascending reading order (one after another in reading order) and so they must be the cells immediately above those in U_λ , and hence the value inside each of those cells in the SSAF created by inserting the row word into U_λ is exactly the value inside the cell just below it. Hence we know what the first two row entries of the corresponding SSAF of $a_{11} a_{12} \dots a_{1c_1} a_{21}$ are. Since we already know the first row entry, which is the lowest entry of the column corresponding to $a_{11} \dots a_{1c_1}$ is, we now know what the lowest entry of the second column (corresponding to $a_{11} \dots a_{1c_1} a_{21}$ and hence the same for $a_{11} \dots a_{1c_1} a_{21} \dots a_{2c_2}$) is.

We can repeat the same process until we get all the last entries of the k columns and hence fix the shape of the SSAF corresponding to w . Since we read those entries just by considering L , this shows that L fixes the shape of the corresponding SSAF of w and result follows. □

Example 18. Pick $\lambda = 4332221$ and $\alpha = (1, 0, 1, 0, 0, 4, 0, 6, 5)$.

Let $\beta = (5, 3, 5, 2, 4, 6, 1, 6, 2)$.

Let $U =$

1									
1	2	3							
1	2	3	4	5	6				
1	2	3	4	5	6	7			
	1	2	3	4	5	6	7	8	9

One can first check that a recording tableaux L (which is also an LRS if we change the basement) of shape β/λ is created when the column word in Example 7 whose corresponding SSAF has shape α is inserted into U . Indeed, we have

$L =$

						5		4	
						5		4	
3					1	5		4	
		2			3	5		4	
		3						5	3
								5	4
	1	2	3	4	5	6	7	8	9


If a column word w would create the same L when inserted in U , we know that it has column lengths 6, 5, 4, 1, 1, and hence we can break it into 5 subsequences:

$$w = a_{11} \dots a_{16} | a_{21} \dots a_{25} | a_{31} \dots a_{34} | a_{41} | a_{51}$$

satisfying the conditions of being a column word in Definition 3.3. Let $F(w)$ be the SSAF corresponding to w , i.e. the SSAF created when inserting w into an empty SSAF with basement being


1 2 3 4

Consider the cells with entry 5 in L :




the largest in reading order (marked as red) is created when a_{11} is inserted in U . Note that as U is a partition shaped SSAF, a_{11} must be placed immediately above the cell (including basement) containing the entry a_{11} . Hence a_{11} must be 8. That means the column in $F(w)$ corresponding to $a_{11} \dots a_{16}$ is above the basement entry 8.

Next consider the cells with entry 5 and the last cell in reading order containing 4:




by applying Lemma 3.11, we know the first two cells created when inserting the row word corresponding to the column word $a_{11} \dots a_{16}a_{21}$ must be the ones marked red as below:



and so we know the two numbers in the row word must be 8 and 9, and since we already know 8 is the basement entry which the new cell created is above, when inserting $a_{11} \dots a_{16}$ and hence we know the second column of $F(w)$ when inserting $a_{21} \dots a_{25}$ is above the basement entry 9.


Apply Lemma 3.11 on the cells with entries 4, 5 and also the last cell in reading order with entry

3, we know the first three cells created when inserting the corresponding row word of the column word $a_{11} \dots a_{16} a_{21} \dots a_{25} a_{31}$ into U , and we mark them red as shown:



That means the first three numbers in the corresponding row word is 6, 8, 9 and since we already know the first two columns are above basement entries 8 and 9, we can conclude that the third column created in $F(w)$ when inserting $a_{31} \dots a_{34}$ is above the basement entry 6.

Continue with the same process and we have:



Therefore we know the shape of $F(w)$ is $(1, 0, 1, 0, 0, 4, 0, 6, 5)$ which is exactly α .

3.4 Decomposition of the product of a dominating monomial and a key into a positive sum of keys

We would adapt the notations and apply the results in [3] to prove the key-positivity property of the product of a dominating monomial and a key. However, we will still use $\bar{\alpha}$ instead of α^* which [3] uses to denote the reverse of α .

Lemma 3.14. For any given partition λ and weak compositions β, γ such that $\lambda \subseteq \beta$, Let

$$S_1 := \left\{ K \left| \begin{array}{l} K \text{ is a LRK of shape } \bar{\delta}/\lambda \text{ with content created } \overline{\omega_\gamma(\gamma)} \text{ and } \phi(K) \text{ is an LRS created} \\ \text{by a column word whose corresponding SSAF has a shape } \geq \bar{\gamma} \text{ and basement shape} \end{array} \right. \right\},$$

$$S_2 := \left\{ L \left| \begin{array}{l} \lambda, \bar{\delta} \leq \beta \\ L \text{ is an LRS of shape } \beta/\lambda \text{ created by a column word whose corresponding SSAF} \\ \text{has shape } \alpha, \alpha \geq \bar{\gamma} \end{array} \right. \right\}$$

(where ϕ is defined in the proof in Theorem 6.1 in [3] (see appendix). Then $\phi|_{S_1} : S_1 \rightarrow S_2$ is a bijection.)

Proof. Let $K \in S_1$. Then by fixing β , we know $\phi(K)$ has an overall shape β . Hence $\phi(K)$ has shape β/λ and thus $\phi(K) \in S_2$.

Therefore we have $\phi(S_1) \subseteq S_2$.

Since ϕ and hence $\phi|_{S_1}$ is injective, it remains to check $\phi|_{S_1} : S_1 \rightarrow S_2$ is surjective.

Let $L \in S_2$. Since L is created by a column word whose corresponding SSAF has shape α and $\alpha \geq \bar{\gamma}$, we know L has content $\overline{\omega_\gamma(\gamma)}$. Then by the proof in Theorem 6.1 in [3], $\phi^{-1}(L)$ is an LRK of shape $\bar{\delta}/\lambda$ for some $\bar{\delta} \leq \beta$ and has the same column sets as L . Hence $\phi^{-1}(L) \in S_1$. Therefore for any $L \in S_2$, we can find an LRK, namely, $\phi^{-1}(L) \in S_1$ such that $\phi|_{S_1}(\phi^{-1}(L)) = L$.

We thus have $\phi|_{S_1}$ is surjective and result follows. \square

Theorem 3.15. *Let λ be a partition and γ be a weak composition. Let \mathcal{A}_λ and κ_γ be an atom of shape λ and a key of shape γ respectively. Then*

$$\mathcal{A}_\lambda \cdot \kappa_\gamma = \sum_{\alpha} b_{\lambda\gamma}^{\alpha} \kappa_{\alpha}$$

where $b_{\lambda\gamma}^{\alpha}$ is the number of distinct LRK of shape $\bar{\alpha}/\lambda$ with content $\overline{\omega_\gamma(\gamma)}$ and the image under ϕ is an LRS with basement shape λ created by a column word whose corresponding SSAF has shape ρ for some $\rho \geq \bar{\gamma}$.

Proof. By Theorem 1 and Theorem 3.13, we have

$$\mathcal{A}_\lambda \cdot \kappa_\gamma = \mathcal{A}_\lambda \cdot \sum_{\alpha \geq \bar{\gamma}} \mathcal{A}_\alpha = \sum_{\alpha \geq \bar{\gamma}} (\mathcal{A}_\lambda \cdot \mathcal{A}_\alpha) = \sum_{\alpha \geq \bar{\gamma}} \sum_{\substack{\beta = |\lambda| + |\alpha| \\ \beta \supseteq \lambda}} c_{\lambda\alpha}^\beta \mathcal{A}_\beta = \sum_{\beta \supseteq \lambda} \sum_{\substack{\alpha \geq \bar{\gamma} \\ \alpha = |\beta| - |\lambda|}} c_{\lambda\alpha}^\beta \mathcal{A}_\beta.$$

Also, by Theorem 1, we have

$$\sum_{\delta} b_{\lambda\gamma}^\delta \kappa_\delta = \sum_{\delta} \sum_{\beta \geq \bar{\delta}} b_{\lambda\gamma}^\delta \mathcal{A}_\beta = \sum_{\beta} \left(\sum_{\bar{\delta} \leq \beta} b_{\lambda\gamma}^\delta \right) \mathcal{A}_\beta.$$

Hence to prove the theorem, we only need to prove

$$\sum_{\substack{\alpha \geq \bar{\gamma} \\ \alpha = |\beta| - |\lambda|}} c_{\lambda\alpha}^\beta = \sum_{\beta \geq \bar{\delta}} b_{\lambda\gamma}^\delta$$

which follows from Lemma 3.14 as $|S_2| = \sum_{\substack{\alpha \geq \bar{\gamma} \\ \alpha = |\beta| - |\lambda|}} c_{\lambda\alpha}^\beta$ and $|S_1| = \sum_{\beta \geq \bar{\delta}} b_{\lambda\gamma}^\delta$

□

3.5 Decomposition of the product of a Schur function and a Demazure character

It is proved in [3] that the product of a Schur function and a Demazure character is key-positive using tableaux-combinatorics. In this section, we give another proof using linear operators.

Lemma 3.16. *If f is atom-positive, then $\pi_i f$ is also atom positive for all positive integers i .*

Proof. Let $f = A_0(x) + \sum_{\substack{I, \\ I \text{ is a reduced word}}} \theta_I A_I(x)$ where $A_0(x) = \sum_{\lambda \in Par} a_\lambda x^\lambda$ with $a_\lambda \in \mathbb{Z}_{\geq 0}$ and $A_I(x) = \sum_{\lambda \in Par} a_\lambda^I x^\lambda$ with $a_\lambda^I \in \mathbb{Z}_{\geq 0}$ for any reduced word I .

Let σ_I be the permutation corresponding to a reduced word I . Then by Lemma 1.7, $|l(s_i \sigma_I) - l(\sigma)| = 1$. If $l(s_i \sigma_I) = l(\sigma_I) - 1$, then by Lemma 1.8, there exists a reduced word of σ_I starting with i . Hence $\pi_i \theta_I = 0$ by item 5. of Proposition 2.1. If $l(s_i \sigma_I) = l(\sigma_I) + 1$, then iI is also a reduced word and hence $\pi_i \theta_I = (1 + \theta_i) \theta_I = \theta_I + \theta_{iI}$.

As a result,

$$\pi_i f = \pi_i(A_0(x) + \sum_{I \text{ is a reduced word}} A_I(x)) = A_0(x) + \theta_i(A_0(x)) + \sum_{iI \text{ is a reduced word}} (\theta_I A_I(x) + \theta_{iI} A_I(x)).$$

Since $A_0(x)$ and $A_I(x)$ are sums of dominating monomials with nonnegative integer coefficients, $\pi_i f$ is also atom positive. □

Lemma 3.17. *Let $\sigma = n, n-1, \dots, 1 \in S_n$. Then $\pi_i \pi_\sigma = \pi_\sigma$ and hence $\theta_i \sigma = 0$.*

Proof. By Corollary 1.10, there exists a reduced word of σ starting with i and hence by item 5. of Proposition 2.1, we have $\pi_i \pi_i = \pi_i$ which implies $\pi_i \pi_\sigma = \pi_\sigma$.

$$\text{Hence } \theta_i \pi_\sigma = (\pi_i - 1) \pi_\sigma = \pi_i \pi_\sigma - \pi_\sigma = \pi_\sigma - \pi_\sigma = 0. \quad \square$$

Theorem 3.18. *The product of a key and a Schur function is key positive.*

Proof. Let $\sigma = n, n-1, \dots, 1 \in S_n$ and λ, μ be partitions with length at most n . By item 2 of Theorem 2.8, we have $s_\lambda = \pi_\sigma(x^\lambda)$.

Let $I = i_1 i_2 \dots i_k$ be a reduced word of some permutation in S_n . We prove that $\pi_I(x^\mu) \times \pi_\sigma(x^\lambda)$ is key positive.

By Lemma 2.6,

$$\begin{aligned} & \pi_{i_k}(x^\mu \times \pi_\sigma(x^\lambda)) \\ &= \pi_{i_k}(x^\mu) \times \pi_\sigma(x^\lambda) + s_{i_k}(x^\mu) \times (\theta_{i_k} \pi_\sigma(x^\lambda)) \\ &= \pi_{i_k}(x^\mu) \times \pi_\sigma(x^\lambda) \end{aligned}$$

$$\begin{aligned}
& \pi_{i_{k-1}} \pi_{i_k} (x^\mu \times \pi_\sigma(x^\lambda)) \\
= & \pi_{i_{k-1}} (\pi_{i_k} (x^\mu) \times \pi_\sigma(x^\lambda)) \\
= & (\pi_{i_{k-1}} \pi_{i_k} (x^\mu)) \times \pi_\sigma(x^\lambda) + s_{i_{k-1}} \pi_{i_k} (x^\mu) \times \theta_{i_{k-1}} \pi_\sigma(x^\lambda) \\
= & (\pi_{i_{k-1}} \pi_{i_k} (x^\mu)) \times \pi_\sigma(x^\lambda)
\end{aligned}$$

Inductively, we get

$$\pi_I(x^\mu) \times \pi_\sigma(x^\lambda) = \pi_{i_1} \pi_{i_2} \cdots \pi_{i_k} (x^\mu) \times \pi_\sigma(x^\lambda) = \pi_{i_1} \pi_{i_2} \cdots \pi_{i_k} (x^\mu \times \pi_\sigma(x^\lambda)).$$

By Lemma 2.5, $\pi_\sigma(x^\lambda) = \sum_{\gamma \leq \sigma} \theta_\gamma(x^\lambda) = \sum_{\gamma \in S_n} \theta_\gamma(x^\lambda)$ which implies $x^\mu \times \pi_\sigma(x^\lambda) = \sum_{\gamma \in S_n} x^\mu \times \theta_\gamma(x^\lambda)$. Therefore by Theorem 3.13, $x^\mu \times \pi_\sigma(x^\lambda)$ is atom positive. By Lemma 3.16, we know $\pi_{i_k} (x^\mu \times \pi_\sigma(x^\lambda))$ is atom-positive. Inductively applying Lemma 3.16, $\pi_{i_1} \pi_{i_2} \cdots \pi_{i_k} (x^\mu \times \pi_\sigma(x^\lambda))$ is also atom-positive. \square

Chapter 4

Atom positivity of the product of two key polynomials whose basements have length at most 3

In this section, we prove Conjectue 1 for $l(\alpha), l(\beta) \leq 3$. Note that if $l(\alpha) < 3$, we can always add zero parts to it to increase the length to 3. Hence, we can assume $l(\alpha) = 3$. Similarly, we can assume $l(\beta) = 3$.

Let $\lambda_\alpha = \omega_\alpha(\alpha)$ and $\lambda_\beta = \omega_\beta(\beta)$. We claim that we can consider $l(\lambda_\alpha), l(\lambda_\beta) \leq 2$, i.e. both α and β have at least one zero part.

First note that for integers $r \geq 0$ and $a \geq b \geq c \geq 0$, $(x_1 x_2 x_3)^r \theta_\tau(x_1^a x_2^b x_3^c) = \theta_\tau(x_1^{a+r} x_2^{b+r} x_3^{c+r})$ and $\pi_\tau(x_1^a x_2^b x_3^c) = (x_1 x_2 x_3)^c \pi_\tau(x_1^{a-c} x_2^{b-c})$ for any $\tau \in S_3$. That means the monomial $(x_1 x_2 x_3)^r$ times any atom is still an atom, and same for the case for key. We can also interpret this by considering fillings, as multiplying $(x_1 x_2 x_3)^r$ to an atom or a key is just adding r bottom rows to the diagram and there is only one way to fill in these cells in an atoms or a key.

Suppose Conjectue 1 is true for $l(\lambda_\alpha), l(\lambda_\beta) \leq 2$. Then for any $\omega, \tau \in S_3$ and integers $a \geq b \geq$

$c \geq 0, s \geq t \geq u \geq 0,$

$$\begin{aligned}
\pi_\omega(x_1^a x_2^b x_3^c) \cdot \pi_\tau(x_1^s x_2^t x_3^u) &= (x_1 x_2 x_3)^{c+u} (\pi_\omega(x_1^{a-c} x_2^{b-c}) \cdot \pi_\tau(x_1^{s-u} x_2^{t-u})) \\
&= (x_1 x_2 x_3)^{c+u} \left(\sum_{\gamma} c_{\gamma} \mathcal{A}_{\gamma} \right) \\
&= \sum_{\gamma} c_{\gamma} ((x_1 x_2 x_3)^{c+u} \mathcal{A}_{\gamma}) \\
&= \sum_{\gamma} c_{\gamma} \mathcal{A}_{\gamma'}
\end{aligned}$$

where c_{γ} are all nonnegative integers and $(x_1 x_2 x_3)^{c+u} \mathcal{A}_{\gamma} = \mathcal{A}_{\gamma'}$ with $\gamma' = \gamma + (c+u, c+u, c+u)$ (i.e. γ' can be obtained by adding $c+u$ to each part of γ).

As a result, we now only consider compositions of length 3 and with at most two nonzero parts.

4.1 Polytopes


In this section, we introduce another way to view Demazure atoms and characters.

For each weak composition with length k , we can view it as a lattice point in $\mathbb{Z}_{\geq 0}^k$. We will focus on the case $k = 3$. (Everything in this section applies for any positive integer k .) Hence we have the bijection:

$$\alpha \leftrightarrow (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{Z}_{\geq 0}^3 \leftrightarrow x^{\alpha} = x_1^{\alpha_1} x_2^{\alpha_2} x_3^{\alpha_3}.$$

Consider the Coxeter arrangement (of type A_2) in \mathbb{R}^3 :

$$\mathbf{Cox}(3) = \{a_i - a_j : 1 \leq i < j \leq 3\}$$



$$R_1 : a_1 \geq a_2 \geq a_3;$$

$$R_2 : a_1 \geq a_3 \geq a_2;$$

$$R_3 : a_2 \geq a_1 \geq a_3;$$

$$R_4 : a_3 \geq a_1 \geq a_2;$$

$$R_5 : a_2 \geq a_3 \geq a_1;$$

$$R_6 : a_3 \geq a_2 \geq a_1;$$


4.1.1 Demazure characters and polytopes

Let α be a weak composition with $\lambda_\alpha = (m, n, 0)$, for some integers $m \geq n \geq 0$. Then there are exactly 6 key polynomials obtained from λ_α , namely:

$$x^{\lambda_\alpha}, \pi_1 x^{\lambda_\alpha}, \pi_2 x^{\lambda_\alpha}, \pi_{21} x^{\lambda_\alpha}, \pi_{12} x^{\lambda_\alpha}, \pi_{121} x^{\lambda_\alpha}.$$


We now plot each of these 6 key polynomials in the Coxeter arrangement:

Case 1. $x^{\lambda_\alpha} = x_1^m x_2^n$, so it corresponds to the point $(m, n, 0)$ in R_1 :




Case 2. $\pi_1 x^{\lambda_\alpha} = \pi_1(x_1^m x_2^n) = x_1^m x_2^n + x_1^{m-1} x_2^{n+1} + \dots + x_1^n x_2^m$

Therefore it corresponds to the line joining $(m, n, 0)$ and $(n, m, 0)$. That is, each monomial corresponds to a lattice point (and vice versa) on the line obtained by joining $(m, n, 0)$ and its reflection along $a_1 = a_2$.



Case 3. $\pi_2 x^{\lambda_\alpha} = \pi_2(x_1^m x_2^n) = x_1^m x_2^n + x_1^m x_2^{n-1} x_3 + \dots + x_1^m x_3^n$


Therefore it corresponds to the line joining $(m, n, 0)$ and $(m, 0, n)$. That is, each monomial corresponds to a lattice point (and vice versa) on the line obtained by joining $(m, n, 0)$ and its reflection along $a_2 = a_3$.




Case 4. $\pi_{21}x^{\lambda_\alpha} = \pi_2\pi_1(x_1^m x_2^n) = \pi_2(x_1^m x_2^n + x_1^{m-1} x_2^{n+1} + \dots + x_1^n x_2^m)$

Since $\pi_2(x_1^m x_2^n + x_1^{m-1} x_2^{n+1} + \dots + x_1^n x_2^m) = \pi_2(x_1^m x_2^n) + \pi_2(x_1^{m-1} x_2^{n+1}) + \dots + \pi_2(x_1^n x_2^m)$, we can apply the same correspondence in Case 3. for each key in the summand. Therefore $\pi_{21}x^{\lambda_\alpha}$ corresponds to the $m - n + 1$ lines obtained by reflecting each lattice point on the line joining $(m, n, 0)$ and $(m, 0, n)$ along $a_2 = a_3$. That is, each monomial corresponds to a lattice point (and vice versa) in the trapezoid obtained by first reflecting $(m, n, 0)$ along the line $a_1 = a_2$ followed by reflecting the resulting line along $a_2 = a_3$ as shown below:


First reflect along $a_1 = a_2$ and get a line joining $(m, n, 0)$ and $(m, 0, n)$:



Then reflect the line along $a_2 = a_3$:



Note that there is another case where the trapezoid does not have any point in R_5 or R_6 :




Indeed, the trapezoid has at least a point in R_5 or R_6 if and only if $m \geq 2n$ (one can prove that by locating the mid-point of the line joining $(n, m, 0)$ and $(n, 0, m)$.)


Case 5. $\pi_{12}x^{\lambda_\alpha} = \pi_1\pi_2(x_1^m x_2^n) = \pi_1(x_1^m x_2^n + x_1^m x_2^{n-1} x_3 + \cdots + x_1^m x_3^n)$

Similar to Case 4., it corresponds to the lattice points in the trapezoid formed by first reflecting $(m, n, 0)$ along $a_2 = a_3$ followed by reflecting the resulting line along $a_1 = a_2$. The trapezoid has at least a point in R_4 or R_6 if and only if $2n \geq m$. The trapezoid is as follows (we just shade the region using dotted pattern for convenience, but the actual correspondence should be lattice points in the shaded region including the boundary):

For $2n < m$:



For $2n \geq m$:




Case 6. $\pi_{121}x^{\lambda\alpha} = \pi_1\pi_2\pi_1(x_1^m x_2^n) = \pi_1(\pi_2\pi_1(x_1^m x_2^n))$


First recall that by Proposition 2.1 $\pi_i s_i = (1 + \theta_i)s_i = s_i + \theta_i s_i = s_i - \theta_i$, hence $\pi_i s_i + \pi_i = s_i - \theta_i + (1 + \theta_i) = s_i + 1$.

For example, when $i = 1$, $\pi_1(x_1^a x_2^b) + \pi_1(x_1^b x_2^a) = x_1^a x_2^b + x_1^b x_2^a$ for any and any integers $a \geq b \geq 0$. Hence $\pi_1(x_1^b x_2^a) = x_1^a x_2^b + x_1^b x_2^a - \pi_1(x_1^a x_2^b)$.

We can plot $\pi_1(x_1^b x_2^a)$ as:




and $\pi_1(x_1^a x_2^b) + \pi_1(x_1^b x_2^a)$ as




Now consider $\pi_{121}(x_1^m x_2^n) = \pi_1(\pi_{21}(x_1^m x_2^n))$.

By Case 4., we have $\pi_{21}(x_1^m x_2^n)$ as a trapezoid as follows:




Apply π_1 to each lattice points in the trapezoid is equivalent to reflecting the trapezoid along $a_1 = a_2$ and get a hexagon with multiplicities on the points. Consider the multiplicity of a lattice point (a_1, a_2, a_3) in the hexagon with $a_1 \geq a_2$. The multiplicity of its ' $a_1 = a_2$ '-reflection, namely, (a_2, a_1, a_3) is the same. Recall that for each ' $a_1 = a_2$ ' - reflection pair in the trapezoid region (i.e. (a, b, c) and (b, a, c)), applying π_1 to both of them results in the two points themselves. So the multiplicity of the lattice points increases by 1 'horizontally' from the boundary, starting from 1, until it first hits the line joining $(m, n, 0)$ and $(0, n, m)$ or the line joining $(n, m, 0)$ and $(n, 0, m)$ (i.e. hit either of the lines in region $a_1 > a_2$), then it becomes stable. Here is an example for $m \geq 2n$:



The monomials corresponding to the lattice points on the red boundary have coefficient 1 in the key polynomial, and those monomial corresponding to the lattice points on the blue boundary have coefficient 2 and so on, while those corresponding to the lattice points in the inner most triangle (the orange region) have coefficient $n + 1$ for $m \geq 2n$ (and $m - n + 1$ if $m \leq 2n$).

If we also plot the multiplicity (with xy -plane being the Coxeter arrangement and z -axis represents the multiplicity), we get a polytope as follows:



Note: One can also start with π_{12} and apply π_2 on each lattice point in the trapezoid corresponding to π_{12} and form π_{212} to check that $\pi_{121} = \pi_{212}$.

4.1.2 Demazure atoms and polytopes


In this section, we will discuss how one can obtain a polytope from a Demazure atom. Again, we focus on the case where the shape of the atom is a weak composition of length 3 with at least one zero part.

Let α be a weak composition with $\lambda_\alpha = (m, n, 0)$, for some integers $m \geq n \geq 0$. Then there are exactly 6 key polynomials obtained from λ_α , namely:

$$x^{\lambda_\alpha}, \theta_1 x^{\lambda_\alpha}, \theta_2 x^{\lambda_\alpha}, \theta_{21} x^{\lambda_\alpha}, \theta_{12} x^{\lambda_\alpha}, \theta_{121} x^{\lambda_\alpha}.$$


We now plot each of these 6 Demazure atoms in the Coxeter arrangement:

Case 1. $x^{\lambda_\alpha} = x_1^m x_2^n$, so it corresponds to the point $(m, n, 0)$ in R_1 :



Case 2. $\theta_1 x^{\lambda_\alpha} = \theta_1(x_1^m x_2^n) = x_1^{m-1} x_2^{n+1} + x_1^{m-2} x_2^{n+2} + \dots + x_1^n x_2^m$


Hence each monomial corresponds to a lattice point except $(m, n, 0)$ (and vice versa) on the line obtained by joining $(m, n, 0)$ and its reflection along $a_1 = a_2$.



This also shows that $\pi_1 = \theta_1 + 1$

Case 3. $\theta_2 x^{\lambda_\alpha} = \theta_2(x_1^m x_2^n) = x_1^m x_2^{n-1} x_3 + x_1^m x_2^{n-2} x_3^2 + \dots + x_1^m x_3^n$

Therefore it corresponds to the line joining $(m, n, 0)$ and $(m, 0, n)$ excluding $(m, n, 0)$. That is, each monomial corresponds to a lattice point except $(m, n, 0)$ (and vice versa) on the line obtained by joining $(m, n, 0)$ and its reflection along $a_2 = a_3$.




This also shows $\pi_2 = \theta_2 + 1$


Case 4. $\theta_{21}x^{\lambda_\alpha} = \theta_2\theta_1(x_1^m x_2^n) = \theta_2(x_1^{m-1}x_2^{n+1} + x_1^{m-2}x_2^{n+2} + \dots + x_1^n x_2^m)$

Since $\theta_2(x_1^{m-1}x_2^{n+1} + x_1^{m-2}x_2^{n+2} + \dots + x_1^n x_2^m) = \theta_2(x_1^{m-1}x_2^{n+1}) + \theta_2(x_1^{m-2}x_2^{n+2}) + \dots + \theta_2(x_1^n x_2^m)$, we can apply the same correspondence in Case 3. for each atom in the summand. Therefore $\theta_{21}x^{\lambda_\alpha}$ corresponds to the $m - n + 1$ lines obtained by reflecting each lattice point except $(m, n, 0)$ on the line joining $(m, n, 0)$ and $(m, 0, n)$ along $a_2 = a_3$. That is, each monomial corresponds to a lattice point (and vice versa) in the ‘semi-open’ trapezoid obtained by first reflecting $(m, n, 0)$ along the line $a_1 = a_2$ followed by reflecting the resulting line along $a_2 = a_3$ as shown below:


First reflect along $a_1 = a_2$ and get a line joining $(m, n, 0)$ and $(m, 0, n)$:



Then reflect the line along $a_2 = a_3$:




Note that there is another case where the trapezoid does not have any point in R_5 or R_6 :



As Case 4 in Section 4.1.1, the trapezoid has at least a point in R_5 or R_6 if and only if $m \geq 2n$ (one can prove that by locating the mid-point of the line joining $(n, m, 0)$ and $(n, 0, m)$.)


Again, we can illustrate $\pi_{21} = 1 + \theta_1 + \theta_2 + \theta_{21}$:




Case 5. $\theta_{12}x^{\lambda\alpha} = \theta_1\theta_2(x_1^m x_2^n) = \theta_1(x_1^m x_2^{n-1}x_3 + x_1^m x_2^{n-2}x_3^2 + \dots + x_1^m x_3^n)$

Similar to Case 4., it corresponds to the lattice points in the semi-open trapezoid formed by first reflecting $(m, n, 0)$ along $a_2 = a_3$ followed by reflecting the resulting line along $a_1 = a_2$. The trapezoid has at least a point in R_4 or R_6 if and only if $2n \geq m$. The trapezoid is as follows (we just shade the region using dotted pattern for convenience, but the actual correspondence should be lattice points in the shaded region including the solid boundaries.):


For $2n < m$:



For $2n \geq m$:



We again illustrate how to decompose π_{12} into $1 + \theta_1 + \theta_2 + \theta_{12}$.




Case 6. $\theta_{121}x^{\lambda\alpha} = \theta_1\theta_2\theta_1(x_1^m x_2^n) = \theta_1(\theta_2\theta_1(x_1^m x_2^n))$

First recall that by Proposition 2.1 $\theta_i s_i = -\theta_i$, hence $\theta_i s_i + \theta_i = 0$.


For example, when $i = 1$, $\theta_1(x_1^a x_2^b) + \theta_1(x_1^b x_2^a) = 0$ for any and any integers $a \geq b \geq 0$.

We can plot $\theta_1(x_1^b x_2^a)$ as:




Now consider $\theta_{121}(x_1^m x_2^n) = \theta_1(\theta_{21}(x_1^m x_2^n))$.

By Case 4., we have $\theta_{21}(x_1^m x_2^n)$ as a trapezoid as follows:



Apply θ_1 to each lattice points in the semi-open trapezoid is equivalent to reflecting the trapezoid along $a_1 = a_2$ and get a hexagon with multiplicities on the points. Recall that each " $a_1 = a_2'$ - reflection pair in the trapezoid region (i.e. (a, b, c) and (b, a, c)) vanishes under θ_1 . Hence the multiplicity stays constant along the horizontal line perpendicular to the line $a_1 = a_2$. Here is an example for $m \geq 2n$:



The monomials corresponding to the lattice points on the red boundary have coefficient 1 in the key polynomial, and those monomial corresponding to the lattice points on the blue boundary have coefficient 2 and so on, while those corresponding to the lattice points in the inner most triangle (the orange region) have coefficient n for $m \geq 2n$ (and $m - n$ for $m \leq 2n$). If we also plot the multiplicity, with xy -plane being the Coxeter arrangement and z -coordinates being the multiplicity, we get a polytope similar to the one in Case 6 in Section 4.1.1 but with different heights (as not all the multiplicity of the monomials in an atom is the same as those in keys).

One can verify that $\theta_{121} = \theta_{212}$ by starting with θ_{12} instead. Also one can get the decomposition $\pi_{121} = 1 + \theta_1 + \theta_2 + \theta_{12} + \theta_{21} + \theta_{121}$ by putting the figures shown in this section together.

4.2 Products of two Demazure characters

Let $m \geq n \geq 0$ and $k \geq l \geq 0$ be integers.

	$x_1^k x_2^l$	$\pi_1(x_1^k x_2^l)$	$\pi_2(x_1^k x_2^l)$	$\pi_{21}(x_1^k x_2^l)$	$\pi_{12}(x_1^k x_2^l)$	$\pi_{121}(x_1^k x_2^l)$
$x_1^m x_2^n$	3.13	3.13	3.13	3.13	3.13	3.13/3.18
$\pi_1(x_1^m x_2^n)$		(i)	4.2.1	4.2.2	(ii)	3.18
$\pi_2(x_1^m x_2^n)$			(iii)	(iv)	4.2.3	3.18
$\pi_{21}(x_1^m x_2^n)$				(v)	4.2.4	3.18
$\pi_{12}(x_1^m x_2^n)$					(vi)	3.18
$\pi_{121}(x_1^m x_2^n)$						3.18

Table 4.1: Decomposition of products of keys into atoms

In the following sections, we will first state the result and one can verify by expanding both sides directly. We will state some other methods (either using operators or polytopes) to verify or interpret the decomposition in the first two sections. These methods are applicable to all cases.


$$\text{Also, we denote the indicator function as } \mathbb{1}_S = \begin{cases} 1 & \text{if } ((m, n), (k, l)) \in S \\ 0 & \text{otherwise.} \end{cases}$$

4.2.1 $\pi_1(x_1^m x_2^n) \times \pi_2(x_1^k x_2^l)$


$$\begin{aligned} \pi_1(x_1^m x_2^n) \times \pi_2(x_1^k x_2^l) = & \sum_{s=0}^{\min\{m-n, k\}} \sum_{t=\max\{0, s-(k-l)\}}^{\min\{l, s+n\}} x_1^{m+k-s} x_2^{n+l+s-t} x_3^t \\ & + \mathbb{1}_{\{m-n > k-l\}} \sum_{t=0}^{\min\{l, (m-n)-(k-l)\}} \theta_1(x_1^{m+l-t} x_2^{k+n} x_3^t) \\ & + \mathbb{1}_{\{l > n\}} \sum_{s=0}^{\min\{l-n, m-n\}} \theta_2(x_1^{m+k-s} x_2^l x_3^{n+s}) \end{aligned}$$

We can also use polytopes to verify the atom positivity of $\pi_1(x_1^m x_2^n) \times \pi_2(x_1^k x_2^l)$.

The polytope corresponding to $\pi_1(x_1^m x_2^n)$ is the line:



The polytope corresponding to $\pi_2(a_1^k x_2^l)$ is the line:




Hence the product is a parallelogram:




There are different possible positions for the parallelogram:

1. The whole parallelogram lies in R_1 :




This case corresponds to $m - n \leq k - l$ and $l \leq n$ in the expansion.

2. The parallelogram lies in R_1 and R_3 :



Then we can decompose the parallelogram as:




Here the green region is obtained by applying θ_1 on each lattice point on the black line. Notice that the black line is in R_1 , meaning that every monomial corresponding to the a lattice point is a dominating monomial.

Hence we can decompose the parallelogram into the yellow region in R_1 and θ_1 of the black

line, all of which are positive sum of atoms.

This corresponds to the case when $m - n > k - l$ and $l \leq n$. Note that there are two different cases for the positions for the line $a_1 = a_2$, either $(n + k, m, l)$ is on the left of the line or on the right of the line. They correspond to the two cases in $\min\{l, (m - n) - (k - l)\}$ in the upper limit of the summation. In fact, one can locate the black line simply by flipping along $a_1 = a_2$ the boundary of the parallelogram which lies in R_3 .

3. The parallelogram lies in R_1 and R_2 :




This case corresponds to $m - n \leq k - l$ and $l > n$ in the expansion.

Similar to the previous case when the parallelogram lies in R_1 and R_3 , we can decompose the parallelogram into the yellow region which corresponds to a sum of dominating monomials and the green region which corresponds to θ_2 of a sum of dominating monomials (corresponding to the lattice points on the black line obtained by reflecting along the line $a_2 = a_3$ the ‘base’

of the parallelogram in R_2 .

Again there are also two cases: either $(n+k, m, l)$ is above or below the line $a_2 = a_3$, corresponding to the upper limit $\min\{l-n, m-n\}$ of the summation in the third summand in the expansion.

4. The parallelogram lies in R_1, R_2 and R_3 :



This case corresponds to $m-n > k-l$ and $l > n$ in the expansion and the three regions correspond to the three summands in the expansion.

4.2.2 $\pi_1(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l)$

We first write the decomposition of $(x_1^m x_2^n) \times \pi_{121}(x_1^k x_2^l)$ as follows:

$$(x_1^m x_2^n) \times \pi_{121}(x_1^k x_2^l) = A_0(x) + \theta_1 A_1(x) + \theta_2 A_2(x) + \theta_{21} A_{21}(x) + \theta_{12} A_{12}(x) + \theta_{121} A_{121}(x),$$

where $A_I(x) = \sum_{\lambda \in Par} a_\lambda^I x^\lambda$ with $a_\lambda^I \in \mathbb{Z}$ for $I \in \{0, 1, 2, 12, 21, 121\}$.


By Theorem 3.18, $(x_1^m x_2^n) \times \pi_{121}(x_1^k x_2^l)$ is key positive and hence atom positive.

i.e. $A_I(x)$ is a sum of dominating monomials with integer coefficients. Then


$$\begin{aligned} \pi_1(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l) = & A_0(x) + \sum_{r=0}^{\min\{m,k\}} \sum_{s=\max\{0,r-(n+l)\}}^{\min\{r,m-n,k-l\}} \theta_1(x_1^{m+k-r} x_2^{n+l+s} x_3^{r-s}) \\ & + \theta_2 A_2(x) + \mathbb{1}_{\{\min\{m,k\} \geq n+l\}} \sum_{r=n+l+1}^{\min\{m,k\}} \theta_{12}(x_1^{m+k-r} x_2^r x_3^{n+l}) \\ & + \theta_{21} A_{21}(x) + \mathbb{1}_{\{k>m>n+l\}} \theta_{121}(x_1^k x_2^m x_3^{n+l}). \end{aligned}$$

One can check the coefficients using polytopes as in Section 4.2.1. We will show a case where $k > m > n + l$ as an example (Note that even $k > m > n + l$ has several subcases). Other cases can be easily deduced similarly.


$\pi_{21}(x_1^k x_2^l)$ corresponds to:



So $\pi_1(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l)$ is equivalent to $m - n + 1$ trapezoids along the line perpendicular to $a_1 = a_2$ as follows:




We can draw the product as:



Here the number next to each line represents the multiplicity of the lattice points lying on that line (i.e. the number of trapezoids that the lattice point lies, which is also equal to the coefficient of the monomial corresponding to that lattice point) like what we have shown in Case 6. in Section 4.1.1. Also all the lattice points in the red region have the maximum multiplicity.

We can now decompose the polytope (with multiplicity) as follows:



One can check that the multiplicity of each lattice point in the original yellow region is at least the multiplicity of the sum of the multiplicities in all other colored regions. This ensures the remaining points in R_1 still corresponds to a positive sum of dominating monomials (i.e. A_0 is a positive sum of some dominating monomials).

We can also check some of the coefficients by using operators.

Suppose

$$\begin{aligned} & \pi_1(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l) \\ = & B_0(x) + \theta_1 B_1(x) + \theta_2 B_2(x) + \theta_{21} B_{21}(x) + \theta_{12} B_{12}(x) + \theta_{121} B_{121}(x). \end{aligned}$$

By Lemma 2.6 and Proposition 2.1, after applying π_1 on both sides, we get:

$$\begin{aligned}
& \pi_1(\pi_1(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l)) \\
&= \pi_1(\pi_{21}(x_1^k x_2^l) \times \pi_1(x_1^m x_2^n)) \\
&= \pi_{121}(x_1^k x_2^l) \times \pi_1(x_1^m x_2^n) + s_1 \pi_{21}(x_1^k x_2^l) \times \theta_1 \pi_1(x_1^m x_2^n) \\
&= \pi_{121}(x_1^k x_2^l) \times \pi_1(x_1^m x_2^n)
\end{aligned}$$

and

$$\begin{aligned}
& \pi_1(B_0(x) + \theta_1 B_1(x) + \theta_2 B_2(x) + \theta_{21} B_{21}(x) + \theta_{12} B_{12}(x) + \theta_{121} B_{121}(x)) \\
&= \pi_1 B_0(x) + \pi_1 \theta_2 B_2(x) + \pi_1 \theta_{21} B_{21}(x).
\end{aligned}$$

Therefore $\pi_1(\pi_1(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l)) = \pi_1 B_0(x) + \pi_1 \theta_2 B_2(x) + \pi_1 \theta_{21} B_{21}(x)$.

Now apply π_1 on both sides of

$$(x_1^m x_2^n) \times \pi_{121}(x_1^k x_2^l) = A_0(x) + \theta_1 A_1(x) + \theta_2 A_2(x) + \theta_{21} A_{21}(x) + \theta_{12} A_{12}(x) + \theta_{121} A_{121}(x),$$

we get

$$\begin{aligned}
& \pi_1((x_1^m x_2^n) \times \pi_{121}(x_1^k x_2^l)) \\
&= \pi_1(A_0(x) + \theta_1 A_1(x) + \theta_2 A_2(x) + \theta_{21} A_{21}(x) + \theta_{12} A_{12}(x) + \theta_{121} A_{121}(x)) \\
&= \pi_1 A_0(x) + \pi_1 \theta_2 A_2(x) + \pi_1 \theta_{21} A_{21}(x).
\end{aligned}$$

Since

$$\begin{aligned}
& \pi_1((x_1^m x_2^n) \times \pi_{121}(x_1^k x_2^l)) \\
&= \pi_1(x_1^m x_2^n) \times \pi_{121}(x_1^k x_2^l) + s_1(x_1^m x_2^n) \times \theta_1 \pi_{121}(x_1^k x_2^l) \\
&= \pi_1(x_1^m x_2^n) \times \pi_{121}(x_1^k x_2^l),
\end{aligned}$$

we can conclude that

$$\pi_1 A_0(x) + \pi_1 \theta_2 A_2(x) + \pi_1 \theta_{21} A_{21}(x) = \pi_1 B_0(x) + \pi_1 \theta_2 B_2(x) + \pi_1 \theta_{21} B_{21}(x).$$

Expand both sides as a sum of atoms:

$$\begin{aligned} & A_0(x) + \theta_1 A_0(x) + \theta_2 A_2(x) + \theta_1 \theta_2 A_2(x) + \theta_{21} A_{21}(x) + \theta_1 \theta_{21} A_{21}(x) \\ &= B_0(x) + \theta_1 B_0(x) + \theta_2 B_2(x) + \theta_1 \theta_2 B_2(x) + \theta_{21} B_{21}(x) + \theta_1 \theta_{21} B_{21}(x) \end{aligned}$$

and thus

$$\begin{aligned} & A_0(x) + \theta_1 A_0(x) + \theta_2 A_2(x) + \theta_{12} A_2(x) + \theta_{21} A_{21}(x) + \theta_{121} A_{21}(x) \\ &= B_0(x) + \theta_1 B_0(x) + \theta_2 B_2(x) + \theta_{12} B_2(x) + \theta_{21} B_{21}(x) + \theta_{121} B_{21}(x). \end{aligned}$$

Since the set of all atoms form a basis by item 3 in Theorem 2.8, we have $A_0 = B_0, A_2 = B_2$ and $A_{21} = B_{21}$.

4.2.3 $\pi_2(x_1^m x_2^n) \times \pi_{12}(x_1^k x_2^l)$

With the same notation in Section 4.2.2, we have

$$\begin{aligned} & \pi_2(x_1^m x_2^n) \times \pi_{12}(x_1^k x_2^l) \\ &= A_0(x) + \theta_1 A_1(x) + \sum_{r=0}^{\min\{m,k\}} \sum_{s=\max\{l,n,r,(n+l)-r\}}^{\min\{n+l,m+k-r\}} \theta_2 (x_1^{m+k+n+l-s-r} x_2^s x_3^r) + \theta_{12} A_{12}(x) \\ &+ \mathbb{1}_{\{n+l \geq \max\{m,k\}\}} \sum_{r=m+k-n-l}^{\min\{m,k\}} \theta_{21} (x_1^{n+l} x_2^{m+k-r} x_3^r) + \mathbb{1}_{\{n+l > k > m\}} \theta_{121} (x_1^{n+l} x_2^k x_3^m). \end{aligned}$$

4.2.4 $\pi_{12}(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l)$

With the same notation in Section 4.2.2, we have

$$\begin{aligned} & \pi_{12}(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l) \\ &= (1 + \theta_1 + \theta_2) A_0(x) + \theta_{21} A_1(x) + \theta_{12} A_2(x) \mathbb{1}_{\{m+l > k > n\}} + \sum_{t=0}^{\min\{m-n,l\}} \theta_{121} (x_1^{m+l-t} x_2^k x_3^{n+t}). \end{aligned}$$

4.2.5 All other cases in the table

We will complete the verification for all other unknown cases in the tables by applying Lemma 3.16 and Lemma 2.6 on verified cases.

(i) As

$$\begin{aligned}
& \pi_1(x_1^m x_2^n) \times \pi_1(x_1^k x_2^l) \\
&= \pi_1(x_1^m x_2^n \times \pi_1(x_1^k x_2^l)) \\
&= \pi_1(x_1^m x_2^n \times x_1^k x_2^l + x_1^m x_2^n \times \theta_1(x_1^k x_2^l)),
\end{aligned}$$

result follows by Theorem 3.13.

(ii)

$$\begin{aligned}
& \pi_1(x_1^m x_2^n) \times \pi_{12}(x_1^k x_2^l) \\
&= \pi_1(x_1^m x_2^n \times \pi_{12}(x_1^k x_2^l)) \\
&= \pi_1\left(x_1^m x_2^n \times (x_1^k x_2^l + \theta_1(x_1^k x_2^l) + \theta_2(x_1^k x_2^l) + \theta_{12}(x_1^k x_2^l))\right),
\end{aligned}$$

result follows by Theorem 3.13.

Alternatively, one can use the fact that

$\pi_1(x_1^m x_2^n) \times \pi_{12}(x_1^k x_2^l) = \pi_1(\pi_1(x_1^m x_2^n) \times \pi_2(x_1^k x_2^l)) = \pi_1(\pi_2(x_1^k x_2^l) \times \pi_1(x_1^m x_2^n))$ by putting $f = \pi_2(x_1^k x_2^l)$, $g = \pi_1(x_1^m x_2^n)$ and $i = 1$ in Lemma 2.6 and claim $\pi_1(x_1^m x_2^n) \times \pi_{12}(x_1^k x_2^l)$ is atom positive by applying Lemma 3.16 on

$\pi_1(x_1^m x_2^n) \times \pi_2(x_1^k x_2^l)$ which is verified as atom positive in Section 4.2.1.

(iii)

$$\begin{aligned}
& \pi_2(x_1^m x_2^n) \times \pi_2(x_1^k x_2^l) \\
&= \pi_2(x_1^m x_2^n \times \pi_2(x_1^k x_2^l)) \\
&= \pi_2(x_1^m x_2^n \times x_1^k x_2^l + x_1^m x_2^n \times \theta_2(x_1^k x_2^l)),
\end{aligned}$$

result follows by Theorem 3.13.

(iv) As

$$\begin{aligned}
& \pi_2(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l) \\
&= \pi_2(x_1^m x_2^n \times \pi_{21}(x_1^k x_2^l)) \\
&= \pi_2(x_1^m x_2^n \times x_1^k x_2^l + x_1^m x_2^n \times \theta_1(x_1^k x_2^l) + x_1^m x_2^n \times \theta_2(x_1^k x_2^l) + x_1^m x_2^n \times \theta_{21}(x_1^k x_2^l))
\end{aligned}$$

and result follows by Theorem 3.13.

Similar to Case (ii), one can also use the fact that

$$\pi_2(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l) = \pi_2(\pi_2(x_1^m x_2^n) \times \pi_1(x_1^k x_2^l)) = \pi_2(\pi_1(x_1^k x_2^l) \times \pi_2(x_1^m x_2^n))$$

and claim $\pi_1(x_1^m x_2^n) \times \pi_{12}(x_1^k x_2^l)$ is atom positive.

(v) $\pi_{21}(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l) = \pi_2(\pi_1(x_1^m x_2^n) \times \pi_{21}(x_1^k x_2^l))$ and result follows by the case in Section 4.2.2.

(vi) $\pi_{12}(x_1^m x_2^n) \times \pi_{12}(x_1^k x_2^l) = \pi_1(\pi_2(x_1^m x_2^n) \times \pi_{12}(x_1^k x_2^l))$ and result follows by the case in Section 4.2.3.

Appendix A

Bijection between LRS and LRK

We will illustrate the map ϕ in Section 6 in [3] without proof. The definition and examples are mostly adapted from [3].

Definition A.1. A word $w = w_1w_2w_3\dots$ is called *contre-lattice* if for any initial sequence $w_1\dots w_i$, there are at least as many occurrences of the number k as the number of $k-1$ for each $1 < k \leq \max\{w_m : 1 \leq m \leq i\}$. We call w a *regular contre-lattice word* if w is *contre-lattice* and contains the number 1.


Example 19. 3231321 is a regular *contre-lattice word* while 3132321 is not.

Definition A.2. Let δ, γ be weak compositions. A *Littlewood-Richardson skew skyline tableau* (LRS) of shape δ/γ is an *SSAF* of shape δ/γ with basement entry of the i^{th} -column is $b_i = 2n + 1 - i$, where $n = l(\delta) = l(\gamma)$, whose reading word (obtained by reading the entries in the cells in ascending reading order) is a regular *contre-lattice word*. We use $\text{LRS}(n)$ to denote the set of LRS with entries in $[n]$.

Definition A.3. Let δ, γ be weak compositions. A *Littlewood-Richardson skew key* (LRK) of shape δ/γ is an *SSAF* of shape δ/γ with basement entry of the i^{th} -column is $b_i = n + i$, where $n = l(\delta) =$

$l(\gamma)$, whose reading word (obtained by reading the entries in the cells in ascending reading order) is a regular contre-lattice word. We use $\text{LRK}(n)$ to denote the set of LRK with entries in $[n]$.

Example 20. An LRS (left) and an LRK (left) with reading word 3231321.



We now describe the map ϕ in Section 6 in [3].

Given an LRK K of shape δ/γ with basement entry of the i^{th} -column is $b_i = n + i$ where $n = l(\delta) = l(\gamma)$, and fix any permutation of δ . Then we can find a unique LRS of overall shape σ whose set of entries of each row is the same as that of the given LRK. We can find this LRS by successively filling the rightmost column strip in the unfilled portion of the diagram for the set of rows containing the smallest entry of K at each step. Here a column strip means a sequence of cells chosen in such a way that they appear in the topmost portion of the diagram (i.e. if a cell is chosen, wither it is the top cell of that column or all cells above it are chosen).

Conversely, given a LRS L of shape δ/β and fix a rearrangement of β , say, γ such that $\bar{\gamma} \leq \beta$, then we can find the corresponding LRK with basement shape $\bar{\gamma}$ as follows:

Consider the basement diagram K^0 with basement shape $\bar{\gamma}$ on which we will build the desired LRK. Consider the bottom row of L and start from the largest entry. Place this entry to the leftmost available cell in the lowest row of K^0 and call the resulting filling as K^1 . Then place the second largest entry of the bottom row of L to the leftmost available cell in the lowest row of K^1 , and so on until all entries of the bottom row of L is filled into the bottom row of the basement diagram. Repeat this process with each column from the bottom to the top until all non-basement entries of L have been placed into the diagram.


We illustrate this process by using the LRK in Example 20.

Example 21. Consider the LRK

3				2
3		1		10
3		2	9	10
6	1	8	9	10
6	7	8	9	10

and fix $\sigma = (2, 4, 1, 4, 3)$. Then we can get a LRS by filling the entries as


shown below:




Conversely, if we start from

3	2			
10	3		1	
10	9		3	2
10	9	1	7	6
10	9	8	7	6

and fix $\gamma = (3, 2, 1, 0, 1)$, we can build the LRS as follows:





and we get back the LRK we started from.

Bibliography

- [1] DEODHAR, V. V. Some characterizations of Bruhat ordering on a Coxeter group and determination of the relative Möbius function. *Invent. Math.* 39, 2 (1977), 187–198.
- [2] HAGLUND, J., HAIMAN, M., AND LOEHR, N. A combinatorial formula for nonsymmetric Macdonald polynomials. *Amer. J. Math.* 130, 2 (2008), 359–383.
- [3] HAGLUND, J., LUOTO, K., MASON, S., AND VAN WILLIGENBURG, S. Refinements of the Littlewood-Richardson rule. *Trans. Amer. Math. Soc.* 363, 3 (2011), 1665–1686.
- [4] HAGLUND, J., MASON, S., AND REMMEL, J. Properties of the nonsymmetric Robinson-Schensted-Knuth algorithm. *J. Algebraic Combin.* 38, 2 (2013), 285–327.
- [5] JOSEPH, A. A decomposition theorem for Demazure crystals. *Journal of Algebra* 265, 2 (2003), 562 – 578.
- [6] LASCoux, A., AND SCHÜTZENBERGER, M.-P. Keys & standard bases. In *Invariant theory and tableaux (Minneapolis, MN, 1988)*, vol. 19 of *IMA Vol. Math. Appl.* Springer, New York, 1990, pp. 125–144.
- [7] LASKU, A., AND SHYUTTSSENBERZHE, M.-P. Noncommutative Schubert polynomials. *Funktional. Anal. i Prilozhen.* 23, 3 (1989), 63–64.

- [8] MACDONALD, I. G. *Symmetric functions and Hall polynomials*, second ed. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [9] MACDONALD, I. G. Affine Hecke algebras and orthogonal polynomials. *Astérisque*, 237 (1996), Exp. No. 797, 4, 189–207. Séminaire Bourbaki, Vol. 1994/95.
- [10] MARSHALL, D. Symmetric and nonsymmetric Macdonald polynomials. *Ann. Comb.* 3, 2-4 (1999), 385–415. On combinatorics and statistical mechanics.
- [11] MASON, S. A decomposition of Schur functions and an analogue of the Robinson-Schensted-Knuth algorithm. *Sém. Lothar. Combin.* 57 (2006/08), Art. B57e, 24.
- [12] MASON, S. An explicit construction of type A Demazure atoms. *J. Algebraic Combin.* 29, 3 (2009), 295–313.
- [13] WATANABE, M. Tensor product of Kraskiewicz and Pragacz’s modules. *ArXiv e-prints* (Oct. 2014).