



5-18-2011

# Neutron Widths and Configuration Mixing in $^{11}\text{Be}$

H. Terry Fortune

*University of Pennsylvania*, [fortune@physics.upenn.edu](mailto:fortune@physics.upenn.edu)

Rubby Sherr

*Princeton University*

Follow this and additional works at: [http://repository.upenn.edu/physics\\_papers](http://repository.upenn.edu/physics_papers)

 Part of the [Physics Commons](#)

---

## Recommended Citation

Fortune, H. T., & Sherr, R. (2011). Neutron Widths and Configuration Mixing in  $^{11}\text{Be}$ . Retrieved from [http://repository.upenn.edu/physics\\_papers/175](http://repository.upenn.edu/physics_papers/175)

### Suggested Citation:

H.T. Fortune and R. Sherr. (2011). Neutron widths and configuration mixing in  $^{11}\text{Be}$ . *Physical Review C*, 054314.

© 2011 The American Physical Society

<http://dx.doi.org/10.1103/PhysRevC.83.054314>

This paper is posted at Scholarly Commons. [http://repository.upenn.edu/physics\\_papers/175](http://repository.upenn.edu/physics_papers/175)

For more information, please contact [repository@pobox.upenn.edu](mailto:repository@pobox.upenn.edu).

---

# Neutron Widths and Configuration Mixing in $^{11}\text{Be}$

## Abstract

We use known widths and branching ratios in  $^{11}\text{Be}$  to discuss  $J^\pi$  and configuration admixtures. Analysis favors  $3/2^-$  for the 3.96-MeV state and three-state mixing for this  $J^\pi$ .

## Disciplines

Physical Sciences and Mathematics | Physics

## Comments

Suggested Citation:

H.T. Fortune and R. Sherr. (2011). Neutron widths and configuration mixing in  $^{11}\text{Be}$ . *Physical Review C*, 054314.

© 2011 The American Physical Society

<http://dx.doi.org/10.1103/PhysRevC.83.054314>

## Neutron widths and configuration mixing in $^{11}\text{Be}$

H. T. Fortune

*Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*

R. Sherr

*Department of Physics, Princeton University, Princeton, New Jersey 08544, USA*

(Received 3 March 2011; published 18 May 2011)

We use known widths and branching ratios in  $^{11}\text{Be}$  to discuss  $J^\pi$  and configuration admixtures. Analysis favors  $3/2^-$  for the 3.96-MeV state and three-state mixing for this  $J^\pi$ .

DOI: [10.1103/PhysRevC.83.054314](https://doi.org/10.1103/PhysRevC.83.054314)

PACS number(s): 21.10.Jx, 21.10.Tg, 27.20.+n

### I. INTRODUCTION

A variety of experiments have been used to establish the properties of the low-lying states of  $^{11}\text{Be}$ . Most of the results are from one- and two-nucleon transfer experiments and from  $\beta$  decay. These include  $^{10}\text{Be}(d,p)$ ,  $^9\text{Be}(t,p)$ ,  $^9\text{Be}(^{16}\text{O},^{14}\text{O})$ ,  $\beta$  decay of  $^{11}\text{Li}$ , and neutron decays following the last two. Early information concerning levels of  $^{11}\text{Be}$  is summarized in a series of compilations [1–3]. The reaction  $^{10}\text{Be}(d,p)$  has been studied at bombarding energies of 12 MeV [4] and 25 MeV [5]. The  $^9\text{Be}(t,p)$  reaction has been performed at energies of 5 MeV [6], 15 MeV [7], 20 MeV [8], and 23 MeV [9]. Early  $\beta$  decay of  $^{11}\text{Li}$  was discussed in Ref. [10]. Much of the  $J^\pi$  information in Refs. [1–3] is incorrect and has been superseded [7,11–13]. There is now general agreement on most values of  $J^\pi$  and dominant configurations of states up to 5.5 MeV. Data for levels at excitation energies below 6 MeV are summarized in Table I, where the  $J^\pi$  values are the most recent ones, most now considered firm [13]. The  $3/2^+$  suggestion for the 3.89-MeV state in Ref. [12] was based on incomplete information and  $J^\pi$  has since been determined to be  $5/2^-$  from  $\beta$  decay. Some information is available from the reaction  $^{13}\text{C}(^6\text{Li},^8\text{B})$  [14]. Early theoretical work was done by Cohen-Kurath (CK) [15] for the  $p$ -shell states and by Teeters-Kurath [16] for positive parity. Other calculations involve an SU(3) cluster model [12]. Additional experimental information is available in Refs. [17–22]. In the present paper, we discuss quantitatively the quality of agreement between the known properties and the supposed structure.

### II. WIDTHS AND SPECTROSCOPIC FACTORS

Unless indicated otherwise, the widths we use (Table II) are the weighted average of those from the compilation and from results of the  $^9\text{Be}(t,p)$  reaction [7]. For the  $5/2^+$  state at 1.778 MeV, the majority of its configuration is a  $d_{5/2}$  single particle (sp) coupled to the ground state (gs) of  $^{10}\text{Be}$ . Presumably most of the remaining strength is  $^{10}\text{Be}(2^+) \times 2s_{1/2}$ . The width of 102(14) keV, combined with the  $\ell = 2$  sp width of 175 keV, gives a spectroscopic factor of  $S = \Gamma/\Gamma_{\text{sp}} = 0.58(8)$ . Values from  $(d,p)$  are similar:  $S = 0.50$  in Ref. [5].

The state at 2.69 MeV has been identified as having  $J^\pi = 3/2^-$ , and is predominantly the lowest  $p$ -shell  $3/2^-$  state. The cross section in  $(t,p)$  is consistent with this interpretation.

(However, see below.) Its total width is 213(15) keV. The  $p$ -wave sp width for this energy is  $\sim 1.8$  MeV, yielding  $S \sim 0.12$ , to be compared with the expected value of 0.106 [12] or 0.168 [15].

The state at 3.41 MeV was assigned  $J^\pi = 3/2^-$  from an apparent  $L = 0$  angular distribution in  $(t,p)$ , and suggested [7] to be the lowest  $(sd)_{0+}^2$  state:  $^9\text{Be}(\text{gs}) \times (sd)_{0+}^2$ . However, the absolute  $(t,p)$  cross section is only  $\sim 28\%$  of the value expected for this configuration. Morrissey *et al.* [11], based on its nonobservation in  $\beta$  decay, suggested it might be a  $3/2^+$  state. Its width is 113(13) keV. An  $\ell = 1$  sp width is difficult to calculate at this energy, but is  $\sim 2.1$  MeV, implying  $S \sim 0.05$  if  $3/2^-$ . The  $\ell = 2$  sp width is 1.1 MeV, giving  $S = 0.10(1)$  if  $3/2^+$ —close to the upper limit for the amount of  $d_{3/2}$  strength expected this low in excitation energy. Of course, a  $3/2^-$  state with configuration  $^9\text{Be}(\text{gs}) \times (sd)_{0+}^2$  would have no gs width either, except through mixing in the initial or final state. If we add this  $S$  to the one for the 2.69-MeV  $3/2^-$  state, we get  $S_{\text{tot}} = 0.17$ , close to the CK value of 0.168 for the  $p$ -shell  $3/2^-$ . Another possibility is mixing in the  $^{10}\text{Be}(\text{gs})$ . A small component  $\varepsilon$  of  $^8\text{Be}(\text{gs}) \times (sd)_{0+}^2$  in  $^{10}\text{Be}(\text{gs})$  would produce a gs decay spectroscopic factor of  $\varepsilon^2 S[^9\text{Be}(\text{gs}) \rightarrow ^8\text{Be}(\text{gs}) + n]$ . In CK, the second factor is 0.58, requiring  $\sim 10\%$  mixing in  $^{10}\text{Be}(\text{gs})$  if this is the sole explanation. Undoubtedly, several effects combine to produce the observed strength. For the 2.69-MeV state, the  $L = 0 + 2$  distorted wave Born approximation (DWBA) curve displayed with the data has the  $L = 0, 2$  mixture appropriate for the  $p$ -shell  $3/2^-$  state. The excess of experimental cross section over the DWBA curve at small angles indicates that the  $L = 0$  component is stronger than expected for the  $p$ -shell state. The  $(sd)_{0+}^2 3/2^-$  state should have pure  $L = 0$ , and the 3.41-MeV state has only  $\sim 30\%$  of the expected cross section. If the  $p$ -shell and  $(sd)_{0+}^2 3/2^-$  states mix, the lower one will acquire a constructive sum of amplitudes and the higher state a destructive sum. So, mixing of these two states could explain both the  $(t,p)$  results and the decay widths. Also, as  $^{11}\text{Li}$  undoubtedly has a constructive sum of  $(sd)_{0+}^2$  and  $p$ -shell neutrons,  $\beta$  decay to the 3.41-MeV state would involve a small overlap, producing a small branch. Thus, this mixing could also explain the  $\beta$  decay. The  $3/2^+$  possibility is discussed further in Sec. III.

Beginning with the state at 3.89 MeV, states of  $^{11}\text{Be}$  can decay to the  $2^+$  first-excited state of  $^{10}\text{Be}$  at 3.368 MeV. For 3.89 MeV, the neutron decay energy is very small—only

TABLE I. Properties of low-lying states of  $^{11}\text{Be}$ .

Compilation <sup>a</sup>		$(t,p)$ <sup>b</sup>		$J^\pi$ <sup>c</sup>	Dominant configuration <sup>b</sup>
$E_x$ (MeV)	$\Gamma$ (keV)	$E_x$ (MeV)	$\Gamma$ (keV)		
0	–	–0.004(3)	–	$1/2^+$	$^{10}\text{Be}(\text{gs}) \times 2s_{1/2}$
0.32004(10)	–	0.320(2)	–	$1/2^-$	$p$ shell
1.778(12)	100(20)	1.748(4)	104(21)	$5/2^+$	$^{10}\text{Be}(\text{gs}) \times 1d_{5/2}$
2.69(2)	200(20)	2.642(9)	228(21)	$3/2^-$	$p$ shell
3.41(2)	125(20)	3.398(6)	104(17)	$3/2^-$ $3/2^+$	$^9\text{Be} \times (\text{sd})_{0+}^2$ $^{10}\text{Be}(2^+) \times 2s_{1/2}$
3.887(12)	<10	3.888(1)	–	$5/2^-$	$p$ shell
3.956(15)	15(5)	3.955(1)	–	$3/2^-$	$^9\text{Be} \times (\text{sd})_{2+}^2$
5.240(21)	45(10)	5.255(3)	29(8) <sup>d</sup>	$5/2^-$	$^9\text{Be} \times (\text{sd})_{2+}^2$
(5.86)	$\sim 300$	5.849(10)	139(17)	$(5/2^-)?$	?

<sup>a</sup>Reference [3].<sup>b</sup>Reference [7].<sup>c</sup>References [7,11–13].<sup>d</sup>Subsequent analysis of data of Ref. [7].

19 keV—and yet a  $2^+$  decay is observed. Following  $\beta$  decay of  $^{11}\text{Li}$ , a branching ratio (BR) of  $\Gamma(2^+)/\Gamma_{\text{tot}} = 0.62^{+0.14}_{-0.21}$  is reported [13]. (See Table III.) As this state is thought to be  $5/2^-$ , this  $2^+$  decay is understandable, because gs decay is hindered by a large centrifugal barrier and by a small amount of expected  $f_{5/2}$  sp strength. For  $\ell = 1$  and  $E_n = 19$  keV, we have  $\Gamma_{\text{sp}} = 2.9$  keV. In CK, the spectroscopic factor for the first  $5/2^-$  state to decay to  $2^+$  is 0.66. (Millener has 0.574.) With the CK value and our sp width, we have  $\Gamma(2^+) = 1.9$  keV as the width expected for decay to  $2^+$ . The BR would then give  $\Gamma_{\text{tot}} = 3.1^{+1.5}_{-0.6}$  keV, leaving 1.2(7) keV for decay to the gs. This total width is consistent with the one given in the compilation as <10 keV. An  $\ell = 3$  sp width for gs decay at this energy is 170 keV—implying  $S(f_{5/2}) = 0.007(4)$ , quite an acceptably small value.

We come now to the 3.96-MeV  $3/2^-$  state. Following  $\beta$  decay its BR for  $2^+$  decay is given [13] as  $\Gamma(2^+)/\Gamma_{\text{tot}} = 0.78^{+0.65}_{-0.31}$ . In the  $^9\text{Be}(^{16}\text{O},^{14}\text{O})$  reaction [20], this ratio is 0.54(7), though this state is not resolved from the 3.89-MeV state. In  $^9\text{Be}(t,p)$  the 3.89-MeV state is approximately half as strong (angle-integrated cross sections) as 3.96 MeV. A similar ratio might be expected in the heavy-ion-induced  $2n$  transfer. Reference [20] points out that no amount of combined 3.89 + 3.96 yield will make their data consistent with those of Ref. [13]. The total width [3] of this state is 15(5) keV. For gs decay, the sp width is difficult to calculate, but we

estimate it to be  $\sim 2.3$  MeV. The gs branch for the 3.96-MeV state is 0.22(4) in Ref. [13] and 0.48(6) in Ref. [20], leading to  $S(\text{gs}) = 1.4(5) \times 10^{-3}$  or  $3.1(11) \times 10^{-3}$ . These are small enough to arise from small, neglected components in the wave function. For decay to the  $2^+$  state, the decay energy is 98 keV, for which  $\Gamma_{\text{sp}}$  is 36 keV. Thus, if we use the  $\beta$  decay BR, we have  $S = 0.32(17)$  for  $2^+$  decay, quite a large value (though with a large uncertainty) for a state thought to be dominated by the configuration  $^9\text{Be}(\text{gs}) \times (\text{sd})_{2+}^2$ . The BR from Ref. [20] gives  $S = 0.22(8)$ , still quite large. Millener has  $S = 0.864$  for the  $p$ -shell  $3/2^-$  to decay to  $2^+$ , but this is presumably the 2.69-MeV state. The next  $p$ -shell  $3/2^-$  has  $S = 0.123$ , but it is expected above 5 MeV. It is very likely that mixing occurs among the lowest three  $3/2^-$  states:  $p$ -shell,  $^9\text{Be} \times (\text{sd})_0^2$ , and  $^9\text{Be} \times (\text{sd})_2^2$ . In  $(t,p)$  there is a hint of an  $L = 0$  contribution to the 3.96-MeV angular distribution (as can be seen by comparing data for 3.96 and 5.24 MeV—the latter is pure  $L = 2$ ). Also, 3.96 is slightly ( $\sim 15\%$ ) weaker than it should be, in comparison with 5.24, if they are both  $^9\text{Be} \times (\text{sd})_{2+}^2$ . The situation is discussed further in Sec. III. Clearly, we need smaller uncertainties here—both in  $\Gamma_{\text{tot}}$  and BR—and a better understanding of the discrepancy between results of Refs. [13] and [20].

The next state is at 5.24 MeV, and has been assigned  $J^\pi = 5/2^-$  and suggested to have the configuration  $^9\text{Be} \times (\text{sd})_{2+}^2$ . Its width is given in the compilation [3] as 45(10) keV.

TABLE II. Widths (keV) and spectroscopic factors for three lowest unbound states of  $^{11}\text{Be}$ .

$E_x$ (MeV)	$J^\pi$	$\Gamma_{\text{exp}}^a$	$\ell$	$\Gamma_{\text{sp}}$	$S = \Gamma_{\text{exp}}/\Gamma_{\text{sp}}$	$S_{\text{th}}$
1.78	$5/2^+$	102(14)	2	175	0.58(8)	$0.67^b$
2.69	$3/2^-$	213(15)	1	$\sim 1800$	$\sim 0.12$	$0.168^c$
3.41	$3/2^-$	113(13)	1	$\sim 2100$	$\sim 0.05$	–
	$3/2^+$		2	1100	0.10(1)	–

<sup>a</sup>Weighted average of values in Table I.<sup>b</sup>Reference [12].<sup>c</sup>Reference [15].TABLE III. Branching ratios of states in  $^{11}\text{Be}$  for neutron decays to  $^{10}\text{Be}$ .

$E_x$ (MeV)	$J^\pi$	$0^+$	$2^+$	Ref.
3.890	$5/2^-$	0.38(9)	$0.62^{+0.14}_{-0.21}$	[13]
3.969	$3/2^-$	0.22(4)	$0.78^{+0.65}_{-0.31}$	[13]
		0.48(6) <sup>a</sup>	$0.54(7)^a$	[20]
5.24	$5/2^-$	–	1.00	[13]
		<0.27	0.81(16)	[20]
5.96	$(5/2^-)?$	<0.13	0.97(16)	[20]

<sup>a</sup>Contains contributions from both 3.89 and 3.96.

No width is listed for this state in the  $(t,p)$  paper, but subsequent analysis of those data yields  $\Gamma = 29(8)$  keV, giving an average of  $35(6)$  keV. The only observed  $n$  decays are to the  $2^+$ . With an energy of 1.38 MeV, the  $\ell = 1$  sp width is 1.45 MeV, giving  $S = 0.024(4)$ . This  $S$  is small enough that it could easily be acquired by mixing with the  $p$ -shell  $5/2^-$ —which is predicted to have  $S = 0.66$ . The limit of  $\text{BR} < 0.27$  [20] for the gs branch corresponds to  $S(f_{5/2}) < 0.02$ .

The state at 5.96 MeV may or may not be the same as the 5.849(10)-MeV state seen in  $(t,p)$ , with a width of 139(17) keV. The compilation lists a questionable state at (5.86) MeV with a width of  $\sim 300$  keV. The 5.96-MeV state in Ref. [20] has  $\Gamma = 400$  keV and only a  $2^+$  branch.

### III. CONFIGURATION MIXING

If the 3.41-MeV state has  $J^\pi = 3/2^+$ , then the lowest  $(sd)^2$  state is the one at 3.96 MeV. The strength of the latter in  $(t,p)$  indicates very little mixing ( $\sim 1\%$ – $2\%$ ) between it and the first  $3/2^-$   $p$ -shell state. Its small gs spectroscopic factor (Sec. II) is also consistent with very little mixing. The weakness of its  $L = 0$  component and the strong decay of the 3.96-MeV state to the  $2^+$  of  $^{10}\text{Be}$  may be more difficult to understand. If  $J^\pi$  (3.41) is  $3/2^+$ , its  $d_{3/2}$  spectroscopic factor of  $\sim 0.10$  (see above) is large enough that it should exhibit a clear  $\ell = 2$  stripping pattern in  $^{10}\text{Be}(d,p)$ . Reference [5] states that they investigated the range of excitation energy up to  $E_x = 7.0$  MeV with the  $^{10}\text{Be}(d,p)$  reaction, but their published spectrum goes only to just above 3 MeV. They state that they did not observe the 3.41-MeV state. However, they also state that the 2.69-MeV state was not excited with measurable strength. The present analysis gives a value of  $S \sim 0.12$  for the latter.

We now consider the possibility that  $J^\pi$  (3.41) is  $3/2^-$ , and summarize the arguments for three-state mixing among the

$3/2^-$  states: The 2.69-MeV state has excess  $L = 0$  strength over that expected for the  $p$ -shell  $3/2^-$ . The 3.41-MeV state has only 28% of the cross section expected for the lowest  $^9\text{Be} \times (sd)_0^2$  state, and it has appreciable width for decay to the gs. The 3.96-MeV state has 15% less strength than that expected for the  $^9\text{Be} \times (sd)_2^2$  configuration, and its angular distribution contains a hint of  $L = 0$  (not present for that configuration). Also it has an appreciable  $S$  for decay to the  $2^+$  state (although with a large uncertainty). Recall that the  $p$ -shell  $3/2^-$  state has a large  $S(2^+)$ . If only the lowest two states had mixed, 12% of the  $p$ -shell  $3/2^-$  state mixed into the 3.41-MeV level is enough to explain the  $(t,p)$  results, while the decay widths suggest 28%, with some uncertainty. For the 3.96-MeV state, the  $2^+$  decay suggests  $\sim 25\%$  mixing of the  $p$ -shell state, whereas only  $\sim 15\%$  of the  $(t,p)$  strength is lacking. It might appear that the absence of any appreciable  $L = 2$  strength in the 3.41-MeV angular distribution might argue against all this. But, this fact actually argues against two-state mixing and for all three states to mix. If the second  $3/2^-$  state mixed with only one of the others, it would necessarily acquire some  $L = 2$  strength. But, with three states, the  $L = 2$  admixtures from the other two states could be destructive for the 3.41-MeV state.

For the  $5/2^-$  states: The 3.89-MeV state is predicted (and observed) to be weak, but it is stronger than expected. As its angular distribution is not a clear  $L = 2$  shape, it could contain some nondirect (e.g., compound nucleus) contribution. If it receives its extra strength from the much stronger 5.24-MeV state, the mixing would have a negligible effect on the latter.

### ACKNOWLEDGMENTS

We acknowledge stimulating correspondence with John Millener.

- 
- [1] F. Ajzenberg-Selove and C. L. Busch, *Nucl. Phys. A* **336**, 1 (1980).  
 [2] F. Ajzenberg-Selove, *Nucl. Phys. A* **433**, 1 (1985).  
 [3] F. Ajzenberg-Selove, *Nucl. Phys. A* **506**, 1 (1990).  
 [4] D. L. Auton, *Nucl. Phys. A* **157**, 305 (1970).  
 [5] B. Zwiaglinski, W. Benenson, R. G. H. Robertson, and W. R. Coker, *Nucl. Phys. A* **315**, 124 (1979).  
 [6] D. J. Pullen, E. R. Litherland, S. Hinds, and R. Middleton, *Nucl. Phys.* **36**, 1 (1962).  
 [7] G.-B. Liu and H. T. Fortune, *Phys. Rev. C* **42**, 167 (1990).  
 [8] F. Ajzenberg-Selove, R. F. Casten, O. Hansen, and T. J. Mulligan, *Phys. Lett. B* **40**, 205 (1972).  
 [9] F. Ajzenberg-Selove, E. R. Flynn, and O. Hansen, *Phys. Rev. C* **17**, 1283 (1978).  
 [10] R. E. Azuma *et al.*, *Phys. Rev. Lett.* **43**, 1652 (1979).  
 [11] D. J. Morrissey *et al.*, *Nucl. Phys. A* **627**, 222 (1997).  
 [12] D. J. Millener, *Nucl. Phys. A* **693**, 394 (2001).  
 [13] Y. Hirayama *et al.*, *Phys. Lett. B* **611**, 239 (2005).  
 [14] R. B. Weisenmiller *et al.*, *Phys. Rev. C* **13**, 1330 (1976).  
 [15] S. Cohen and D. Kurath, *Nucl. Phys. A* **101**, 1 (1967); **141**, 145 (1970).  
 [16] W. D. Teeters and D. Kurath, *Nucl. Phys. A* **275**, 61 (1975); **283**, 1 (1977).  
 [17] I. Mukha *et al.*, *Phys. Lett. B* **367**, 65 (1996).  
 [18] M. J. G. Borge *et al.*, *Phys. Rev. C* **55**, R8 (1997).  
 [19] N. Aoi *et al.*, *Nucl. Phys. A* **616**, 181c (1997); **358**, 253 (1997).  
 [20] P. J. Haigh *et al.*, *Phys. Rev. C* **79**, 014302 (2009).  
 [21] F. Sarazin *et al.*, *Phys. Rev. C* **70**, 031302(R) (2004).  
 [22] H. O. U. Fynbo *et al.*, *Nucl. Phys. A* **736**, 39 (2004).