

BioCell Your Trusted Supplier of *in vivo* MAbs
 α -PD-1 · α -PD-L1 · α -CTLA-4 · α -CD20 · α -NK1.1 · α -IFNAR-1

DISCOVER MORE



C3a Enhances Nerve Growth Factor-Induced NFAT Activation and Chemokine Production in a Human Mast Cell Line, HMC-1

This information is current as of September 27, 2022.

Jasimuddin Ahamed, Rampura T. Venkatesha, E. Berla Thangam and Hydar Ali

J Immunol 2004; 172:6961-6968; ;
doi: 10.4049/jimmunol.172.11.6961
<http://www.jimmunol.org/content/172/11/6961>

References This article cites **77 articles**, 39 of which you can access for free at:
<http://www.jimmunol.org/content/172/11/6961.full#ref-list-1>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

*average

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>

The Journal of Immunology is published twice each month by
The American Association of Immunologists, Inc.,
1451 Rockville Pike, Suite 650, Rockville, MD 20852
Copyright © 2004 by The American Association of
Immunologists All rights reserved.
Print ISSN: 0022-1767 Online ISSN: 1550-6606.



C3a Enhances Nerve Growth Factor-Induced NFAT Activation and Chemokine Production in a Human Mast Cell Line, HMC-1¹

Jasimuddin Ahamed,² Rampura T. Venkatesha, E. Berla Thangam, and Hydar Ali³

Activation of cell surface G protein-coupled receptors leads to transphosphorylation and activation of a number of receptor tyrosine kinases. Human mast cells express G protein-coupled receptors for the complement component C3a (C3aR) and high affinity nerve growth factor (NGF) receptor tyrosine kinase, TrkA. To determine whether C3a cross-regulates TrkA signaling and biological responses, we used a human mast cell-line, HMC-1, that natively expresses both receptors. We found that NGF caused tyrosine phosphorylation of TrkA, resulting in a sustained Ca^{2+} mobilization, NFAT activation, extracellular-signal regulated kinase (ERK) phosphorylation, and chemokine, macrophage inflammatory protein-1 β (MIP-1 β) production. In contrast, C3a induced a transient Ca^{2+} mobilization and ERK phosphorylation but failed to stimulate TrkA phosphorylation, NFAT activation, or MIP-1 β production. Surprisingly, C3a significantly enhanced NGF-induced NFAT activation, ERK phosphorylation, and MIP-1 β production. Pertussis toxin, a $G_{i/o}$ inhibitor, selectively blocked priming by C3a but had no effect on NGF-induced responses. Mitogen-activated protein/ERK kinase inhibitor U0126 caused ~30% inhibition of NGF-induced MIP-1 β production but had no effect on priming by C3a. However, cyclosporin A, an inhibitor of calcineurin-mediated NFAT activation, caused substantial inhibition of NGF-induced MIP-1 β production both in the absence and presence of C3a. These data demonstrate that NGF caused tyrosine phosphorylation of TrkA to induce chemokine production in HMC-1 cells via a pathway that mainly depends on sustained Ca^{2+} mobilization and NFAT activation. Furthermore, C3a enhances NGF-induced transcription factor activation and chemokine production via a G protein-mediated pathway that does not involve TrkA phosphorylation. *The Journal of Immunology*, 2004, 172: 6961–6968.

Nerve growth factor (NGF)⁴ was originally described as a neurotrophic factor that plays a critical role in the development and maintenance of sensory and sympathetic neurons (1–3). There is now a substantial body of evidence that demonstrates an important role of NGF in allergic and inflammatory diseases. For example, activation of mast cells via the cross-linking of cell surface Fc ϵ RI results in the synthesis and release of NGF (4). Furthermore, increased levels of NGF have been demonstrated in plasma, and nasal and bronchoalveolar lavage fluids obtained from allergic patients (5–7). Several lines of evidence suggest that NGF released from mast cells mediates autocrine/paracrine effects on mast cell development and activation. NGF causes mast cell hypertrophy and hyperplasia in many tissues and organs (8). NGF is a mast cell chemoattractant and it also induces histamine release from these cells (9–12). NGF has recently been shown to stimulate the expression of IL-6 in bone marrow-derived mast cells (13). Mast cells

are a rich source of chemokines, and these proinflammatory cytokines play important roles in allergic and inflammatory diseases (14–18). However, whether NGF stimulates chemokine production in mast cells has not been determined.

The biological effects of NGF are mediated via two types of cell surface receptors, p75 and TrkA (19). Mast cells express TrkA but not p75 (11, 20, 21). TrkA-mediated signaling pathways have been studied most extensively in neuronal cell lines. In this system, NGF causes tyrosine phosphorylation of TrkA leading to the activation of two main signaling pathways. Phosphorylation of TrkA at Tyr⁴⁹⁰ leads to the binding of Shc, resulting in transient activation of the Ras/mitogen-activated protein (MAP) kinase signaling cascade (22, 23). In contrast, phosphorylation of TrkA at the other major phosphorylation site, Tyr⁷⁸⁵, mediates docking and sustained activation of phospholipase C γ (PLC γ) (23, 24). Whereas the transient Ras/MAP kinase pathway mediates neuronal cell survival (25, 26), the sustained PLC γ activation leads to the expression of Na⁺ channel gene (27). NFAT are a family of transcription factors expressed in a wide variety of cell types (28–31). Ag stimulation of T cells and mast cells leads to tyrosine phosphorylation of PLC γ , resulting in a sustained Ca^{2+} mobilization. The subsequent activation of NFAT regulates the expression of cytokine and chemokine genes (28, 32). Furthermore, NGF-induced outgrowth of embryonic axons requires NFAT activation (29). Whether NGF stimulates NFAT activation in mast cells and what biological response it induces is not known.

In addition to NGF, the complement component C3a plays an important role in the pathogenesis of allergic and inflammatory diseases (33–35). In leukocytes, C3a mediates its biological effects via the activation of a cell surface G protein-coupled receptor (GPCR) (18, 36). Cross-regulation of receptor tyrosine kinases by GPCRs has been studied quite extensively. In most cases, GPCR

Department of Pathology, School of Dental Medicine, University of Pennsylvania, Philadelphia, PA 19104

Received for publication October 20, 2003. Accepted for publication March 1, 2004.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by National Institutes of Health Grant HL63372.

² Current address: The Scripps Research Institute, La Jolla, CA 92037.

³ Address correspondence and reprint requests to Dr. Hydar Ali, Department of Pathology, School of Dental Medicine, University of Pennsylvania, 240 South 40th Street, Philadelphia, PA 19014-6002. E-mail address: ali@path.dental.upenn.edu

⁴ Abbreviations used in this paper: NGF, nerve growth factor; CsA, cyclosporin A; ERK, extracellular signal-regulated kinase; GAIP, G α -interacting protein; GIPC, GAIP-interacting protein C terminus; GPCR, G protein-coupled receptor; HMC, human mast cell; MAP, mitogen-activated protein; MCP-1, monocyte chemoattractant protein-1; MIP-1 β , macrophage inflammatory protein-1 β ; PDZ, PSD-95/Dig/ZO-1; PLC γ , phospholipase C γ ; PTX, pertussis toxin; PC12, pheochromocytoma 12.

activation leads to phosphorylation of receptor tyrosine kinases, resulting in their activation (37). Indeed, in pheochromocytoma 12 (PC12) cells, adenosine and neuropeptides use their respective cell surface GPCRs to cause tyrosine phosphorylation of TrkA, resulting in the activation of cell survival signaling pathway (38–40). Whether C3a also cross-regulates TrkA-mediated responses in mast cells has not been determined. To address this issue, we have used a human mast cell line (HMC-1), that natively expresses both C3aR and TrkA (18, 20, 41). In this study, we show that NGF stimulates chemokine production in HMC-1 cells via TrkA phosphorylation, sustained Ca^{2+} mobilization, and NFAT activation. Surprisingly, we found that C3a enhanced NGF-induced Ca^{2+} mobilization, NFAT activation, and chemokine production without causing TrkA phosphorylation.

Materials and Methods

Materials

NGF (2.5S) was obtained from Upstate Biotechnology (Lake Placid, NY). Purified C3a was obtained from Advance Research Technologies (San Diego, CA). Recombinant C5a was purchased from Sigma-Aldrich (St. Louis, MO). U0126, K252a, PMA, Ca^{2+} -ionophore A23187, cyclosporin A (CsA), and recombinant *Escherichia coli* LPS were purchased from Calbiochem (La Jolla, CA). Pertussis toxin (PTX) and all tissue culture reagents were purchased from Invitrogen (Gaithersburg, MD). Indo-1 AM and pluronic F-127 were from Molecular Probes (Eugene, OR). ECL Western blotting analysis kits were purchased from Amersham (Arlington, IL). Dual Luciferase Assay kit was obtained from Promega (Madison, WI). TrkA, phospho-TrkA, and extracellular signal-regulated kinase (ERK) Abs were purchased from Cell Signaling Technology (Beverly, MA). Capture Abs to macrophage inflammatory protein-1 β (MIP-1 β ; MAB 271), monocyte chemoattractant protein-1 (MCP-1; MAB 679), and IL-8 (DY 208), and polyclonal biotinylated Abs to MIP-1 β , MCP-1, and IL-8 were purchased from R&D Systems (Minneapolis, MN).

Calcium measurement

Ca^{2+} mobilization was determined as described previously (18, 42). Briefly, HMC-1 cells (1×10^6) were loaded with $1 \mu\text{M}$ Indo-1 AM in the presence of $1 \mu\text{M}$ pluronic F-127 for 30 min at room temperature. Cells were washed and resuspended in 1.5 ml of HEPES-buffered saline. Ca^{2+} mobilization was measured in a Hitachi F-2500 spectrophotometer (San Jose, CA) with an excitation wavelength of 355 nm and an emission wavelength of 410 nm (18).

TrkA phosphorylation

HMC-1 cells (5×10^6) were stimulated with C3a (10 nM), NGF (30 ng/ml), and C3a plus NGF at 37°C for 5 min. Reaction was stopped by the

addition of 3 vol of ice-cold PBS containing 1 mM sodium orthovanadate. Cells were lysed with 1 ml of lysis buffer (150 mM NaCl, 50 mM Tris, pH 8.0, 1% Nonidet P-40, 0.5% deoxycholate, 0.1% SDS, 5 mM EDTA, 10 mM sodium fluoride, 10 mM sodium pyrophosphate, 10 $\mu\text{g}/\text{ml}$ leupeptin, 1 $\mu\text{g}/\text{ml}$ pepstatin, 10 $\mu\text{g}/\text{ml}$ PMSF) for 1 h at 4°C, and centrifuged at $14,000 \times g$ for 10 min; supernatants were transferred to 1.5-ml tube and incubated with 2 $\mu\text{g}/\text{ml}$ TrkA Ab for 1 h on a rotating wheel, followed by incubation with 20 μl of protein G agarose beads to immunoprecipitate TrkA receptor. SDS-sample buffer (50 μl) was added to the pellet and heated to 90°C for 10 min. Proteins were separated on a 7.5% SDS-polyacrylamide gel and Western blotting was performed with phosphotyrosine-specific Ab, and TrkA phosphorylation was detected by ECL.

Transient transfection of HMC-1 cells and NFAT/NF- κ B luciferase activity

HMC-1 cells (2×10^7) were mixed with 10 μg of pNFAT or pNF- κ B luciferase reporter construct (Stratagene, La Jolla, CA) in a total volume of 200 μl of transfection medium. To standardize the expression level of the transfected constructs, 0.5 μg of pRL-SV40 *Renilla* luciferase construct (Promega) was added to the transfection mixture. Cells were then electroporated with a single pulse using a Bio-Rad gene pulser (Hercules, CA) as described previously (43). Cells (0.2×10^6) were seeded to each well of a 48-well plate. The following day, cells were incubated in serum-free medium and stimulated with C3a, NGF, or C3a plus NGF for 6 h at 37°C, unless otherwise indicated. More than 80% of the cells were viable at the time of the assay as determined by trypan blue staining. Cells were lysed and assayed for luciferase activity with Dual-Luciferase Reporter Assay System (Promega) according to manufacturer's instructions. The *Firefly* luciferase activity was normalized on the basis of *Renilla* luciferase activity.

Phosphorylation of ERK

Cells were stimulated with C3a, NGF, or C3a plus NGF for 5 min and the reactions were stopped by the addition of a 3-fold excess ice-cold PBS containing 1 mM sodium orthovanadate. The cells were lysed with $2 \times$ SDS sample buffer, and ERK phosphorylation was determined by Western blotting with a phospho-ERK1/2 Ab. The membrane was stripped and re-probed with an Ab that reacts with unphosphorylated ERK1/2, as described previously (18).

Assay of chemokine protein production by ELISA

HMC-1 cells (0.2×10^6 /well) were cultured in complete growth medium overnight. Cells were serum starved and stimulated with C3a, NGF, or C3a plus NGF for 6 h, unless otherwise stated. Chemokine protein levels (MIP-1 β , MCP-1, and IL-8) were quantified by sandwich ELISA using matched Ab pairs as described previously (18).





FIGURE 1. C3a- and NGF-induced Ca^{2+} mobilization in HMC-1 cells. Cells were cultured overnight in normal medium (Control; A and B), medium supplemented with PTX (100 ng/ml; C and D), or TrkA-specific Ab (2 $\mu\text{g}/\text{ml}$; E and F). Cells were then washed and loaded with Indo-1 AM, stimulated with C3a (10 nM; A, C, and E) or NGF (30 ng/ml; B, D, and F), and intracellular Ca^{2+} mobilization was determined. The data shown are representative of three similar experiments.

FIGURE 2. C3a does not cause TrkA phosphorylation but enhances NGF-induced NFAT activation and ERK phosphorylation. *A*, HMC-1 cells were exposed to buffer (Con), C3a (10 nM), NGF (30 ng/ml), or C3a + NGF for 5 min. Cell lysates were immunoprecipitated with anti-TrkA Ab, and receptor phosphorylation was determined by Western blotting with a phosphotyrosine-specific Ab (p-TrkA). *B*, HMC-1 cells were cotransfected with NFAT-dependent luciferase reporter and *Renilla* luciferase plasmids. Cells were stimulated for 6 h with C3a, NGF, or C3a + NGF. The NFAT luciferase activity was measured from cell lysate. Data presented are relative luciferase activity, normalized to *Renilla* luciferase activity as relative light units (RLU) ($n = 3$). *C*, Cells were stimulated with C3a, NGF, or C3a + NGF for 5 min. ERK activation was determined by Western blotting with phospho-specific ERK1/2 Ab (p-ERK1/2). *D*, Bar graph represents summary of ERK phosphorylation from four independent experiments, where C3a + NGF stimulated cells (100%). The data shown in *B* are mean \pm SEM of five experiments. *, $p < 0.05$ in the absence or presence of NGF and **, $p < 0.01$ NGF in the absence and presence of C3a.



Results

Regulation of C3aR and TrkA in HMC-1 cells

We initially used a Ca^{2+} mobilization assay to characterize endogenous C3aR and TrkA in HMC-1. As shown in Fig. 1A, C3a (10 nM) stimulated a rapid and transient Ca^{2+} spike that returned to basal within 2 min. In contrast, NGF (30 ng/ml) induced a slower response that remained elevated for a longer time period (Fig. 1B). To test the role of G protein signaling on C3a and NGF responses, cells were treated with PTX (100 ng/ml, overnight) and Ca^{2+} mobilization was determined. PTX caused a substantial inhibition of C3a-induced Ca^{2+} mobilization (Fig. 1C) but had no effect on NGF response (Fig. 1D). In contrast, anti-TrkA Ab did not inhibit C3a-induced Ca^{2+} mobilization but completely blocked the response to NGF (Fig. 1, E and F). The inhibitory effect of anti-TrkA Ab was specific for TrkA, as an isotype-matched Ab had no effect on the NGF response (data not shown). These data suggest that C3aR and TrkA use G protein and tyrosine kinase signaling pathways, respectively, to induce Ca^{2+} mobilization in HMC-1 cells.

C3a does not cause TrkA phosphorylation but enhances NGF-induced NFAT activation and ERK phosphorylation

Lee and Chao (38) recently showed that adenosine receptor activation resulted in Ca^{2+} mobilization in PC12 cells. Furthermore, adenosine caused transphosphorylation of TrkA via a Ca^{2+} -dependent pathway. Given that C3a stimulated Ca^{2+} mobilization in HMC-1 cells (Fig. 1), we hypothesized that C3a could also cause TrkA phosphorylation in HMC-1 cells. To test this hypothesis, cells were stimulated with C3a for 5 min and cell lysate was immunoprecipitated with anti-TrkA Ab. Tyrosine phosphorylation of TrkA was determined by Western blotting using a phosphotyrosine-specific Ab. As shown in Fig. 2A, C3a did not cause phosphorylation of TrkA in HMC-1 cells. TrkA phosphorylation was not evident even 1 h after C3a stimulation (data not shown). In contrast, NGF (30 ng/ml, 5 min) induced a robust tyrosine phosphorylation of TrkA. To determine whether C3a could enhance NGF-induced response, we stimulated HMC-1 cells simultaneously with C3a and NGF. We found that C3a had little or no effect on NGF-induced TrkA phosphorylation (Fig. 2A).





FIGURE 3. Effects of PTX on C3a- and NGF-induced NFAT activation and ERK phosphorylation. *A*, HMC-1 cells transiently expressing NFAT luciferase and *Renilla* constructs were incubated with buffer or PTX (100 ng/ml, overnight). Cells were stimulated with C3a (10 nM), NGF (30 ng/ml), or C3a + NGF for 6 h and NFAT luciferase activity (normalized to *Renilla* activity as relative light units (RLU)) was measured from the cell lysate. *B*, PTX-treated or untreated HMC-1 cells were stimulated with C3a, NGF, or C3a + NGF for 5 min and ERK1/2 activation was determined by Western blotting using phospho-ERK1/2 Ab. Bands were quantified by densitometry and expressed as percentage vs C3a plus NGF-stimulated cells (100%). The data shown are mean \pm SEM of four experiments. *, $p < 0.05$ and **, $p < 0.01$ in the absence and presence of PTX treatment.

FIGURE 4. C3a enhances NGF-induced MIP-1 β production in HMC-1 cells. *A*, Cells were stimulated with different concentrations of NGF for 6 h in the absence and presence of C3a (10 nM). MIP-1 β production was determined by sandwich ELISA. *B*, HMC-1 cells were preincubated with buffer or PTX (100 ng/ml, overnight), TrkA Ab (2 μ g/ml, overnight), or K252a (200 nM, 60 min). Cells were then exposed to C3a (10 nM), NGF (30 ng/ml), or C3a + NGF for 6 h, and MIP-1 β production was determined. The data shown are mean \pm SEM of three experiments. **, $p < 0.01$ and ***, $p < 0.001$ in the absence and presence of C3a.



Using an EMSA, we have recently shown that a sustained Ca²⁺ mobilization is associated with the activation of transcription factor NFAT in HMC-1 cells (18). As NGF stimulated a sustained Ca²⁺ mobilization (Fig. 1), we hypothesized that NGF could stimulate NFAT activation. To test this possibility, we generated transient transfectants in HMC-1 cells expressing NFAT linked to a luciferase reporter construct and determined the effects of NGF on NFAT-luciferase reporter activity. As shown in Fig. 2*B*, NGF caused a significant increase in NFAT-luciferase reporter activity. In agreement with our previous studies using EMSA (18), we found that C3a did not stimulate NFAT reporter activity. Interestingly, C3a caused significant enhancement of the response to NGF (Fig. 2*B*).

NGF has been shown to activate ERK phosphorylation in HMC-1 cells (20). We therefore sought to determine whether C3a could also enhance NGF-induced ERK phosphorylation. For these experiments, cells were stimulated with C3a, NGF, or C3a plus NGF for 5 min, and ERK phosphorylation was determined. As shown in Fig. 2, *C* and *D*, C3a and NGF caused ERK phosphorylation in HMC-1 cells. However, the response to C3a was \sim 50% greater than that of NGF. Interestingly, when C3a and NGF were added together, there was a synergistic activation of ERK phosphorylation. These findings demonstrate that although C3a did not stimulate TrkA phos-

phorylation, it enhanced NGF-induced downstream signaling such as NFAT activation and ERK phosphorylation.

In PC12 cells, NGF-induced ERK phosphorylation is mediated, at least in part, by a G_{αi/o} family of G proteins (44, 45). To determine the role of G proteins on NGF-induced signaling in HMC-1 cells, we tested the effects of PTX on C3a- and NGF-induced responses. PTX had no effects on NGF-induced NFAT activation (Fig. 3*A*) or ERK phosphorylation (Fig. 3*B*). However, PTX inhibited C3a-induced ERK phosphorylation and priming of NFAT activation and ERK phosphorylation by C3a (Fig. 3, *A* and *B*).

C3a enhances NGF-induced MIP-1 β and MCP-1 production in HMC-1 cells

Our next goal was to determine the functional consequence of the cross-regulation between C3aR and TrkA for NFAT activation and ERK phosphorylation in HMC-1 cells. We have previously shown that MIP-1 β and MCP-1 are up-regulated both at the levels of mRNA and protein by a variety of stimulants in HMC-1 cells (18). We decided to initially examine the effects of NGF and C3a plus NGF on MIP-1 β production. As shown in Fig. 4*A*, NGF caused a dose-dependent increase in MIP-1 β production in HMC-1 cells. The EC₅₀ value for NGF was \sim 10 ng/ml and the maximal response was obtained at a concentration of \sim 30 ng/ml. We have





FIGURE 5. C3a but not C5a or LPS enhances NGF-induced Ca²⁺ mobilization, NFAT activation, and MCP-1 production in HMC-1 cells. *A*, Cells were loaded with Indo-1 AM and the effects of C3a (10 nM), C5a (10 nM), and LPS (100 ng/ml) on Ca²⁺ mobilization were determined. *B*, Cells were stimulated with NGF, C3a + NGF, or C5a + NGF, and Ca²⁺ mobilization was determined. *C*, Transient transfectants were generated in HMC-1 cells coexpressing NFAT luciferase and *Renilla* plasmids. Cells were then stimulated with C3a (10 nM), C5a (10 nM), or LPS (100 ng/ml) in the absence and presence of NGF (30 ng/ml). NFAT luciferase activity was determined in cell lysate. The data are expressed as relative NFAT luciferase activity. *D*, MCP-1 production was measured by ELISA from culture supernatants at 6 h after stimulation. The data shown are mean \pm SEM of three to four experiments performed in triplicate. **, $p < 0.01$ and ***, $p < 0.001$ in the absence and presence of C3a, C5a, or LPS.

FIGURE 6. NGF does not cause NF- κ B activation or IL-8 production in HMC-1 cells. Transient transfectants were generated in HMC-1 cells coexpressing NF- κ B luciferase and *Renilla* plasmids. Cells were then stimulated with C3a (10 nM), C5a (10 nM), or LPS (100 ng/ml) in the absence and presence of NGF (30 ng/ml). Cells were also stimulated with PMA (100 nM) + A23187 (100 nM). *A*, Relative NF- κ B luciferase activity was determined in cell lysate. The data are expressed as fold increase over control. *B*, IL-8 production was measured by ELISA from culture supernatants at 6 h after stimulation. The data shown are mean \pm SEM of three to four experiments performed in triplicate. ***, $p < 0.001$ in the absence and presence of PMA + A23187.



previously shown that transient Ca^{2+} mobilization induced by C3a did not provide sufficient signal for NFAT activation or MIP-1 β production in HMC-1 cells (18). However, as shown in Fig. 4A, C3a (10 nM) enhanced NGF-induced MIP-1 β production by 2- to 3-fold. Chemokine production in response to C3a/NGF required protein synthesis, as treatment of cells with cycloheximide completely blocked the response (data not shown).

To determine the role of the G_i family of G proteins on chemokine production, cells were treated with PTX and its effects on NGF and C3a plus NGF-induced MIP-1 β production were determined. PTX selectively blocked the priming by C3a but had no effect on the response to NGF (Fig. 4B). In contrast, a TrkA-specific Ab or an inhibitor for TrkA signaling, K252a, blocked NGF-induced MIP-1 β production both in the absence and presence of C3a. These data suggest that NGF-induced MIP-1 β production is mediated via the activation of TrkA, whereas the enhanced response by C3a requires the activation of PTX-sensitive G protein.

To determine whether C3a selectively enhances NGF-induced responses in HMC-1 cells, we tested the effects of C5a and LPS on NGF-induced responses in HMC-1 cells. We found that C5a caused a transient Ca^{2+} mobilization in HMC-1 cells whereas LPS had no effect (Fig. 5A). Interestingly, only C3a but not C5a or LPS enhanced NGF-induced Ca^{2+} mobilization (Fig. 5B), NFAT activation (Fig. 5C), MCP-1 production (Fig. 5D), and MIP-1 β (data not shown) in HMC-1 cells.

Effects of C3a/NGF on MIP-1 β and MCP-1 production in HMC-1 cells does not require NF- κ B activation

NF- κ B is a crucial transcription factor that regulates the expression of many proinflammatory cytokines and immunoregulatory molecules (46–48). Although NGF causes NFAT activation in neuronal tissues (29, 49) and HMC-1 cells (Fig. 2B), whether it also stimulates NF- κ B activation has not been determined. We therefore generated transient transfectants in HMC-1 cells expressing NF- κ B luciferase construct and tested the effect of NGF on NF- κ B reporter activity. As shown in Fig. 6A, NGF alone or in combination with C3a, C5a, and LPS did not stimulate NF- κ B luciferase activity. In contrast, PMA plus Ca^{2+} ionophore A23187 resulted in \sim 5-fold increase in NF- κ B reporter activity (Fig. 6A). NF- κ B activation is required for chemokine IL-8 production in a variety of cell types (50–53). As shown in Fig. 6B, NGF alone or in combination with C3a, C5a, or LPS did not stimulate IL-8 production. In contrast, PMA/A23187 caused a substantial generation of this chemokine in HMC-1 cells. These data suggest that C3a selectively enhances NGF-induced MIP-1 β and MCP-1 production in HMC-1 cells via the activation of NFAT but not NF- κ B.

Effects of ERK phosphorylation and NFAT activation on MIP-1 β production in HMC-1 cells

As C3a enhanced NGF-induced ERK phosphorylation in HMC-1 cells (Fig. 2), we tested the role of this pathway on the cross-regulation of MIP-1 β production. U0126 (MAP/ERK-1/2 inhibitor) completely blocked NGF-induced ERK phosphorylation both in the absence and presence of C3a (Fig. 7A). However, it inhibited NGF-induced chemokine production by $38 \pm 1.04\%$ and blocked C3a- plus NGF-induced response by $31 \pm 12\%$ (Fig. 7B). These data suggest that ERK phosphorylation pathway plays a minor role in NGF-induced chemokine production and has little or no effect on C3a-induced priming.





FIGURE 7. Role of ERK phosphorylation on C3a- and NGF-induced MIP-1 β production in HMC-1 cells. *A*, Cells were preincubated with buffer or U0126 (10 μ M, 30 min) and stimulated with C3a (10 nM), NGF (30 ng/ml), or C3a + NGF for 5 min, and ERK phosphorylation was determined by Western blotting using phospho-ERK1/2 Ab (p-ERK1/2, upper panel). The membrane was stripped and reprobed with ERK1/2 Ab (lower panel). *B*, Cells were preincubated with buffer or U0126 (10 μ M, 30 min) and stimulated with C3a, NGF, or C3a + NGF for 6 h, and MIP-1 β production was determined. Data presented are mean \pm SEM of five experiments. *, $p < 0.05$ in the absence and presence of U0126.

FIGURE 8. NFAT activation is required for C3a- and NGF-induced MIP-1 β production in HMC-1 cells. Transient transfectants were generated in HMC-1 cells coexpressing NFAT luciferase and *Renilla* plasmids. Cells were then preincubated with or without CsA, (100 nM, 30 min) and stimulated with C3a (10 nM), NGF (30 ng/ml), or C3a + NGF. NFAT luciferase activity from the lysate (A) and MIP-1 β production (B) was measured from culture supernatants 6 h after stimulation. C, Cells were preincubated with CsA and stimulated with C3a, NGF, or C3a + NGF for 5 min, and ERK phosphorylation was determined by Western blotting using a phospho-ERK specific Ab (p-ERK1/2, upper panel). Protein loading was confirmed by reprobating the membrane with ERK1/2 Ab (lower panel).



Given that C3a enhanced NGF-induced NFAT activation (Fig. 2 and 3A), we hypothesized that the ability of C3a to prime NGF-induced MIP-1 β production could be mediated via the activation of NFAT. To test this possibility, we used CsA, a potent inhibitor of calcineurin-mediated NFAT activation. CsA (100 nM) completely inhibited both NGF-induced NFAT activation (Fig. 8A) and MIP-1 β production (Fig. 8B) in the presence and absence of C3a. The inhibitory effects of CsA were specific for NFAT, as it had no effect on C3a-, NGF-, or C3a- plus NGF-induced ERK phosphorylation (Fig. 8C).

Discussion

Although NGF exerts its biological effects predominantly on neurons, it can also influence the development and activation of many hemopoietic cell types. The mast cell was the first nonneuronal cell type identified as a target for NGF (8). In addition to TrkA, mast cells express GPCRs for C3a (18, 36). Given that both C3aR and TrkA play important roles in allergic and inflammatory diseases, we hypothesized that these receptors could cross-regulate each other to enhance signaling and cell activation. In the present study, we used HMC-1 that natively expresses both C3aR and TrkA, and demonstrated a novel cross-talk between these receptors for Ca²⁺ mobilization, NFAT activation, and chemokine MIP-1 β production. Interestingly, this type of cross-regulation is specific for C3aR and TrkA receptors as C5a, which mediates many biological responses similar to C3a, failed to enhance NGF-induced responses in HMC-1 cells.

The biological effects of NGF are mediated via two types of cell surface receptors, p75 and TrkA (19). Previous studies have shown that mast cells express TrkA but not p75 (11, 20, 21). In the present study, we demonstrated that anti-TrkA Ab and a specific Trk antagonist, K252a, blocked NGF-induced Ca²⁺ mobilization and MIP-1 β production. These data are consistent with the notion that the biological effects of NGF on mast cells are mediated via the activation of TrkA. Despite extensive reports on NGF-induced responses in mast cells, very little information is available on TrkA-mediated signaling pathways in these cells. In the present study, we have shown that NGF caused ERK phosphorylation in HMC-1 cells. However, U0126, which completely blocked NGF-induced

ERK phosphorylation, had little effect on MIP-1 β production. These data suggest that NGF-induced chemokine production in HMC-1 cells requires the activation of additional signaling pathways.

The data presented herein demonstrate that Ca²⁺ mobilization and the subsequent activation of the transcription factor NFAT provides an important mechanism for NGF-induced chemokine production in HMC-1 cells. NFAT is a cytosolic transcription factor that was originally shown to regulate the expression of cytokine genes in Ag-stimulated mast cells and T cells (28, 54). The mechanism of NFAT activation involves a Ca²⁺-dependent activation of the phosphatase calcineurin, which dephosphorylates NFAT resulting in its translocation to the nucleus. The immunosuppressive drug CsA inhibits NFAT activation by blocking calcineurin activation. We have recently shown fMLP, via the activation of a cell surface GPCR, stimulated a sustained Ca²⁺ mobilization and chemokine MIP-1 β production in HMC-1 cells (18). We also demonstrated that the ability of fMLP to induce chemokine production was dependent on NFAT activation. The findings in the present study that NGF also stimulated a sustained Ca²⁺ mobilization and NFAT luciferase activity suggest that this transcription factor could also be involved in NGF-induced MIP-1 β production. This contention is supported by the observations that both NGF-induced NFAT activation and chemokine production were inhibited by CsA.

A novel finding of the present study was that C3a, which caused a transient Ca²⁺ mobilization, did not stimulate MIP-1 β production by itself but significantly enhanced the response to NGF in HMC-1 cells. The studies on the cross-regulation of C3aR and TrkA provided additional support for the notion that chemokine production in HMC-1 cells depends mainly on NFAT activation. This contention is supported by the following observations. First, although C3a enhanced both ERK phosphorylation and NFAT activation, blocking ERK phosphorylation had no effect on C3a-induced priming of chemokine production. Second, CsA blocked both NGF-induced NFAT activation and chemokine production, but not ERK phosphorylation in the absence or presence of C3a.

The molecular mechanism by which C3a enhances NGF-induced NFAT activation and MIP-1 β production is not known. In PC12 cells, adenosine, lysophosphatidic acid, and neuropeptides

use their respective GPCRs to cause transactivation of TrkA (38–40, 55). We therefore hypothesized that the ability of C3a to enhance NGF-induced NFAT activation and chemokine production in mast cells could involve the tyrosine phosphorylation of TrkA. However, this possibility is unlikely, as C3a did not cause TrkA phosphorylation in HMC-1 cells. Furthermore, although NGF caused tyrosine phosphorylation of TrkA, C3a had no effect on this response. These findings suggest that the locus for priming by C3a is downstream of TrkA phosphorylation.

An interesting finding of the present study was that although C3a and C5a activate HMC-1 cells via their respective GPCRs, only C3a enhanced NGF-induced signaling and chemokine production. These findings raise the intriguing possibility that C3a could regulate TrkA signaling via the formation of a signaling complex downstream of TrkA phosphorylation. G_{α} -interacting protein (GAIP)-interacting protein C terminus (GIPC) is a PSD-95/Dig/ZO-1 (PDZ)-binding protein that is linked to GPCR signaling pathway via its interaction with GAIP, a regulator of G protein signaling. It has recently been shown that GIPC binds to both phosphorylated TrkA and GAIP to form a macromolecular signaling complex in PC12 cells (56). Hu et al. (57) recently showed that β_1 -adrenergic receptor interacts with GIPC and that this interaction requires the presence of serine residue in the PDZ motif T/SXV. It is interesting to note that only C3aR but not C5aR (CD88) possesses the PDZ motif T/SXV at its carboxyl terminus. Thus, the possibility that C3aR interacts with GIPC or other adapter molecules (58, 59) and that this interaction leads to the formation of a macromolecular signaling complex with phosphorylated TrkA resulting in enhanced Ca^{2+} mobilization, NFAT activation, and chemokine production remains to be determined.

Cross-regulation between GPCRs and receptor tyrosine kinases has been studied quite extensively. It has been shown that epidermal growth factor, platelet-derived growth factor, and insulin-like growth factor receptors become tyrosine phosphorylated after GPCR activation leading to MAP kinase activation (37, 60–64). Furthermore, adenosine and neuropeptides, via their respective GPCRs, cause transphosphorylation of TrkA resulting in the enhancement of both Akt phosphorylation and cell survival (39, 40). In the present study, we described a novel finding that C3a enhances NGF-induced transcription factor NFAT activation and chemokine production via a pathway that does not require transphosphorylation of a receptor tyrosine kinase, TrkA. Although HMC-1 is an immature mast cell line and does not display all properties of primary mast cells, it is widely used as a model to study mast cell signaling and biology in vitro (65–73). It is important to note that human eosinophils, which express both C3a and TrkA, respond to NGF for cytokine production via a pathway that depends on NFAT activation (74, 75). Furthermore, NGF-induced outgrowth of embryonic axons requires NFAT activation (29). A large number of other immune and neuronal cell types that express C3aR and other GPCRs also possess TrkA and NFAT (29, 74, 76, 77). Thus, the type of cross-regulation described in the present study between C3aR and TrkA in HMC-1 cells is likely to be a general phenomenon.

Acknowledgments

We are grateful to Dr. Joseph Butterfield (Mayo Clinic, Rochester, MN) for supplying us with HMC-1 cells.

References

- Bothwell, M. 1995. Functional interactions of neurotrophins and neurotrophin receptors. *Annu. Rev. Neurosci.* 18:223.
- Ip, N. Y., and G. D. Yancopoulos. 1996. The neurotrophins and CNTF: two families of collaborative neurotrophic factors. *Annu. Rev. Neurosci.* 19:491.
- Levi-Montalcini, R. 1987. The nerve growth factor 35 years later. *Science* 237:1154.
- Xiang, Z., and G. Nilsson. 2000. IgE receptor-mediated release of nerve growth factor by mast cells. *Clin. Exp. Allergy* 30:1379.
- Bonini, S., A. Lambiase, F. Angelucci, L. Magrini, L. Manni, and L. Aloe. 1996. Circulating nerve growth factor levels are increased in humans with allergic diseases and asthma. *Proc. Natl. Acad. Sci. USA* 93:10955.
- Sanico, A. M., A. M. Stanisz, T. D. Gleeson, S. Bora, D. Proud, J. Bienenstock, V. E. Koliatsos, and A. Togias. 2000. Nerve growth factor expression and release in allergic inflammatory disease of the upper airways. *Am. J. Respir. Crit. Care Med.* 161:1631.
- Virchow, J. C., P. Julius, M. Lommatzsch, W. Luttmann, H. Renz, and A. Braun. 1998. Neurotrophins are increased in bronchoalveolar lavage fluid after segmental allergen provocation. *Am. J. Respir. Crit. Care Med.* 158:2002.
- Aloe, L. 1988. The effect of nerve growth factor and its antibody on mast cells in vivo. *J. Neuroimmunol.* 18:1.
- Sawada, J., A. Itakura, A. Tanaka, T. Furusaka, and H. Matsuda. 2000. Nerve growth factor functions as a chemoattractant for mast cells through both mitogen-activated protein kinase and phosphatidylinositol 3-kinase signaling pathways. *Blood* 95:2052.
- Pearce, F. L., and H. L. Thompson. 1986. Some characteristics of histamine secretion from rat peritoneal mast cells stimulated with nerve growth factor. *J. Physiol.* 372:379.
- Horigome, K., J. C. Pryor, E. D. Bullock, and E. M. Johnson, Jr. 1993. Mediator release from mast cells by nerve growth factor. Neurotrophin specificity and receptor mediation. *J. Biol. Chem.* 268:14881.
- Kawamoto, K., J. Aoki, A. Tanaka, A. Itakura, H. Hosono, H. Arai, Y. Kiso, and H. Matsuda. 2002. Nerve growth factor activates mast cells through the collaborative interaction with lysophosphatidylserine expressed on the membrane surface of activated platelets. *J. Immunol.* 168:6412.
- Marshall, J. S., K. Gomi, M. G. Blennerhassett, and J. Bienenstock. 1999. Nerve growth factor modifies the expression of inflammatory cytokines by mast cells via a prostanoic-dependent mechanism. *J. Immunol.* 162:4271.
- Alam, R. 1997. Chemokines in allergic inflammation. *J. Allergy Clin. Immunol.* 99:273.
- Teran, L. M., and D. E. Davies. 1996. The chemokines: their potential role in allergic inflammation. *Clin. Exp. Allergy* 26:1005.
- Yano, K., M. Yamaguchi, F. Demora, C. S. Lantz, J. H. Butterfield, J. J. Costa, and S. J. Galli. 1997. Production of macrophage inflammatory protein-1 α by human mast cells: increased anti-IgE-dependent secretion after IgE-dependent enhancement of mast cell IgE-binding ability. *Lab. Invest.* 77:185.
- Selvan, R. S., J. H. Butterfield, and M. S. Krangel. 1994. Expression of multiple chemokine genes by a human mast cell leukemia. *J. Biol. Chem.* 269:13893.
- Ali, H., J. Ahmed, C. Hernandez-Munain, J. L. Baron, M. S. Krangel, and D. D. Patel. 2000. Chemokine production by G protein-coupled receptor activation in a human mast cell line: roles of extracellular signal-regulated kinase and NFAT. *J. Immunol.* 165:7215.
- Sutter, A., R. J. Riopelle, R. M. Harris-Warrick, and E. M. Shooter. 1979. Nerve growth factor receptors: characterization of two distinct classes of binding sites on chick embryo sensory ganglia cells. *J. Biol. Chem.* 254:5972.
- Tam, S. Y., M. Tsai, M. Yamaguchi, K. Yano, J. H. Butterfield, and S. J. Galli. 1997. Expression of functional TrkA receptor tyrosine kinase in the HMC-1 human mast cell line and in human mast cells. *Blood* 90:1807.
- Burgi, B., U. H. Otten, B. Ochensberger, S. Rihs, K. Heese, P. B. Ehrhard, C. F. Ibanez, and C. A. Dahinden. 1996. Basophil priming by neurotrophic factors: activation through the trk receptor. *J. Immunol.* 157:5582.
- Obermeier, A., R. Lammers, K. H. Wiesmuller, G. Jung, J. Schlessinger, and A. Ullrich. 1993. Identification of Trk binding sites for SHC and phosphatidylinositol 3'-kinase and formation of a multimeric signaling complex. *J. Biol. Chem.* 268:22963.
- Stephens, R. M., D. M. Loeb, T. D. Copeland, T. Pawson, L. A. Greene, and D. R. Kaplan. 1994. Trk receptors use redundant signal transduction pathways involving SHC and PLC- γ 1 to mediate NGF responses. *Neuron* 12:691.
- Obermeier, A., H. Halfter, K. H. Wiesmuller, G. Jung, J. Schlessinger, and A. Ullrich. 1993. Tyrosine 785 is a major determinant of Trk-substrate interaction. *EMBO J.* 12:933.
- Bonni, A., A. Brunet, A. E. West, S. R. Datta, M. A. Takasu, and M. E. Greenberg. 1999. Cell survival promoted by the Ras-MAPK signaling pathway by transcription-dependent and -independent mechanisms. *Science* 286:1358.
- Watson, F. L., H. M. Heerssen, A. Bhattacharyya, L. Klesse, M. Z. Lin, and R. A. Segal. 2001. Neurotrophins use the Erk5 pathway to mediate a retrograde survival response. *Nat. Neurosci.* 4:981.
- Choi, D. Y., J. J. Toledo-Aral, R. Segal, and S. Halegoua. 2001. Sustained signaling by phospholipase C- γ mediates nerve growth factor-triggered gene expression. *Mol. Cell. Biol.* 21:2695.
- Rao, A., C. Luo, and P. G. Hogan. 1997. Transcription factors of the NFAT family: regulation and function. *Annu. Rev. Immunol.* 15:707.
- Graef, I. A., F. Wang, F. Charron, L. Chen, J. Neilson, M. Tessier-Lavigne, and G. R. Crabtree. 2003. Neurotrophins and netrins require calcineurin/NFAT signaling to stimulate outgrowth of embryonic axons. *Cell* 113:657.
- Hill-Eubanks, D. C., M. F. Gomez, A. S. Stevenson, and M. T. Nelson. 2003. NFAT regulation in smooth muscle. *Trends Cardiovasc. Med.* 13:56.
- Crabtree, G. R., and E. N. Olson. 2002. NFAT signaling: choreographing the social lives of cells. *Cell* 109(Suppl.):S67.
- Turner, H., and D. A. Cantrell. 1997. Distinct Ras effector pathways are involved in Fc ϵ R1 regulation of the transcriptional activity of Elk-1 and NFAT in mast cells. *J. Exp. Med.* 185:43.

33. Drouin, S. M., D. B. Corry, T. J. Hollman, J. Kildsgaard, and R. A. Wetsel. 2002. Absence of the complement anaphylatoxin C3a receptor suppresses Th2 effector functions in a murine model of pulmonary allergy. *J. Immunol.* 169:5926.
34. Humbles, A. A., B. Lu, C. A. Nilsson, C. Lilly, E. Israel, Y. Fujiwara, N. P. Gerard, and C. Gerard. 2000. A role for the C3a anaphylatoxin receptor in the effector phase of asthma. *Nature* 406:998.
35. Bautsch, W., H. G. Hoymann, Q. Zhang, I. Meier-Wiedenbach, U. Raschke, R. S. Ames, B. Sohns, N. Flemme, A. Meyer Zu Vilsendorf, M. Grove, et al. 2000. Cutting edge: guinea pigs with a natural C3a-receptor defect exhibit decreased bronchoconstriction in allergic airway disease: evidence for an involvement of the C3a anaphylatoxin in the pathogenesis of asthma. *J. Immunol.* 165:5401.
36. Nilsson, G., M. Johnell, C. H. Hammer, H. L. Tiffany, K. Nilsson, D. D. Metcalfe, A. Siegbahn, and P. M. Murphy. 1996. C3a and C5a are chemotaxins for human mast cells and act through distinct receptors via a pertussis toxin-sensitive signal transduction pathway. *J. Immunol.* 157:1693.
37. Luttrell, L. M., Y. Daaka, and R. J. Lefkowitz. 1999. Regulation of tyrosine kinase cascades by G-protein-coupled receptors. *Curr. Opin. Cell Biol.* 11:177.
38. Lee, F. S., and M. V. Chao. 2001. Activation of Trk neurotrophin receptors in the absence of neurotrophins. *Proc. Natl. Acad. Sci. USA* 98:3555.
39. Lee, F. S., R. Rajagopal, and M. V. Chao. 2002. Distinctive features of Trk neurotrophin receptor transactivation by G protein-coupled receptors. *Cytokine Growth Factor Rev.* 13:11.
40. Lee, F. S., R. Rajagopal, A. H. Kim, P. C. Chang, and M. V. Chao. 2002. Activation of Trk neurotrophin receptor signaling by pituitary adenylate cyclase-activating polypeptides. *J. Biol. Chem.* 277:9096.
41. Nilsson, G., K. Forsberg-Nilsson, Z. Xiang, F. Hallbook, K. Nilsson, and D. D. Metcalfe. 1997. Human mast cells express functional TrkA and are a source of nerve growth factor. *Eur. J. Immunol.* 27:2295.
42. Ali, H., R. M. Richardson, E. D. Tomhave, J. R. Didsbury, and R. Snyderman. 1993. Differences in phosphorylation of formylpeptide and C5a chemoattractant receptors correlate with differences in desensitization. *J. Biol. Chem.* 268:24247.
43. Ahamed, J., and H. Ali. 2002. Distinct roles of receptor phosphorylation, G protein usage, and mitogen-activated protein kinase activation on platelet activating factor-induced leukotriene C4 generation and chemokine production. *J. Biol. Chem.* 277:22685.
44. Knipper, M., A. Beck, J. Rylett, and H. Breer. 1993. Neurotrophin induced cAMP and IP3 responses in PC12 cells: different pathways. *FEBS Lett.* 324:147.
45. Rakhit, S., S. Pyne, and N. J. Pyne. 2001. Nerve growth factor stimulation of p42/p44 mitogen-activated protein kinase in PC12 cells: role of G_{i/o}, G protein-coupled receptor kinase 2, β -arrestin I, and endocytic processing. *Mol. Pharmacol.* 60:63.
46. Green, D. R. 2003. Death and NF- κ B in T cell activation: life at the edge. *Mol. Cell.* 11:551.
47. Chen, F., L. M. Demers, and X. Shi. 2002. Upstream signal transduction of NF- κ B activation. *Curr. Drug Targets Inflamm. Allergy.* 1:137.
48. Kravchenko, V. V., Z. Pan, J. Han, J. M. Herbert, R. J. Ulevitch, and R. D. Ye. 1995. Platelet-activating factor induces NF- κ B activation through a G protein-coupled pathway. *J. Biol. Chem.* 270:14928.
49. Groth, R. D., and P. G. Mermelstein. 2003. Brain-derived neurotrophic factor activation of NFAT (nuclear factor of activated T-cells)-dependent transcription: a role for the transcription factor NFATc4 in neurotrophin-mediated gene expression. *J. Neurosci.* 23:8125.
50. Hsu, M. H., M. Wang, D. D. Browning, N. Mukaida, and R. D. Ye. 1999. NF- κ B activation is required for C5a-induced interleukin-8 gene expression in mononuclear cells. *Blood* 93:3241.
51. Schwarz, M., and P. M. Murphy. 2001. Kaposi's sarcoma-associated herpesvirus G protein-coupled receptor constitutively activates NF- κ B and induces proinflammatory cytokine and chemokine production via a C-terminal signaling determinant. *J. Immunol.* 167:505.
52. Shepard, L. W., M. Yang, P. Xie, D. D. Browning, T. Voyno-Yasenetskaya, T. Kozasa, and R. D. Ye. 2001. Constitutive activation of NF- κ B and secretion of interleukin-8 induced by the G protein-coupled receptor of Kaposi's sarcoma-associated herpesvirus involve G_{α13} and RhoA. *J. Biol. Chem.* 276:45979.
53. Zhao, D., S. Kuhn-Moore, H. Zeng, J. S. Wu, M. P. Moyer, and C. Pothoulakis. 2003. Neurotensin stimulates IL-8 expression in human colonic epithelial cells through Rho GTPase-mediated NF- κ B pathways. *Am. J. Physiol.* 284:C1397.
54. Turner, H., M. Gomez, E. McKenzie, A. Kirchem, A. Lennard, and D. A. Cantrell. 1998. Rac-1 regulates nuclear factor of activated T cells (NFAT) C1 nuclear translocation in response to Fc ϵ receptor type 1 stimulation of mast cells. *J. Exp. Med.* 188:527.
55. Moughal, N. A., C. Waters, B. Sambhi, S. Pyne, and N. J. Pyne. 2004. Nerve growth factor signaling involves interaction between the Trk A receptor and lysophosphatidate receptor 1 systems: nuclear translocation of the lysophosphatidate receptor 1 and Trk A receptors in pheochromocytoma 12 cells. *Cell. Signal.* 16:127.
56. Lou, X., H. Yano, F. Lee, M. V. Chao, and M. G. Farquhar. 2001. GIPC and GAIP form a complex with TrkA: a putative link between G protein and receptor tyrosine kinase pathways. *Mol. Biol. Cell* 12:615.
57. Hu, L. A., W. Chen, N. P. Martin, E. J. Whalen, R. T. Premont, and R. J. Lefkowitz. 2003. GIPC interacts with the β_1 -adrenergic receptor and regulates β_1 -adrenergic receptor-mediated ERK activation. *J. Biol. Chem.* 278:26295.
58. Nakazawa, T., I. Nakano, M. Sato, T. Nakamura, M. Tamai, and N. Mori. 2002. Comparative expression profiles of Trk receptors and Shc-related phosphotyrosine adapters during retinal development: potential roles of N-Shc/ShcC in brain-derived neurotrophic factor signal transduction and modulation. *J. Neurosci. Res.* 68:668.
59. Nakamura, T., M. Komiya, K. Sone, E. Hirose, N. Gotoh, H. Morii, Y. Ohta, and N. Mori. 2002. Grit, a GTPase-activating protein for the Rho family, regulates neurite extension through association with the TrkA receptor and N-Shc and CrkL/Crk adapter molecules. *Mol. Cell. Biol.* 22:8721.
60. Daub, H., F. U. Weiss, C. Wallasch, and A. Ullrich. 1996. Role of transactivation of the EGF receptor in signalling by G-protein-coupled receptors. *Nature* 379:557.
61. Prenzel, N., E. Zwick, H. Daub, M. Leserer, R. Abraham, C. Wallasch, and A. Ullrich. 1999. EGF receptor transactivation by G-protein-coupled receptors requires metalloproteinase cleavage of proHB-EGF. *Nature* 402:884.
62. Linseman, D. A., C. W. Benjamin, and D. A. Jones. 1995. Convergence of angiotensin II and platelet-derived growth factor receptor signaling cascades in vascular smooth muscle cells. *J. Biol. Chem.* 270:12563.
63. Rao, G. N., P. Delafontaine, and M. S. Runge. 1995. Thrombin stimulates phosphorylation of insulin-like growth factor-1 receptor, insulin receptor substrate-1, and phospholipase C- γ 1 in rat aortic smooth muscle cells. *J. Biol. Chem.* 270:27871.
64. Ahmed, I., D. Gesty-Palmer, M. K. Drezner, and L. M. Luttrell. 2003. Transactivation of the epidermal growth factor receptor mediates parathyroid hormone and prostaglandin F2 α -stimulated mitogen-activated protein kinase activation in cultured transgenic murine osteoblasts. *Mol. Endocrinol.* 17:1607.
65. Li, L., Y. Yang, and R. L. Stevens. 2003. RasGRP4 regulates the expression of prostaglandin D₂ in human and rat mast cell lines. *J. Biol. Chem.* 278:4725.
66. Feoktistov, I., S. Ryzhov, A. E. Goldstein, and I. Biaggioni. 2003. Mast cell-mediated stimulation of angiogenesis: cooperative interaction between A2B and A3 adenosine receptors. *Circ. Res.* 92:485.
67. Forsythe, P., and A. D. Befus. 2003. Inhibition of calpain is a component of nitric oxide-induced down-regulation of human mast cell adhesion. *J. Immunol.* 170:287.
68. Hollander, C., M. Nystrom, S. Janciauskiene, and U. Westin. 2003. Human mast cells decrease SLPI levels in type II: like alveolar cell model, in vitro. *Cancer Cell Int.* 3:14.
69. Berger, P., P. O. Girodet, H. Begeret, O. Ousova, D. W. Perng, R. Marthan, A. F. Walls, and J. M. Tunon De Lara. 2003. Tryptase-stimulated human airway smooth muscle cells induce cytokine synthesis and mast cell chemotaxis. *FASEB J.* 17:2139.
70. Alexandrakis, M. G., D. S. Kyriakou, D. Seretakis, W. Boucher, R. Letourneau, D. Kempuraj, and T. C. Theoharides. 2003. Inhibitory effect of retinoic acid on proliferation, maturation and tryptase level in human leukemic mast cells (HMC-1). *Int. J. Immunopathol. Pharmacol.* 16:43.
71. Gronenberg, D. A., P. Welker, T. C. Fischer, Q. T. Dinh, A. Grutzkau, C. Peiser, U. Wahn, B. M. Henz, and A. Fischer. 2003. Down-regulation of vasoactive intestinal polypeptide receptor expression in atopic dermatitis. *J. Allergy Clin. Immunol.* 111:1099.
72. Gallagher, S. J., J. S. Marshall, and D. W. Hoskin. 2003. Human mast cells induce caspase-independent DNA fragmentation in leukemic T cells. *Oncol. Rep.* 10:1019.
73. Kang, O. H., H. J. Jeong, D. K. Kim, S. C. Choi, T. H. Kim, Y. H. Nah, H. M. Kim, and Y. M. Lee. 2003. Trypsin induces tumour necrosis factor- α secretion from a human leukemic mast cell line. *Cell Biochem. Funct.* 21:161.
74. Noga, O., C. Englmann, G. Hanf, A. Grutzkau, S. Guhl, and G. Kunkel. 2002. Activation of the specific neurotrophin receptors TrkA, TrkB and TrkC influences the function of eosinophils. *Clin. Exp. Allergy* 32:1348.
75. Cron, R. Q., S. J. Bort, Y. Wang, M. W. Brunvand, and D. B. Lewis. 1999. T cell priming enhances IL-4 gene expression by increasing nuclear factor of activated T cells. *J. Immunol.* 162:860.
76. Plyte, S., M. Boncrisiano, E. Fattori, F. Galvagni, S. R. Paccani, M. B. Majolini, S. Oliviero, G. Ciliberto, J. L. Telford, and C. T. Baldari. 2001. Identification and characterization of a novel nuclear factor of activated T-cells-1 isoform expressed in mouse brain. *J. Biol. Chem.* 276:14350.
77. Ricci, A., L. Felici, S. Mariotta, F. Mannino, G. Schmid, C. Terzano, G. Cardillo, F. Amenta, and E. Bronzetti. 2004. Neurotrophin and neurotrophin receptor protein expression in the human lung. *Am. J. Respir. Cell Mol. Biol.* 30:12.